

A Polynomial-Scale Gap Between the Chromatic and Cochromatic Numbers of a Random Graph

Candidate all-n solution to Erdős Problem 625; not externally peer reviewed

SAMUIL PETKOV

École normale supérieure–PSL (ENS–PSL)

samuil.petkov@ens.psl.eu

Preprint version of 12 July 2026

Abstract

For a graph G , let $\chi(G)$ be its chromatic number and let $\zeta(G)$ be the least number of parts in a vertex partition in which every part induces either an empty graph or a complete graph. We present a candidate proof that, for $G_n \sim G(n, 1/2)$,

$$\Pr \left\{ \chi(G_n) - \zeta(G_n) \geq \frac{(\ln 2)^2}{32} \ln \left(\frac{200}{153} \right) \frac{n}{(\ln n)^3} \right\} \longrightarrow 1. \quad (0.1)$$

The construction uses classes of the four sizes $\alpha - 2, \dots, \alpha - 5$, where α is the usual independence-number phase. Signing each class as independent or complete moves its first-moment root down by order $n/(\ln n)^3$, uniformly through the whole phase cycle. An exact sign-summed overlap identity reduces the second moment to a bipartite configuration model. We sum partial diagonals, unequal-type dense containments, all intermediate high cells, and every residual even-subgraph attachment with logarithmic cost $o(n/(\ln n)^4)$. Paley–Zygmund gives a possibly rare cocolouring, and a proved Alon–Scott-type leftover argument amplifies it to high probability without losing the root separation.

Introduction

The divergence question was posed by Erdős and Gimbel (1993, p. 263) and restated by Gimbel (2016, Section 7.4, p. 100); it is also recorded as Erdős Problem 625 on the problem website of Bloom (2026). For historical comparison, the first-order dense-random-graph chromatic asymptotic is due to Bollobás (1988, Theorem 4); Janson et al. (2000, Section 7.4, Theorem 7.14 and Remark 7.15) give a textbook treatment.

The recent work of Heckel (2024, Theorem 1) and, independently, Steiner (2024) gave the first quantitative evidence toward divergence. More precisely, Heckel proved that if a deterministic integer sequence $g(n)$ satisfies $\Pr\{\chi(G_n) - \zeta(G_n) \leq g(n)\} > 0.999$, then along an unbounded sequence n^* ,

$$g(n^*) > c \frac{\sqrt{n^*} \ln \ln n^*}{(\ln n^*)^3}.$$

This is an obstruction to a uniformly small high-probability upper bound; it is not itself a high-probability lower bound for the gap at every n . Heckel also conjectured the scale $\chi(G_n) - \zeta(G_n) = \Theta(n/(\ln n)^3)$ with high probability (Heckel, 2024, Conjecture 4).

The subsequent theorem of Heckel (2025, Theorem 1) proved that, for every fixed $\varepsilon > 0$,

$$n^{0.05+\varepsilon} \leq \mu_\alpha \leq n^{1-\varepsilon} \implies \chi(G_n) - \zeta(G_n) \geq n^{1-\varepsilon} \text{ with high probability.}$$

The phrase “roughly 95%” refers to the integers satisfying this phase-dependent window (with a fixed buffer), not to an all- n theorem; see Heckel (2025, Section 2). The relationship between the two Heckel papers and the present manuscript is as follows.

- (i) The anti-concentration theorem of Heckel (2024, Theorem 1) is contextual evidence and motivation; it is not a logical input to the proof below.
- (ii) The one-partition factor 2^k and the principle that joint signs must be counted according to overlap compatibility come from Heckel (2025, Proposition 6). Section V rederives the first identity, and Section VI replaces the cited pairwise upper bound by an exact component formula for every overlap matrix.
- (iii) The strategy of obtaining a rare second-moment seed and then applying concentration amplification follows Heckel (2025, Proposition 5 and the proof of Theorem 1). Sections VI–IX prove a different signed second-moment estimate, and Section X proves the quantitative amplifier used here rather than invoking Heckel’s theorem.
- (iv) The four-size profile, its uniform entropy certificate, the treatment of all partial diagonals and residual attachments, the claimed all-phase uniformity, and the explicit constant in (0.1) are new claims of this manuscript. In particular, the range-restricted tame-profile existence theorem used by Heckel is not imported.

Main theorem (candidate statement)

For $G_n \sim G(n, 1/2)$,

$$\Pr \left\{ \chi(G_n) - \zeta(G_n) \geq \frac{(\ln 2)^2}{32} \ln \left(\frac{200}{153} \right) \frac{n}{(\ln n)^3} \right\} \longrightarrow 1.$$

In particular, $\chi(G_n) - \zeta(G_n)$ diverges with high probability along the full sequence of integers n .

Proof roadmap. The candidate argument has the following structure. Sections II–IV establish an unrestricted lower location for $\chi(G_n)$. Section V constructs a four-size signed cocolouring profile whose first moment is positive at a threshold lower by order $n/(\ln n)^3$. Sections VI–IX establish the corresponding normalized second-moment estimate: Section VI gives the exact overlap and sign decomposition, Section VII controls common whole classes, Section VIII sums the high-cell skeletons, and Section IX bounds the remaining conditional local and cycle contributions. Section X amplifies the resulting possibly rare cocolouring event to a high-probability statement. Finally, Section XI combines the chromatic lower location with the cocolouring upper location. The roadmap records dependencies only; each asymptotic estimate is proved at the point where it is used.

I. Notation and elementary facts

All logarithms are natural unless a base is displayed.

For an integer s , let

$$\mu_s(v) = \binom{v}{s} 2^{-\binom{s}{2}}, \quad \mu_s = \mu_s(n). \quad (1.1)$$

We use the following standard inequalities in their stated forms.

1. For integers $m \geq 1$,

$$\ln(m!) = m \ln m - m + \frac{1}{2} \ln(2\pi m) + O(1/m). \quad (1.2)$$

In particular, uniformly for $m \geq 0$, with $0 \ln 0 = 0$, the last two terms may be replaced by $O(\ln(m+1))$.

2. If a random variable Y is a function of r independent blocks and changing one block changes Y by at most one, then

$$\Pr\{|Y - \mathbb{E}Y| \geq t\} \leq 2 \exp(-2t^2/r). \quad (1.3)$$

We will use the corresponding one-sided bounds as well.

3. If $Z \geq 0$ and $0 < \mathbb{E}Z < \infty$, then

$$\Pr\{Z > 0\} \geq \frac{(\mathbb{E}Z)^2}{\mathbb{E}Z^2}. \quad (1.4)$$

4. If $X \sim \text{Bin}(m, 1/2)$, then

$$\Pr\{X \leq m/4\} \leq e^{-m/16}. \quad (1.5)$$

Indeed, exponential Markov with $t = \ln 3$ bounds the probability by $[3^{1/4}(2/3)]^m \leq e^{-m/16}$.

5. For every nonnegative random variable X and $a > 0$, $\Pr\{X \geq a\} \leq \mathbb{E}X/a$.

The first is Stirling's estimate, the second is McDiarmid's bounded-differences inequality, and the third is the zero-threshold case of Paley-Zygmund. The fourth is the only binomial-tail estimate used below; the fifth is Markov's inequality. The bounded-differences formulation is the standard one recorded by McDiarmid (1989, pp. 148–188).

II. The complete independence-number phase

Define

$$\alpha_0 = 2 \log_2 n - 2 \log_2 \log_2 n + 2 \log_2(e/2) + 1, \quad \alpha = \lfloor \alpha_0 \rfloor, \quad \delta = \alpha_0 - \alpha. \quad (2.1)$$

Thus $0 \leq \delta < 1$. The estimates below are uniform on this entire interval, including sequences approaching either endpoint.

Lemma 2.1 (phase expansion)

There is a bounded continuous function $K(\delta)$ such that

$$\ln \mu_\alpha = \delta \ln n + \left(\frac{2}{\ln 2} - \frac{1}{2} - \delta \right) \ln \ln n + K(\delta) + O((\ln \ln n)^2 / \ln n). \quad (2.2)$$

In particular,

$$\mu_{\alpha+2} = n^{\delta-2+o(1)} \leq n^{-1+o(1)} = o(1), \quad (2.3)$$

and, with $a = \alpha - 2$,

$$\mu_a \geq cn^2(\ln n)^{2/\ln 2 - 5/2} \quad (2.4)$$

for an absolute $c > 0$ and all sufficiently large n .

Proof

Write

$$C = 1 + \ln(\ln 2) - \ln 2, \quad S = \ln n - \ln \ln n + C, \quad b = 1 - \delta.$$

Then $\alpha = 2S/\ln 2 + b$. Since $\alpha = O(\ln n)$, (1.2) gives

$$\ln(n)_\alpha = \alpha \ln n + \sum_{j=0}^{\alpha-1} \ln(1 - j/n) = \alpha \ln n + O(\alpha^2/n),$$

and hence

$$\ln \mu_\alpha = \ln(n)_\alpha - \ln(\alpha!) - \ln 2 \binom{\alpha}{2}.$$

Applying Stirling's formula to $\alpha!$ in this identity gives

$$\ln \mu_\alpha = \alpha \left(\ln n - \ln \alpha + 1 - \frac{\ln 2}{2}(\alpha - 1) \right) - \frac{1}{2} \ln(2\pi\alpha) + O(1/\ln n). \quad (2.5)$$

Expanding $\ln \alpha$ at $2\ln n/\ln 2$, uniformly in $b \in (0, 1]$, gives

$$\ln n - \ln \alpha + 1 - \frac{\ln 2}{2}(\alpha - 1) = \frac{\ln 2\delta}{2} + \frac{\ln \ln n - C - b \ln 2/2}{\ln n} + O((\ln \ln n)^2 / (\ln n)^2). \quad (2.6)$$

Multiplication by α proves (2.2). For reference, the bounded constant is

$$\begin{aligned} K(\delta) = & \delta C + \frac{\ln 2}{2} \delta(1 - \delta) - \frac{2C}{\ln 2} - (1 - \delta) \\ & - \frac{1}{2} \ln(2\pi) - \frac{\ln 2}{2} + \frac{1}{2} \ln(\ln 2). \end{aligned} \quad (2.7)$$

The exact adjacent-size ratios are

$$\frac{\mu_{s+1}}{\mu_s} = \frac{n-s}{s+1} 2^{-s}, \quad \frac{\mu_{s-1}}{\mu_s} = \frac{s}{n-s+1} 2^{s-1}. \quad (2.8)$$

At $s = \alpha + O(1)$, the first is $\Theta(\ln n / n)$ and the second is $\Theta(n / \ln n)$, uniformly in δ . Thus

$$\mu_{\alpha+2} = \mu_\alpha \Theta\left(\frac{(\ln n)^2}{n^2}\right), \quad \mu_{\alpha-2} = \mu_\alpha \Theta\left(\frac{n^2}{(\ln n)^2}\right). \quad (2.8a)$$

The first relation, together with (2.2), proves (2.3). More explicitly, $\delta(\ln n - \ln \ln n) \geq 0$, while K is bounded, so (2.2) gives $\ln \mu_\alpha \geq (2/\ln 2 - 1/2) \ln \ln n + \inf_\delta K(\delta) + o(1)$ uniformly in the phase. Hence $\mu_\alpha \geq c_1 (\ln n)^{2/\ln 2 - 1/2}$. The second relation in (2.8a) now proves (2.4). $\square$

Let $X_{\alpha+2}$ count independent sets of size $\alpha + 2$. Then $\mathbb{E}X_{\alpha+2} = \mu_{\alpha+2}$, and $\alpha(G_n) > \alpha + 1$ implies $X_{\alpha+2} \geq 1$. Thus (2.3) and Markov's inequality give

$$\Pr\{\alpha(G_n) > \alpha + 1\} = o(1). \quad (2.9)$$

III. Continuous profile roots

For a class size u , put

$$d_u = 2^{\binom{u}{2}} u!.$$

Write $u = \alpha - i$, so that i is the deficit from α . Let

$$S_+ = \{-1, 0, 1, 2, \dots\}, \quad S_4 = \{2, 3, 4, 5\}. \quad (3.1)$$

At finite n , S_+ is cut off at $i = \alpha - 1$, since class sizes are positive. For a real class count k , define

$$L_k = n \ln n - n + k - \sum_i k_i \ln(k_i d_{\alpha-i}), \quad (3.2)$$

and let $L_S(n, k)$ be its maximum over real $k_i \geq 0$ satisfying

$$\sum_{i \in S} k_i = k, \quad \sum_{i \in S} (\alpha - i) k_i = n. \quad (3.3)$$

Set

$$T = \alpha - \frac{n}{k}. \quad (3.4)$$

The following lemma supplies all root geometry needed later.

Lemma 3.1 (root phase, derivative, and support comparison)

Let

$$s_0 = \alpha_0 - 1 - \frac{2}{\ln 2}, \quad T_0 = \alpha - s_0 = 1 + \frac{2}{\ln 2} - \delta. \quad (3.5)$$

For $S \in \{S_+, S_4\}$, and uniformly for $0 \leq c \leq \ln 2$, the equation $L_S(n, k) + ck = 0$ has a unique zero $r_{S,c}$ in the root corridor described below, and

$$\frac{n}{r_{S,c}} = s_0 + O(\ln \ln n / \ln n), \quad T = \alpha - \frac{n}{r_{S,c}} = T_0 + O(\ln \ln n / \ln n). \quad (3.6)$$

For every fixed $A > 0$, uniformly when $|n/k - s_0| \leq A \ln \ln n / \ln n$,

$$\frac{\partial}{\partial k} \{L_S(n, k) + ck\} = \frac{2}{\ln 2} (\ln n)^2 + O(\ln n \ln \ln n). \quad (3.7)$$

For two supports R, S , at the same feasible k ,

$$\frac{L_R(n, k) - L_S(n, k)}{k} = \mathcal{F}_{n,R}(T) - \mathcal{F}_{n,S}(T), \quad (3.8)$$

where

$$\mathcal{F}_{n,S}(T) = \max_{\substack{\sum p_i = 1 \\ \sum i p_i = T}} \left[- \sum_i p_i \ln p_i + \sum_i p_i h_n(i) \right]. \quad (3.8a)$$

To make the uniformity precise, fix

$$K_* = [2/\ln 2 - 1/10, 1 + 2/\ln 2 + 1/10] \subset (2, 5). \quad (3.9a)$$

Then, uniformly for $T \in K_*$,

$$\mathcal{F}_{n,S}(T) \longrightarrow \mathcal{F}_S(T) := \max_{\substack{\sum p_i = 1 \\ \sum i p_i = T}} \left[- \sum p_i \ln p_i - \frac{\ln 2}{2} \sum i^2 p_i \right]. \quad (3.9)$$

The effective Lagrange tilts are uniformly bounded on K_* . For S_4 , if $p_{n,i}(T)$ and $p_i(T)$ denote the finite and limiting optimizers, then

$$\sup_{T \in K_*} \max_{2 \leq i \leq 5} |p_{n,i}(T) - p_i(T)| = o(1), \quad \inf_{T \in K_*} \min_{2 \leq i \leq 5} p_{n,i}(T) > 0. \quad (3.9b)$$

All convergence and error terms are uniform in δ .

Proof

Put $s = n/k$ and $p_i = k_i/k$. Dividing (3.2) by k gives

$$\frac{L_S(n, n/s)}{n/s} = (s-1) \ln n - s + 1 + \ln s + \max_p \left[- \sum p_i \ln p_i - \sum p_i \ln d_{\alpha-i} \right]. \quad (3.10)$$

There is an exact affine-plus-curved decomposition

$$-\ln d_{\alpha-i} = A_n + B_n i + h_n(i), \quad (3.11)$$

where

$$A_n = -\ln d_\alpha, \quad B_n = \ln 2\alpha - \ln 2/2 + \ln \alpha, \quad (3.12)$$

and, for $i \geq 0$,

$$h_n(i) = -\frac{\ln 2}{2}i^2 + \sum_{r=0}^{i-1} \ln(1 - r/\alpha). \quad (3.13)$$

For $i = -1$, direct subtraction gives $h_n(-1) = -\ln 2/2 + \ln(\alpha/(\alpha + 1))$. Thus, for fixed i ,

$$h_n(i) = -\ln 2i^2/2 + O(i^2/\alpha), \quad (3.14)$$

while (3.13) also supplies the uniform bound $h_n(i) \leq -\ln 2i^2/2$. Since the target mean T_0 stays in the compact interval

$$\frac{2}{\ln 2} \leq T_0 \leq 1 + \frac{2}{\ln 2}, \quad (3.15)$$

the mean maps at two fixed tilts bracket all of K_* , for both supports and all sufficiently large n . The effective Lagrange tilt is therefore in a fixed compact interval. Its optimizer is proportional to $\exp(\lambda i + h_n(i))$. On every bounded tilt interval, the weights and their first moments are dominated by a constant multiple of $\exp(C|i| - \ln 2i^2/2)$, a summable sequence. Hence the partition functions and their first derivatives converge uniformly. Their mean maps are strictly increasing, with derivative equal to a positive variance; compactness then gives uniform convergence of the inverse tilts. This proves (3.9), and on the finite support S_4 also proves (3.9b). The affine terms in (3.11) prove the exact difference identity (3.8).

Substitute (3.11) into (3.10). If $T = \alpha - s$, the scalar part at $s = s_0$ is, by Stirling,

$$s_0 \left(\ln n - \ln \alpha - \frac{\ln 2}{2}s_0 + \frac{\ln 2}{2} \right) - \ln n + T_0 + 1 + \ln s_0 + \frac{\ln 2}{2}T_0^2 - \frac{1}{2} \ln(2\pi\alpha) + O(1/\ln n). \quad (3.15a)$$

Now $\ln 2s_0/2 = \ln n - \ln \ln n + \ln(\ln 2) - \ln 2$ exactly, while $\ln \alpha = \ln s_0 + T_0/s_0 + O(1/(\ln n)^2)$ and $\ln s_0 = \ln \ln n + \ln 2 - \ln(\ln 2) + O(\ln \ln n / \ln n)$. These relations reduce (3.15a) to $O(\ln \ln n)$, uniformly in the phase. The entropy term is $O(1)$ by the Gaussian tail just proved. Hence

$$L_S(n, n/s_0)/(n/s_0) = O(\ln \ln n). \quad (3.16)$$

Write $\Psi_S(s) = L_S(n, n/s)/(n/s)$, and let $\lambda_{n,S}(T)$ be the effective tilt of the maximizing profile at mean $T = \alpha - s$. Differentiating (3.10) by the envelope theorem gives

$$\Psi'_S(s) = \ln n - 1 + s^{-1} - B_n + \lambda_{n,S}(T).$$

The tilt is uniformly bounded on the fixed target interval, and

$$B_n = 2\ln n - \ln \ln n + O(1), \quad (3.17)$$

therefore (3.17) gives

$$\Psi'_S(s) = -\ln n + O(\ln \ln n) \quad (3.18)$$

uniformly in a fixed neighbourhood of s_0 . For $\Psi_{S,c}(s) = \Psi_S(s) + c$, (3.16) gives $\Psi_{S,c}(s_0) = O(\ln \ln n)$ and (3.18) gives a negative derivative of magnitude at least $\ln n/2$. Taking a sufficiently large fixed corridor constant brackets zero at $s = s_0 - A \ln \ln n / \ln n$ and $s = s_0 + A \ln \ln n / \ln n$, and strict monotonicity gives the unique corridor zero and (3.6), uniformly for $0 \leq c \leq \ln 2$.

Throughout any fixed corridor $|s - s_0| \leq A \ln \ln n / \ln n$, the same equations give $\Psi_{S,c}(s) = O_A(\ln \ln n)$. Since $L_S + ck = k\Psi_{S,c}(s)$ and $ds/dk = -s/k$,

$$\frac{d}{dk}\{L_S + ck\} = \Psi_{S,c}(s) - s\Psi'_S(s) = \frac{2}{\ln 2}(\ln n)^2 + O_A(\ln n \ln \ln n). \quad (3.19)$$

This is (3.7). $\square$

IV. A valid unrestricted lower location for χ

Let $r_+(n) = r_{S_+,0}$ be the zero in Lemma 3.1, and put

$$k_\chi^- := \lfloor r_+(n) \rfloor - \lfloor \ln n \rfloor. \quad (4.1)$$

For an integer profile, direct enumeration gives

$$\mathbb{E}X_{\mathbf{k}} = \frac{n!}{\prod_i ((\alpha - i)!)^{k_i} k_i!} 2^{-\sum_i k_i \binom{\alpha-i}{2}}. \quad (4.2)$$

There are at most $(n+1)^{\alpha+1} = \exp(O((\ln n)^2))$ bounded profiles. Applying (1.2) to (4.2), uniformly even when some $k_i = 0$, yields

$$\ln E_{n,k,\alpha+1} \leq L_+(n, k) + O((\ln n)^2), \quad (4.3)$$

where the left side is the logarithm of the expected total number of unordered, exactly k -part, $(\alpha+1)$ -bounded colourings. By Lemma 3.1,

$$L_+(n, k_\chi^-) \leq -c(\ln n)^3, \quad (4.4)$$

so the expectation in (4.3) is $o(1)$. On the event (2.9), any colouring with at most k_χ^- nonempty parts can be refined, by splitting parts, to exactly k_χ^- nonempty independent parts, all of size at most $\alpha+1$. Therefore

$$\Pr\{\chi(G_n) \leq k_\chi^-\} = o(1). \quad (4.5)$$

This is an unrestricted chromatic lower bound; the size cap has been removed probabilistically by (2.9).

V. The four-size signed first-moment advantage

For $S \in \{S_+, S_4\}$, introduce the tilted weights, partition function, and mean map

$$w_i(\lambda) := \exp\left(\lambda i - \frac{\ln 2}{2} i^2\right), \quad Z_S(\lambda) := \sum_{i \in S} w_i(\lambda),$$

$$M_S(\lambda) := \frac{\sum_{i \in S} i w_i(\lambda)}{Z_S(\lambda)} = \frac{d}{d\lambda} \ln Z_S(\lambda).$$

The Gaussian factor makes both sums finite. Moreover, $M'_S(\lambda) = \text{Var}_{S,\lambda}(i) > 0$, so for every target mean T in the interior of the convex hull of S there is a unique tilt $\lambda_S(T)$ with $M_S(\lambda_S(T)) = T$. The optimizer in (3.9) is therefore

$$p_i = \frac{w_i(\lambda_S(T))}{Z_S(\lambda_S(T))} = \frac{e^{\lambda_S(T)i - \ln 2i^2/2}}{\sum_{j \in S} e^{\lambda_S(T)j - \ln 2j^2/2}}. \quad (5.1)$$

The same calculation gives the dual representation

$$\mathcal{F}_S(T) = \inf_{\lambda \in \mathbb{R}} \{\ln Z_S(\lambda) - \lambda T\} = \ln Z_S(\lambda_S(T)) - \lambda_S(T)T.$$

Define

$$D_4(\delta) = \mathcal{F}_{S_+}(T_0) - \mathcal{F}_{S_4}(T_0), \quad \gamma_4 = \ln \frac{200}{153}. \quad (5.2)$$

Lemma 5.1 (uniform entropy certificate)

For every $0 \leq \delta \leq 1$,

$$0 \leq D_4(\delta) < \ln \frac{153}{100}, \quad \ln 2 - D_4(\delta) > \gamma_4. \quad (5.3)$$

Proof

Fix $\delta \in [0, 1]$, put $T_0 = 1 + 2/\ln 2 - \delta$, and abbreviate

$$\lambda_4 := \lambda_{S_4}(T_0).$$

We first bracket this tilt. At $\lambda = 2\ln 2$, we make the reindexing completely explicit. The old summation index ranges over $i \in S_4 = \{2, 3, 4, 5\}$. Define the new index by

$$j := i - 2.$$

Then $i \mapsto j = i - 2$ is a bijection from $\{2, 3, 4, 5\}$ onto $\{0, 1, 2, 3\}$, with inverse $i = j + 2$. Thus the lower and upper bounds change from $2 \leq i \leq 5$ to $0 \leq j \leq 3$, respectively. Substituting $i = j + 2$ in the weight gives

$$\begin{aligned} w_{j+2}(2\ln 2) &= \exp\left(2\ln 2(j+2) - \frac{\ln 2}{2}(j+2)^2\right) \\ &= \exp\left(2\ln 2 - \frac{\ln 2}{2}j^2\right) = 4 \cdot 2^{-j^2/2}. \end{aligned}$$

Consequently, reindexing both sums in the definition of the mean—and only reindexing them, without discarding any term—gives

$$\begin{aligned}
Z_{S_4}(2\ln 2) &= \sum_{i=2}^5 w_i(2\ln 2) = 4 \sum_{j=0}^3 2^{-j^2/2}, \\
\sum_{i=2}^5 i w_i(2\ln 2) &= 4 \sum_{j=0}^3 (j+2)2^{-j^2/2} \\
&= 8 \sum_{j=0}^3 2^{-j^2/2} + 4 \sum_{j=0}^3 j2^{-j^2/2}.
\end{aligned}$$

Dividing the second equality by the first explains the additive constant 2 in the following formula:

$$\begin{aligned}
M_{S_4}(2\ln 2) &= 2 + \frac{\sum_{j=0}^3 j2^{-j^2/2}}{\sum_{j=0}^3 2^{-j^2/2}} \\
&< 2 + \frac{\sum_{j \geq 0} j2^{-j^2/2}}{\sum_{j \geq 0} 2^{-j^2/2}} < 2 + \frac{4}{5} < \frac{2}{\ln 2} \leq T_0.
\end{aligned} \tag{5.4}$$

For completeness, the first strict inequality in (5.4) has the following direct verification. Put

$$A = \sum_{j=0}^3 2^{-j^2/2}, \quad B = \sum_{j=0}^3 j2^{-j^2/2}, \quad C = \sum_{j=4}^{\infty} 2^{-j^2/2}, \quad D = \sum_{j=4}^{\infty} j2^{-j^2/2}.$$

Here $B/A \leq 3$, while $D/C \geq 4$. Since $A, C > 0$,

$$\frac{B+D}{A+C} - \frac{B}{A} = \frac{AD-BC}{A(A+C)} > 0.$$

Thus adjoining the terms whose new indices satisfy $j \geq 4$ strictly increases the weighted mean. For the next inequality, retain the first three denominator terms, sum the numerator through $j = 3$, and bound its remaining tail by $1/60$. Finally, $2/3 < \ln 2 < 5/7$ gives $2 + 4/5 < 2/\ln 2$.

At $\lambda = 9\ln 2/2$, put $t = 2^{-1/8}$. Completing the square shows that $w_i(9\ln 2/2)$ is proportional to $t^{(2i-9)^2}$. Thus the four weights, in the order $i = 2, 3, 4, 5$, are proportional to

$$t^{25}, \quad t^9, \quad t, \quad t.$$

The numerator of $M_{S_4}(9\ln 2/2) - 4$ is therefore

$$t - t^9 - 2t^{25} = t(1 - t^8 - 2t^{24}) = t/4 > 0, \tag{5.5}$$

and the denominator is positive. Hence this mean is greater than four, whereas $T_0 \leq 1 + 2/\ln 2 < 4$. Since M_{S_4} is strictly increasing, the two mean comparisons give

$$2\ln 2 < \lambda_4 < 9\ln 2/2. \tag{5.6}$$

For a tilt λ , define the omitted low- and high-deficit ratios

$$L(\lambda) := \frac{\sum_{i=-1}^1 w_i(\lambda)}{Z_{S_4}(\lambda)}, \quad H(\lambda) := \frac{\sum_{i \geq 6} w_i(\lambda)}{Z_{S_4}(\lambda)}.$$

The logarithmic derivative of L is a weighted mean on $\{-1, 0, 1\}$ minus one on $\{2, 3, 4, 5\}$, and is therefore negative. Similarly, the logarithmic derivative of H is a weighted mean on $\{6, 7, \dots\}$ minus one on $\{2, 3, 4, 5\}$, and is positive. Direct summation and Gaussian-tail bounds give

$$L(2\ln 2) < \frac{51}{100}, \quad H(3\ln 2) < \frac{1}{50}, \quad L(3\ln 2) < \frac{3}{25}, \quad H(9\ln 2/2) < \frac{1}{4}. \quad (5.7)$$

Here are explicit checks. At $2\ln 2$, with $b = 2^{-1/2}$,

$$L(2\ln 2) = \frac{b + 1/4 + b/16}{1 + b + 1/4 + b/16} < 51/100.$$

At $3\ln 2$, after centring at deficit three, the denominator is $5/4 + 2b$, the low numerator is $1/256 + b/16 + 1/4$, and the high numerator is at most $b/16 + 1/256 + 1/3968$. For the last term, the first omitted summand is $b^{25} = b/4096$, and each successive ratio is at most $1/32$, so the entire remaining tail is less than $1/3968$. The two inequalities at $3\ln 2$ now follow from $7/10 < b < 71/100$.

Finally, after division by t , the high tail at $9\ln 2/2$ begins with t^8, t^{24}, t^{48} . From the third term onward the successive ratios are at most $1/16$, so that residual tail is less than $1/60$. Since $t^8 = 1/2$ and $t^{24} = 1/8$,

$$3(t^8 + t^{24}) + 4/60 < 2, \quad (5.8)$$

and this is equivalent to $H(9\ln 2/2) < 1/4$.

If $\lambda_4 \leq 3\ln 2$, monotonicity and (5.7) give $L(\lambda_4) + H(\lambda_4) < 53/100$; if $\lambda_4 \geq 3\ln 2$, they give $L(\lambda_4) + H(\lambda_4) < 37/100$. By the dual representation above,

$$\mathcal{F}_{S_4}(T_0) = \ln Z_{S_4}(\lambda_4) - \lambda_4 T_0,$$

whereas evaluating the S_+ dual function at the same parameter gives

$$\mathcal{F}_{S_+}(T_0) \leq \ln Z_{S_+}(\lambda_4) - \lambda_4 T_0.$$

Subtracting these two relations yields

$$\begin{aligned} D_4(\delta) &\leq \ln \frac{Z_{S_+}(\lambda_4)}{Z_{S_4}(\lambda_4)} \\ &= \ln(1 + L(\lambda_4) + H(\lambda_4)) < \ln(153/100). \end{aligned}$$

Since S_4 is a subset of S_+ , the loss is nonnegative. Subtracting from $\ln 2$ completes the proof.

□

The following 2^k gain is the one-partition identity isolated by Heckel (2025, Proposition 6). We repeat its short argument because it is an input to the new root calculation. For a fixed partition into k classes, assigning each class the declaration “independent” or “complete” multiplies its first moment by 2^k : every declaration prescribes all internal edge bits, and all 2^k

declarations have probability $2^{-\Sigma \binom{u}{2}}$. The declarations are disjoint here because all four allowed class sizes tend to infinity (in particular, there are no singleton classes). Let $r_4^{\text{co}} = r_{S_4, \ln 2}$ be the unique corridor zero of

$$L_{S_4}(n, k) + \ln 2k. \quad (5.9)$$

Put

$$\Phi_n(k) := L_{S_4}(n, k) + \ln 2k.$$

Then $\Phi_n(r_4^{\text{co}}) = 0$ by definition, while at $k = r_+$, Lemma 3.1 and (5.2) give

$$L_{S_4}(n, r_+) + \ln 2r_+ = r_+ \{\ln 2 - D_4(\delta) + o(1)\}. \quad (5.10)$$

Both roots lie in the corridor (3.6), so the whole interval between them is inside a fixed corridor. The mean-value theorem therefore gives a point $\xi_n \in (r_4^{\text{co}}, r_+)$ such that

$$\Phi_n(r_+) - \Phi_n(r_4^{\text{co}}) = \Phi'_n(\xi_n)(r_+ - r_4^{\text{co}}).$$

Now $r_+ = (\ln 2/2 + o(1))n/\ln n$, and (3.7), uniformly at ξ_n , gives

$$\Phi'_n(\xi_n) = \frac{2}{\ln 2}(\ln n)^2 + O((\ln n)(\ln \ln n)).$$

Substituting this and (5.10) into the preceding identity yields

$$r_+ - r_4^{\text{co}} = \left(\frac{(\ln 2)^2}{4} \{\ln 2 - D_4(\delta)\} + o(1) \right) \frac{n}{(\ln n)^3}. \quad (5.11)$$

In particular, for all sufficiently large n ,

$$r_+ - r_4^{\text{co}} \geq \frac{(\ln 2)^2 \gamma_4}{8} \frac{n}{(\ln n)^3}. \quad (5.12)$$

Choose the midpoint integer

$$k_{\text{co}} = \left\lceil \frac{r_4^{\text{co}} + r_+}{2} \right\rceil. \quad (5.13)$$

At this exact k , let $p_i^{(n)}$ be the exact finite- n maximizer of L_{S_4} . Let $\lambda_{n, \text{def}}$ denote the Lagrange multiplier for the deficit-mean constraint, with the sign convention that its contribution is $e^{\lambda_{n, \text{def}} i}$. The Lagrange equations and the decomposition (3.11) then give

$$p_i^{(n)} \propto d_{\alpha-i}^{-1} e^{\lambda_{n, \text{def}} i} \propto e^{\widehat{\lambda}_n i + h_n(i)}, \quad \widehat{\lambda}_n := B_n + \lambda_{n, \text{def}}. \quad (5.14)$$

Thus $\lambda_{n, \text{def}}$ is the multiplier before the affine term $B_n i$ is absorbed, while $\widehat{\lambda}_n$ is the effective tilt in the limiting Gaussian coordinates. (If a size-coordinate multiplier τ_n is used instead, then $\lambda_{n, \text{def}} = -\tau_n$.) The target deficit mean is $\alpha - n/k_{\text{co}}$. Lemma 3.1 makes these proportions converge uniformly to (5.1), with $\widehat{\lambda}_n \rightarrow \lambda_4(T_0)$. Equation (5.6) also shows directly that the four limiting weights differ by at most a fixed factor. Thus

$$\min_{2 \leq i \leq 5} p_i^{(n)} \geq c_p > 0 \quad (5.15)$$

uniformly in the phase.

Round $k_{\text{co}} p_i^{(n)}$ to preliminary integers $\tilde{k}_i$. Define the signed constraint errors

$$e_0 = \sum_i \tilde{k}_i - k_{\text{co}}, \quad e_1 = \sum_i i \tilde{k}_i - (\alpha k_{\text{co}} - n),$$

so $e_0, e_1 = O(1)$, and add

$$\Delta k_2 = e_1 - 3e_0, \quad \Delta k_3 = 2e_0 - e_1. \quad (5.16)$$

Indeed, $\Delta k_2 + \Delta k_3 = -e_0$ and $2\Delta k_2 + 3\Delta k_3 = -e_1$. This enforces both constraints exactly, changes only $O(1)$ counts, and preserves positivity by (5.15). Put $\mathbf{k}^* = (k_{\text{co}} p_i^{(n)})_{i=2}^5$ and let $\Delta^{\text{tot}} = \mathbf{k} - \mathbf{k}^*$ be the total rounding and correction displacement. It has bounded norm and satisfies both homogeneous constraint equations, because $\mathbf{k}^*$ and $\mathbf{k}$ obey the same two constraints. Thus it is tangent to the feasible affine plane. At the exact optimizer the linear term vanishes, while the Hessian of the entropy term is $-\text{diag}(1/k_i)$. Hence

$$L_{\mathbf{k}^* + \Delta^{\text{tot}}} - L_{\mathbf{k}^*} = O\left(\sum_i \frac{(\Delta_i^{\text{tot}})^2}{k_i^*}\right) = O(1/k_{\text{co}}).$$

Applying (1.2) to the factorial terms gives an exact integer vector

$$\mathbf{k} = (k_2, k_3, k_4, k_5), \quad u_i = \alpha - i, \quad k_i = \Theta(n/\ln n), \quad (5.17)$$

whose signed first moment is

$$\mathbb{E} Z_{\mathbf{k}}^{\text{sgn}} = 2^{k_{\text{co}}} \frac{n!}{\prod_i (u_i!)^{k_i} k_i!} 2^{-\sum_i k_i \binom{u_i}{2}}, \quad (5.18)$$

and

$$\ln \mathbb{E} Z_{\mathbf{k}}^{\text{sgn}} = L_{S_4}(n, k_{\text{co}}) + \ln 2k_{\text{co}} + O(\ln n) \geq c_Z k_{\text{co}} \quad (5.19)$$

for a phase-independent $c_Z > 0$. Indeed, the midpoint is at distance at least $(r_+ - r_4^{\text{co}})/2 - O(1)$ to the right of r_4^{co} . Integrating the positive derivative (3.7) over that interval and using (5.12) gives

$$L_{S_4}(n, k_{\text{co}}) + \ln 2k_{\text{co}} \geq c_Z k_{\text{co}},$$

after decreasing c_Z if necessary. Every u_i tends to infinity, so the independent and complete declarations of one part are disjoint. Thus $Z_{\mathbf{k}}^{\text{sgn}} > 0$ implies an actual cocolouring with k_{co} parts.

Finally, (4.1), (5.12), and (5.13) give

$$k_{\chi}^- - k_{\text{co}} \geq \left(\frac{(\ln 2)^2 \gamma_4}{16} - o(1) \right) \frac{n}{(\ln n)^3}. \quad (5.20)$$

VI. Exact signed second-moment representation

For two partitions sharing exactly ℓ whole classes, [Heckel \(2025, Proposition 6\)](#) bounded the number of joint sign declarations by $2^{2k-\ell}$, thereby transferring selected ordinary-colouring overlap bounds to coccolourings. Lemma 6.1 below does not import that inequality. It keeps the entire overlap matrix and counts the compatible signs exactly: compatibility is controlled by the components of the graph of cells of size at least two. Whole common classes appear as one special case of this component formula.

Temporarily label all slots within each of the four types. This multiplies the unordered witness by the deterministic factor $\prod_i k_i!$ and hence does not change its normalized second moment. Choose two independent uniform ordered partitions of the profile. If their slots are indexed by a and b , put

$$r_{ab} = |V_a \cap W_b|. \quad (6.1)$$

The overlap matrix has the exact law

$$p(r) = \frac{\prod_a s_a! \prod_b t_b!}{n! \prod_{a,b} r_{ab}!}, \quad (6.2)$$

where here both degree lists are the multiset given by (5.17). Equivalently, place s_a stubs at each row slot and t_b stubs at each column slot and take a uniform perfect matching of the two n -stub sets.

Let $H(r)$ be the simple bipartite graph whose edges are cells with $r_{ab} \geq 2$, omitting isolated slot vertices. Write

$$W = \sum_{a,b} \binom{r_{ab}}{2}, \quad \beta(H) = |E(H)| - |V(H)| + c(H). \quad (6.3)$$

Lemma 6.1 (exact sign sum)

The normalized local factor of the pair of partitions is

$$A_\zeta(r) = 2^{W+c(H)-|V(H)|} = \left(\prod_{a,b} g(r_{ab}) \right) 2^{\beta(H)}, \quad (6.4)$$

where

$$g(0) = g(1) = g(2) = 1, \quad g(x) = 2^{\binom{x}{2}-1} \quad (x \geq 3). \quad (6.5)$$

Consequently

$$\frac{\mathbb{E}[(Z_{\mathbf{k}}^{\text{sgn}})^2]}{(\mathbb{E}Z_{\mathbf{k}}^{\text{sgn}})^2} = \sum_{r \in \mathcal{R}(\mathbf{s}, \mathbf{t})} p(r) A_\zeta(r). \quad (6.6)$$

where $\mathcal{R}(\mathbf{s}, \mathbf{t})$ is the finite set of nonnegative integer matrices with the prescribed row margins $\mathbf{s}$ and column margins $\mathbf{t}$.

Proof

Let $B = \sum_a \binom{s_a}{2}$, which is the number of internal edge bits prescribed by one partition. A row sign and a column sign are compatible on a cell of size at least two exactly when they agree. Therefore signs must be constant on every component of H . There are

$$2^{2k_{\text{co}} - |V(H)| + c(H)}$$

compatible pairs of sign assignments, including the free signs on slots isolated from H . For each compatible pair, W edge bits are prescribed twice, so the probability of all constraints is $2^{-(2B-W)}$. Division by the square of the one-partition signed probability $(2^{k_{\text{co}}} 2^{-B})^2$ gives

$$\frac{2^{2k_{\text{co}} - |V(H)| + c(H)} 2^{-(2B-W)}}{(2^{k_{\text{co}}} 2^{-B})^2} = 2^{W + c(H) - |V(H)|},$$

which is the first expression in (6.4).

For every support edge, split $\binom{r}{2}$ as $(\binom{r}{2} - 1) + 1$. Since $|E| - |V| + c = \beta$,

$$\prod_{a,b} g(r_{ab}) = 2^{W - |E(H)|},$$

and multiplying by $2^{\beta(H)}$ gives the second expression. Averaging over (6.2) proves (6.6). $\square$
The identity

$$2^{\beta(H)} = \#\{F \subseteq E(H) : \deg_F(v) \text{ is even for every } v\} \quad (6.7)$$

will be used repeatedly. It is the elementary fact that the even subgraphs form the binary cycle space of dimension $\beta(H)$.

We also need one exact configuration-model estimate.

Lemma 6.2 (joint prescribed-cell bound)

In a bipartite configuration model of total degree m_0 , with row degrees d_a and column degrees d'_b , prescribe distinct cells D and demands $x_{ab} \geq 1$. Put

$$x = \sum_D x_{ab}, \quad D_a = \sum_b x_{ab}, \quad D'_b = \sum_a x_{ab}.$$

In the feasible case

$$x \leq m_0, \quad D_a \leq d_a \quad (\forall a), \quad D'_b \leq d'_b \quad (\forall b),$$

we have

$$\Pr\{r_{ab} \geq x_{ab} \quad \forall ab \in D\} \leq \frac{\prod_a (d_a)_{D_a} \prod_b (d'_b)_{D'_b}}{(m_0)_x \prod_{ab \in D} x_{ab}!}. \quad (6.8)$$

If any feasibility condition fails, the prescribed event is empty and is handled separately. In particular, in all cases with $m_0 > 0$, with $e := \exp(1)$ and $\theta_{ab} = e d_a d'_b / m_0$, its probability is at most

$$\prod_{ab \in D} \frac{\theta_{ab}^{x_{ab}}}{x_{ab}!}. \quad (6.9)$$

Proof

If a feasibility condition fails, no matching realizes all demands. In the feasible case, for each row a , assigning disjoint row stubs to its demanded cells gives $(d_a)_{D_a} / \prod_b x_{ab}!$ choices. The analogous column choice gives $(d'_b)_{D'_b} / \prod_a x_{ab}!$ choices, and there are $x_{ab}!$ bijections between the selected stubs inside cell ab . Thus the total number of demand witnesses is

$$\frac{\prod_a (d_a)_{D_a} \prod_b (d'_b)_{D'_b}}{\prod_{ab \in D} x_{ab}!}.$$

Any fixed witness occurs with probability $1/(m_0)_x$, because after its x prescribed pairs are exposed the remaining matching is uniform on the unused stubs. A union bound proves (6.8).

It remains to simplify the global denominator. For $1 \leq x \leq m_0$,

$$\binom{m_0}{x} = \prod_{j=0}^{x-1} \frac{m_0 - j}{x - j} \geq \left(\frac{m_0}{x}\right)^x, \quad x! \geq \left(\frac{x}{e}\right)^x.$$

The factorial inequality follows, for example, from $\sum_{j=1}^x \ln j \geq \int_1^x \ln t \, dt = x \ln x - x + 1$. Consequently

$$(m_0)_x = x! \binom{m_0}{x} \geq (m_0/e)^x, \quad (6.10)$$

with the case $x = 0$ immediate. Combining (6.10) with $(d)_r \leq d^r$ proves (6.9). Notice that (6.8) retains the single global falling factorial before (6.9) is taken. $\square$

VII. Exact partial diagonals

Purpose of this section. The exact overlap representation of Section VI still permits two partitions to share whole classes. The definitions below assign a nonnegative marked weight to each such common subprofile. We prove that the total marked contribution is $1 + o(1)$, uniformly through the phase. This partial-diagonal estimate is the input used in Section VIII to control the dense transportation sum. The counting identities in (7.1)–(7.4) are finite- n identities; asymptotic estimates are explicitly labelled below.

For a selected common subprofile $\ell = (\ell_i)$, put

$$\ell_\bullet = \sum_i \ell_i, \quad m = \sum_i u_i \ell_i, \quad b_\ell = \sum_i \binom{u_i}{2} \ell_i. \quad (7.1)$$

Here $0 \leq \ell_i \leq k_i$ coordinatewise. Since the full profile has $\sum_i u_i k_i = n$, this automatically gives $m \leq n$; thus every factorial in the counting formula below has its ordinary combinatorial meaning.

The signed first moment of such partial partitions occupying any m vertices of $[n]$ is

$$Y_\ell^{\text{sgn}} = 2^\ell \frac{n!}{(n-m)! \prod_i (u_i!)^{\ell_i} \ell_i!} 2^{-b_\ell}. \quad (7.2)$$

Summing the possible matchings between selected row and column slots gives the exact *marked* common-subprofile weight

$$D(\ell) = \frac{\prod_i \binom{k_i}{\ell_i}^2}{Y_\ell^{\text{sgn}}}. \quad (7.3)$$

Different marked choices need not be disjoint: an overlap containing several common blocks contributes to each corresponding marked choice. The sums below are sums of these nonnegative marked weights, so no disjointness is used. In particular, $D(0) = 1$ and $D(k) = 1/\mathbb{E} Z_k^{\text{sgn}}$. To add one common block of type i , first note that

$$\frac{\binom{k_i}{\ell_i+1}^2}{\binom{k_i}{\ell_i}^2} = \frac{(k_i - \ell_i)^2}{(\ell_i + 1)^2},$$

while (7.2) gives

$$\frac{Y_{\ell+e_i}^{\text{sgn}}}{Y_\ell^{\text{sgn}}} = \frac{2(n-m)_{u_i}}{u_i! (\ell_i + 1) 2^{\binom{u_i}{2}}} = \frac{2\mu_{u_i}(n-m)}{\ell_i + 1}.$$

Dividing these two ratios gives

$$\frac{D(\ell + e_i)}{D(\ell)} = \frac{(k_i - \ell_i)^2}{2(\ell_i + 1)\mu_{u_i}(n-m)}. \quad (7.4)$$

At the opposite endpoint, put $h = k - \ell$, $v = \sum_i u_i h_i$, and let $Z_h^{\text{sgn}}(v)$ be the signed first moment of a complete profile h on v vertices. Exact factorial cancellation gives

$$D(k - h) = \frac{B(h)}{\mathbb{E} Z_k^{\text{sgn}}}, \quad B(h) = \left(\prod_i \binom{k_i}{h_i} \right) Z_h^{\text{sgn}}(v), \quad (7.5)$$

and

$$\frac{\binom{k_i}{h_i+1}}{\binom{k_i}{h_i}} = \frac{k_i - h_i}{h_i + 1}, \quad \frac{Z_{h+e_i}^{\text{sgn}}(v + u_i)}{Z_h^{\text{sgn}}(v)} = \frac{2\mu_{u_i}(v + u_i)}{h_i + 1}.$$

Therefore

$$\frac{B(h + e_i)}{B(h)} = \frac{2(k_i - h_i)\mu_{u_i}(v + u_i)}{(h_i + 1)^2}. \quad (7.6)$$

In Heckel's admissible-phase argument, the analogous control of partial profiles enters through the tame-colouring input used to prove Heckel (2025, Proposition 5); its hypotheses are the source of the lower μ_α -window in that theorem. The next lemma is the replacement

needed here. It is proved directly for the four-size signed profile, including both corners and every intermediate mass, and no tame-profile theorem is invoked.

Lemma 7.1 (all common subprofiles)

For the exact midpoint profile,

$$\sum_{0 \leq \ell_i \leq k_i} D(\ell) = 1 + o(1). \quad (7.7)$$

Proof: the empty corner

Let

$$\eta = \frac{\ln \ln n}{32 \ln n}, \quad \xi_i = \frac{k_i^2}{2\mu_{u_i}(n)}, \quad \Xi_{\text{empty}} = \sum_i \xi_i. \quad (7.8)$$

Equations (2.4), (2.8), and $k_i = \Theta(n/\ln n)$ imply

$$\Xi_{\text{empty}} = O((\ln n)^{-(2/\ln 2 - 1/2)}) = o(1). \quad (7.9)$$

If the final selected mass is at most ηn , every intermediate mass m' in (7.4) satisfies

$$\frac{\mu_{u_i}(n)}{\mu_{u_i}(n - m')} \leq (1 - 2\eta)^{-u_i} \leq (\ln n)^{3/8}. \quad (7.10)$$

Iterating (7.4), in any order, yields

$$D(\ell) \leq \prod_i \frac{((\ln n)^{3/8} \xi_i)^{\ell_i}}{\ell_i!}.$$

Thus

$$1 \leq \sum_{m \leq \eta n} D(\ell) \leq \exp((\ln n)^{3/8} \Xi_{\text{empty}}) = 1 + o(1). \quad (7.11)$$

Proof: the central range

Write

$$\begin{aligned} p_i &= k_i/k_{\text{co}}, & y_i &= \ell_i/k_{\text{co}}, & Y &= \sum_i y_i, & I &= \sum_i i y_i, \\ z_i &= p_i - y_i, & R &= \sum_i z_i = 1 - Y, & I_r &= \sum_i i z_i = T - I. \end{aligned} \quad (7.12)$$

The residual vertex fraction is

$$\rho = \frac{n - m}{n} = R + \frac{I - TY}{\alpha - T}. \quad (7.13)$$

Uniform Stirling control. Uniform Stirling estimates in (7.3), retaining zeros, give

$$\ln D(\ell) = n\rho \ln \rho + k_{\text{co}} \sum_i \{2p_i \ln p_i - 2(p_i - y_i) \ln(p_i - y_i) - y_i \ln y_i - y_i + y_i E_i\} + O(\ln n), \quad (7.14)$$

where

$$E_i = \ln k_{\text{co}} + \ln(u_i!) + u_i - u_i \ln n + \ln 2 \binom{u_i}{2} - \ln 2. \quad (7.15)$$

Exact subtraction and the definition of α_0 give

$$E_{i+1} - E_i = \ln n - \ln(\alpha - i) - \ln 2(\alpha - i) + \ln 2 - 1 = -\ln 2\alpha/2 + O(1). \quad (7.16)$$

Let $\bar{E} = \sum p_i E_i$. Applying Stirling to the complete moment (5.18) gives

$$-\frac{1}{k_{\text{co}}} \ln \mathbb{E} Z_k^{\text{sgn}} = \sum_i p_i \ln p_i - 1 + \bar{E} + o(1). \quad (7.17)$$

The positive margin (5.19) bounds $\bar{E}$ above. Indeed, (7.17) and $\ln \mathbb{E} Z_k^{\text{sgn}} \geq c_Z k_{\text{co}}$ give $\bar{E} = O(1)$ from above, uniformly in the phase. Since the support consists of four consecutive indices, summing (7.16) over at most three steps and using $\sum_i p_i(i - T) = 0$ gives

$$E_i = \bar{E} - \frac{\ln 2\alpha}{2}(i - T) + O(1) \quad (2 \leq i \leq 5).$$

Multiplying by y_i and summing yields

$$\sum_i y_i E_i \leq \frac{\ln 2\alpha}{2}(TY - I) + O(Y). \quad (7.18)$$

The term $Y\bar{E}$ has been absorbed into $O(Y)$ in (7.18). The remaining entropy expression in (7.14) is $O(Y \ln(e/Y))$, uniformly because every p_i is bounded below. If $\rho \geq 1/32$, then $R \geq 1/64$ for large n , and (7.13) gives

$$n\rho \ln \rho = k_{\text{co}} \alpha R \ln R + O(k_{\text{co}} Y). \quad (7.19)$$

Consequently

$$\ln D(\ell) \leq k_{\text{co}} \alpha \Phi_T(z) + C k_{\text{co}} Y \ln(e/Y) + C \ln n, \quad (7.20)$$

where

$$\Phi_T(z) = R \ln R + \frac{\ln 2}{2}(I_r - TR). \quad (7.21)$$

Uniform rate negativity. This rate is uniformly negative away from its two corners. Indeed,

$$I_r - TR \leq (5 - T)R, \quad I_r - TR = \sum_i (T - i)y_i \leq (T - 2)(1 - R), \quad (7.22)$$

and hence

$$\Phi_T \leq R\{\ln R + \ln 2(5 - T)/2\}, \quad \Phi_T \leq R \ln R + \ln 2(T - 2)(1 - R)/2. \quad (7.23)$$

Here is the numerical check used in this split. On $1/64 \leq R \leq 47/100$, add $Y/5000 = (1 - R)/5000$ to the first bound in (7.23). The resulting function

$$R \ln R + \frac{\ln 2}{2}(5 - T)R + \frac{1 - R}{5000}$$

is convex in R . Its largest possible coefficient occurs at $T = 2/\ln 2$, and direct substitution at $R = 1/64$ and $R = 47/100$, using $\ln 2 < 0.6932$, gives a negative value at both endpoints. Hence $\Phi_T \leq -Y/5000$ throughout this interval.

On $47/100 \leq R \leq 1$, use the second bound in (7.23) and $T \leq 1 + 2/\ln 2$. After adding $Y/200$, it is at most

$$R \ln R + (1 - \ln 2/2 + 1/200)(1 - R)$$

This function is convex, is negative at $47/100$, and equals zero at $R = 1$. It follows that $\Phi_T \leq -Y/200$ on the second interval. Thus, whenever

$$m > \eta n, \quad n - m > n/32, \quad (7.24)$$

we have $Y \geq \eta/2$. The leading term is at most $-k_{\text{co}}\alpha Y/5000$, whereas $\ln(e/Y) = O(\ln \ln n)$, $\ln \ln n/\alpha = o(1)$, and $\ln n = o(k_{\text{co}}\alpha\eta)$. It therefore dominates both error terms in (7.20). Uniformly in this range,

$$D(\ell) \leq e^{-c k_{\text{co}} \ln \ln n}. \quad (7.25)$$

There are only $(k_{\text{co}} + 1)^4$ vectors, so their total is $o(1)$.

Proof: the full corner

It remains to take $v = n - m \leq n/32$. Uniformly for the four sizes, $\mu_{u_i}(n) \leq n^{6+o(1)}$ by (2.2) and (2.8), while

$$\mu_{u_i}(v) \leq \mu_{u_i}(n)(v/n)^{u_i} \leq n^{-4+o(1)}. \quad (7.26)$$

Here $u_i = (2/\ln 2 + o(1))\ln n$ and $32^{-u_i} = n^{-10+o(1)}$. Consequently $\mu_{u_i}(v + u_i) \leq n^{-4+o(1)}$ at every intermediate step while $v \leq n/32$. Since $k_i \leq n$ and $(h_i + 1)^2 \geq 1$, the right side of (7.6) is at most $2n^{-3+o(1)} < 1$. Thus, in any order of adding residual blocks, $B(h)$ never increases from $B(0) = 1$, and hence $B(h) \leq 1$. Equations (5.19) and (7.5) give

$$D(k - h) \leq e^{-cz k_{\text{co}}}. \quad (7.27)$$

Summing $(k_{\text{co}} + 1)^4$ residual vectors proves an $o(1)$ contribution. Together with (7.11) and (7.25), this proves (7.7). $\square$

VIII. Canonical high cells and dense endpoint transportation

Purpose of this section. Section VII controls the diagonal contribution coming from common whole classes. We now decompose every remaining overlap canonically according to its high cells and sum the resulting high-cell contribution, including unequal-type and nonendpoint cells. The local and cycle factors of the unexposed residual configuration are deliberately deferred: Section IX will bound them uniformly after the present bare high-skeleton sum has been established.

Put

$$u_{\max} = \alpha - 2, \quad R_0 = \lfloor u_{\max}/2 \rfloor. \quad (8.1)$$

In every overlap matrix, the cells of multiplicity greater than R_0 will be called *high cells*. They form a matching: two high cells in one row or column would use more than $u_{\max}$ stubs. Write the resulting canonical high-cell skeleton and its multiplicities as

$$\mathcal{M} = \{a_r b_r : 1 \leq r \leq h\}, \quad r_{a_r b_r} = j_r > R_0, \quad J = \sum_r j_r. \quad (8.2)$$

Selecting the exact stub pairs in these cells has incidence

$$\pi(\mathcal{M}, j) = \frac{\prod_r (s_{a_r})_{j_r} (t_{b_r})_{j_r}}{(n)_J \prod_r j_r!}. \quad (8.3)$$

Conditional on the exposed pairs, the remaining matching is a uniform configuration model with total degree $n - J$ and the residual row and column degrees. It is restricted to put no further pair in a cell of $\mathcal{M}$ and no cell above R_0 . Multiplication of (8.3) by the exact residual contingency-table law cancels to (6.2). Thus every overlap matrix occurs exactly once.

We first sum the bare high-skeleton contribution, postponing all local and topological factors in the capped residual model. Index the four sizes as

$$u_i = a - i, \quad 0 \leq i \leq 3, \quad a = \alpha - 2. \quad (8.4)$$

For a typed full-containment matrix $\mathbf{L} = (\ell_{ij})$, put

$$r_i = \sum_j \ell_{ij}, \quad c_j = \sum_i \ell_{ij}, \quad x_{ij} = \min(u_i, u_j), \quad J(\mathbf{L}) = \sum_{ij} x_{ij} \ell_{ij}. \quad (8.5)$$

Its exact incidence, including its local signed rewards but not residual attachments, is

$$W(\mathbf{L}) = \frac{\prod_i (k_i)_{r_i} \prod_j (k_j)_{c_j}}{\prod_{ij} \ell_{ij}!} \frac{1}{(n)_{J(\mathbf{L})}} \prod_{ij} \left[\frac{(u_i)_{x_{ij}} (u_j)_{x_{ij}}}{x_{ij}!} g(x_{ij}) \right]^{\ell_{ij}}. \quad (8.6)$$

For a margin vector r , define

$$D(r) = \frac{\prod_i (k_i)_{r_i}^2 \prod_i [u_i! g(u_i)]^{r_i}}{\prod_i r_i! (n)_{m_r}}, \quad m_r = \sum_i u_i r_i. \quad (8.7)$$

This is exactly the common-subprofile weight (7.3).

Lemma 8.1 (geometric-mean transportation comparison)

For every feasible $\mathbf{L}$,

$$W(\mathbf{L}) \leq \sqrt{D(r)D(c)} \frac{\sqrt{\prod_i r_i! c_i!}}{\prod_{ij} \ell_{ij}!} \prod_{ij} Q_{ij}^{\ell_{ij}}, \quad (8.8)$$

where $Q_{ii} = 1$ and, if the two sizes are $t = s + d$, $1 \leq d \leq 3$,

$$Q_{ij} = (n+1)^{d/2} \frac{\sqrt{(t)_d}}{d!} 2^{-\{ds + \binom{d}{2}\}/2} \leq \frac{\eta_n^d}{d!}, \quad (8.9)$$

with

$$\eta_n = 2^{3/2} \sqrt{\frac{(n+1)^a}{2^a}} = O((\ln n)^{3/2}/\sqrt{n}) = o(1). \quad (8.10)$$

Proof

For integer x , let $f(x) = \ln(n)_x$ and interpolate linearly between successive integers. Its successive slopes are $\ln(n-x)$, which decrease; therefore f is concave, and every slope is at most $\ln(n+1)$. Moreover,

$$\frac{m_r + m_c}{2} = J(\mathbf{L}) + \frac{1}{2} \sum_{ij} |i-j| \ell_{ij}. \quad (8.11)$$

Concavity first gives

$$\frac{f(m_r) + f(m_c)}{2} \leq f\left(\frac{m_r + m_c}{2}\right).$$

Using (8.11) and then moving from $J(\mathbf{L})$ along a distance $\frac{1}{2} \sum_{ij} |i-j| \ell_{ij}$, with slope at most $\ln(n+1)$, gives

$$\frac{f(m_r) + f(m_c)}{2} - f(J(\mathbf{L})) \leq \frac{\ln(n+1)}{2} \sum_{ij} |i-j| \ell_{ij}.$$

Exponentiating this inequality gives

$$\frac{\sqrt{(n)_{m_r} (n)_{m_c}}}{(n)_{J(\mathbf{L})}} \leq (n+1)^{\frac{1}{2} \sum_{ij} |i-j| \ell_{ij}}. \quad (8.12)$$

For $t = s + d$, put $b_u = u!g(u)$. A direct calculation gives

$$\frac{(t)_s (s)_s}{s!} g(s) = b_s \binom{t}{d}, \quad \frac{b_t}{b_s} = (t)_d 2^{ds + \binom{d}{2}}. \quad (8.13)$$

Thus the local factor divided by $\sqrt{b_s b_t}$ is

$$\frac{\sqrt{(t)_d}}{d!} 2^{-\{ds + \binom{d}{2}\}/2}. \quad (8.14)$$

Dividing (8.6) by $\sqrt{D(r)D(c)}$, then using (8.12) and (8.14), proves (8.8)–(8.9). Finally $t \leq a$, $s \geq a - 3$, and

$$2^a = \Theta(n^2/(\ln n)^2) \quad (8.15)$$

uniformly in the phase. This proves (8.10). $\square$

Lemma 8.2 (sum of all endpoint tables)

Let $\mathcal{P}$ be the finite set of feasible one-sided margin vectors. The first sum below is over all typed full-containment tables $\mathbf{L}$ whose row and column margins lie in $\mathcal{P}$. Then

$$\begin{aligned} \sum_{\mathbf{L}} W(\mathbf{L}) &\leq \exp\{O(\eta_n k_{\text{co}})\} \left(\sum_{\mathbf{r} \in \mathcal{P}} \sqrt{D(\mathbf{r})} \right)^2 \\ &\leq \exp\{O(\sqrt{n \ln n}) + O(\ln n)\} \sum_{\mathbf{r} \in \mathcal{P}} D(\mathbf{r}) \\ &= \exp\{O(\sqrt{n \ln n}) + O(\ln n)\}. \end{aligned} \quad (8.16)$$

Proof

For fixed margins, write

$$A_{\mathbf{L}} = \frac{\prod_i r_i!}{\prod_{ij} \ell_{ij}!}, \quad C_{\mathbf{L}} = \frac{\prod_j c_j!}{\prod_{ij} \ell_{ij}!}. \quad (8.17)$$

Write $Q^{\mathbf{L}} := \prod_{ij} Q_{ij}^{\ell_{ij}}$. Cauchy's inequality gives

$$\sum_{\mathbf{L}} \sqrt{A_{\mathbf{L}} C_{\mathbf{L}}} Q^{\mathbf{L}} \leq \left(\sum_{\mathbf{L}} A_{\mathbf{L}} Q^{\mathbf{L}} \right)^{1/2} \left(\sum_{\mathbf{L}} C_{\mathbf{L}} Q^{\mathbf{L}} \right)^{1/2}. \quad (8.18)$$

All summands in (8.18) are nonnegative. In the first sum, remove the column-margin constraints while retaining the fixed row margins; this enlarges the summation domain and therefore gives an upper bound. The unrestricted allocation of the r_i row objects among the column types is then counted exactly by the multinomial theorem. The second inequality is obtained symmetrically by removing the row-margin constraints while retaining the column margins. Thus

$$\sum_{\mathbf{L}} A_{\mathbf{L}} Q^{\mathbf{L}} \leq \prod_i \left(\sum_j Q_{ij} \right)^{r_i}, \quad \sum_{\mathbf{L}} C_{\mathbf{L}} Q^{\mathbf{L}} \leq \prod_j \left(\sum_i Q_{ij} \right)^{c_j}. \quad (8.19)$$

Every row and column sum of Q is at most $1 + C\eta_n$, and the total margin is at most k_{co} . Hence the fixed-margin cost is $\exp(O(\eta_n k_{\text{co}}))$. There are only $O(k_{\text{co}}^4)$ margin vectors, and

$$\left(\sum_{\mathbf{r} \in \mathcal{P}} \sqrt{D(\mathbf{r})} \right)^2 \leq O(k_{\text{co}}^4) \sum_{\mathbf{r} \in \mathcal{P}} D(\mathbf{r}). \quad (8.20)$$

Use Lemma 7.1 and $\eta_n k_{\text{co}} = O(\sqrt{n \ln n})$ to finish. $\square$

Lemma 8.3 (all nonendpoint high multiplicities)

The full-containment sum (8.16) remains $\exp(o(n/(\ln n)^4))$ after all high-cell multiplicities $R_0 < j < \min(u_i, u_j)$ are inserted.

Proof

Let the smaller slot size be m , the larger $m + d$, with $0 \leq d \leq 3$. Replacing the endpoint multiplicity m by $m - e$ has exact local ratio

$$R_{m,d}(e) = \frac{\binom{m}{e}}{(d+1) \cdots (d+e)} 2^{-em+e(e+1)/2}. \quad (8.21)$$

For many decorated cells, with total deficit Q , the ratio of the one global denominators is

$$\frac{\binom{n}{J_0}}{\binom{n}{J_0-Q}} = (n - J_0 + Q)_Q \leq n^Q. \quad (8.22)$$

Put

$$A_e := n^e R_{m,d}(e), \quad 0 \leq e < m/4.$$

For $0 \leq e < m/4$, its consecutive ratio is

$$\frac{n(m-e)2^{-m+e+1}}{(e+1)(d+e+1)}. \quad (8.23)$$

The ratio of (8.23) at $e+1$ to its value at e is

$$2^{\frac{m-e-1}{m-e} \frac{e+1}{e+2} \frac{d+e+1}{d+e+2}}, \quad (8.24)$$

which increases for $e \leq m/4$ and $d \leq 3$. Here is a direct check of the one-turn assertion. If $r(e)$ denotes (8.23), interpolated for real e , then

$$\frac{d^2}{de^2} \ln r(e) = -\frac{1}{(m-e)^2} + \frac{1}{(e+1)^2} + \frac{1}{(d+e+1)^2} > 0 \quad (8.24a)$$

for $0 \leq e \leq m/4$ and all sufficiently large m . Thus the consecutive ratios are log-convex and their maximum is at an endpoint. At $e=0$ the ratio is $O((\ln n)^3/n)$, by (8.15), and at $e = \lfloor m/4 \rfloor$ it is $n^{-1/2+o(1)}$. The series therefore decreases geometrically and

$$\sum_{1 \leq e < m/4} n^e R_{m,d}(e) = O((\ln n)^3/n). \quad (8.25)$$

Near-containment decorations. The passage from this per-cell estimate to the full skeleton sum is as follows. For a full-containment table $\mathbf{L}$, temporarily distinguish its endpoint cells, write $\text{cells}(\mathbf{L})$ for the resulting multiset, and let $\mathcal{S}(\mathbf{L})$ be all near-containment skeletons obtained by assigning to each such cell either $e = 0$ or $1 \leq e < m/4$. If $w(S)$ is its bare incidence and local weight, (8.21)–(8.25) give the nonnegative bound

$$\sum_{S \in \mathcal{S}(\mathbf{L})} w(S) \leq W(\mathbf{L}) \prod_{c \in \text{cells}(\mathbf{L})} \left(1 + \sum_{1 \leq e < m_c/4} n^e R_{m_c, d_c}(e) \right). \quad (8.25a)$$

Distinguishing and then forgetting identical typed cells is exactly the multinomial expansion, so there is no additional multiplicity. Since a high skeleton is a matching and has at most k_{co} cells, (8.25a) is at most $W(\mathbf{L})$ times

$$(1 + O((\ln n)^3/n))^{k_{\text{co}}} = \exp(O((\ln n)^2)). \quad (8.26)$$

Middle-strip residual factor. It remains to cover the middle strip, rather than silently importing an equitable-profile estimate. First expose a near-containment skeleton S . Its underlying set of high cells is a matching; write $\mathcal{M}(S)$ for that matching after multiplicities are forgotten, and let m_0 be the residual stub mass. Let $\mathcal{N}(S)$ be the residual event that there is no further near-containment cell. On $\mathcal{N}(S)$, every unexposed high cell is in the range $R_0 < j \leq 3a/4 + O(1)$. Define $E_{\text{mid}}(S)$ as the residual expectation of the product of the local factors truncated to this middle range, multiplied by $\mathbf{1}_{\mathcal{N}(S)}$. If $\tilde{g}$ denotes this truncated factor, then on $\mathcal{N}(S)$

$$\tilde{g}(r_{ab}) \leq 1 + \sum_{R_0 < j \leq 3a/4 + O(1)} g(j) \mathbf{1}_{\{r_{ab} \geq j\}}.$$

Expanding the product produces nonnegative monomials, each of which chooses distinct residual cells and at most one threshold in each selected cell. Apply Lemma 6.2 jointly to the entire threshold family before changing any residual restriction. Only afterwards do we discard $\mathbf{1}_{\mathcal{N}(S)}$: this can only increase each nonnegative monomial. The local factor remains truncated to the stated middle range, so the relaxation does not create a new charge for a newly admitted near-containment cell. If $m_0 \geq n/(\ln n)^6$, expand over distinct residual cells and their threshold demands. With $\theta_{ab} = e d_a d'_b / m_0 \leq e a^2 / m_0$, Lemma 6.2 is applied jointly before the constraints are dropped, and gives

$$\begin{aligned} E_{\text{mid}}(S) &\leq \sum_{\substack{D \text{ distinct residual cells} \\ R_0 < j_{ab} \leq 3a/4 + O(1)}} \prod_{ab \in D} g(j_{ab}) \Pr\{r_{ab} \geq j_{ab} \mid \forall ab \in D\} \\ &\leq \prod_{ab} \left(1 + \sum_{R_0 < j \leq 3a/4 + O(1)} g(j) \frac{\theta_{ab}^j}{j!} \right) \leq \exp(\Xi_4), \end{aligned} \quad (8.26a)$$

where

$$\Xi_4 \leq k_{\text{co}}^2 \sum_{a/2 < j \leq 3a/4 + O(1)} g(j) \frac{(e a^2 / m_0)^j}{j!}. \quad (8.27)$$

Put $\ell_n = \log_2 n$ and $j = x\ell_n$. Uniformly for $1 + o(1) \leq x \leq 3/2 + o(1)$,

$$\begin{aligned} \log_2 \left[k_{\text{co}}^2 g(j) \frac{(e a^2 / m_0)^j}{j!} \right] &\leq 2\ell_n + \frac{j^2}{2} - j \log_2 m_0 + O(j \log_2 a) \\ &\leq -c\ell_n^2. \end{aligned} \quad (8.28)$$

Indeed, the quadratic coefficient is $x^2/2 - x \leq -3/8 + o(1)$ on this interval. Hence $\Xi_4 = 2^{-\Omega(\ell_n^2)}$.

If instead $m_0 < n/(\ln n)^6$, leave the entire residual matching unexposed. For an upper bound, replace $\mathbf{1}_{N(S)}$ by 1 and the truncated local product by the full product $\prod_e g(r_e)$; both replacements enlarge the nonnegative integrand. Every residual degree is at most a , so every residual cell has $r_e \leq a$ and therefore $\binom{r_e}{2} \leq ((a-1)/2)r_e$. The near skeleton is a matching, so pointwise

$$\beta(\mathcal{M}(S) \cup H_{\text{res}}) \leq |E(H_{\text{res}})| \leq m_0/2, \quad \sum_e \binom{r_e}{2} \leq (a-1)m_0/2. \quad (8.29)$$

All remaining local and topological factors together are therefore at most $\exp(O(am_0)) = \exp(O(n/(\ln n)^5))$. Since $g \geq 1$ and $\beta \geq 0$, summing the residual matching probability first gives the explicit conditional bound

$$E_{\text{mid}}(S) \leq \mathbb{E}_{\text{res}} \left[\prod_e g(r_e) 2^{\beta(\mathcal{M}(S) \cup H_{\text{res}})} \right] \leq \exp(Cam_0). \quad (8.29a)$$

Combining (8.16), (8.25a)–(8.26a), and the two residual-mass branches now gives the global inequality

$$\sum_{\text{all high skeletons}} \text{bare weight} \leq e^{O((\ln n)^2)} \left(\sum_{\mathbf{L}} W(\mathbf{L}) \right) \max\{e^{\Xi_4}, e^{Cn/(\ln n)^5}\}. \quad (8.29b)$$

Therefore

$$\sum_{\text{all high skeletons}} \text{bare weight} \leq \exp\{O(\sqrt{n \ln n}) + O((\ln n)^2) + O(n/(\ln n)^5)\} = \exp\{o(n/(\ln n)^4)\}. \quad (8.30)$$

□

IX. All residual local and cycle attachments

Purpose of this section. Fixing a canonical high skeleton does not yet control the local multiplicity factors or the cycle-space factor of the residual overlap graph. The conditional quantity defined below collects precisely these residual contributions, subject to the residual cap and the prohibition on adding a further pair in a high cell. We prove a bound uniform over all feasible skeletons; Proposition 9.2 then combines that bound with the bare high-skeleton estimate from Section VIII.

Fix an actual canonical high skeleton $(\mathcal{M}, j)$ as in (8.2), and let $m_0 = n - J$ be the residual total degree. Its residual degrees are at most $u_{\max}$. Define

$$\mathcal{A}(\mathcal{M}, j) = \mathbb{E}_{\text{res}} \left[\left(\prod_e g(r'_e) \right) 2^{\beta(\mathcal{M} \cup H_{\text{res}})} \mathbf{1}_{\mathcal{E}(\mathcal{M}, j)} \right], \quad (9.1)$$

where $\mathcal{E}(\mathcal{M}, j)$ imposes the residual cap $r'_e \leq R_0$ and forbids a further pair in a cell of $\mathcal{M}$. By (6.4), (8.3), and the exact conditional law, the normalized moment is exactly

$$\sum_{(\mathcal{M}, j)} \pi(\mathcal{M}, j) \left(\prod_{e \in \mathcal{M}} g(j_e) \right) \mathcal{A}(\mathcal{M}, j). \quad (9.2)$$

The sum in (9.2) is over the finite set of feasible canonical high skeletons: typed matchings $\mathcal{M}$, allowed multiplicities $j_e > R_0$, and the induced nonnegative residual row and column degrees.

Lemma 9.1 (uniform attachment bound)

There is a deterministic sequence $\varepsilon_n \rightarrow 0$, independent of the canonical high skeleton, such that for every feasible skeleton and all sufficiently large n ,

$$\mathcal{A}(\mathcal{M}, j) \leq \exp \left\{ \varepsilon_n \frac{n}{(\ln n)^4} \right\}. \quad (9.3)$$

More precisely, for an absolute C , it is at most $\exp(C(\ln n)^8)$ when $m_0 \geq n/(\ln n)^6$ and at most $\exp(Cn/(\ln n)^5)$ otherwise. No lower bound on $\mathcal{A}$ is asserted or needed.

Proof: large residual degree

Suppose $m_0 \geq n/(\ln n)^6$. Since

$$u_{\max} = (2 + o(1)) \log_2 n, \quad \log_2 m_0 = \log_2 n - O(\log_2 \ln n), \quad (9.4)$$

we have $u_{\max} < 3 \log_2 m_0$ for all sufficiently large n . For a residual cell outside $\mathcal{M}$, put

$$\theta_{ab} = \frac{e d_a d'_b}{m_0}, \quad \Delta_x = g(x) - g(x-1), \quad (9.5)$$

and

$$\lambda_{ab}^{\text{loc}} = \sum_{x=3}^{R_0} \Delta_x \frac{\theta_{ab}^x}{x!}, \quad q_{ab} = \frac{\theta_{ab}^2}{2} + \lambda_{ab}^{\text{loc}}. \quad (9.6)$$

Set both quantities to zero on $\mathcal{M}$. Let $\theta_* = e u_{\max}^2 / m_0$. The ratio of consecutive majorants $g(x)\theta_*^x/x!$ is

$$\frac{2^x \theta_*}{x+1}, \quad (9.7)$$

and increases with x . Hence the sequence is log-convex. Its decreasing head is dominated by the $x = 3$ term. Consecutive values of (9.7) grow by the factor $2(x+1)/(x+2) \geq 8/5$, so only

$O(1)$ transition terms can have ratio between $1/2$ and 1 . Its increasing tail is dominated by its right endpoint, and

$$\log_2 \left(g(R_0) \frac{\theta_*^{R_0}}{R_0!} \right) \leq \frac{R_0^2}{2} - R_0 \log_2 m_0 + O(R_0 \log_2 u_{\max}) = -\Omega(R_0 \log n), \quad (9.8)$$

using $R_0 = u_{\max}/2$ and (9.4). Thus, uniformly in the cell,

$$\lambda_{ab}^{\text{loc}} \leq C \theta_{ab}^3, \quad q_{ab} \leq C \theta_{ab}^2. \quad (9.9)$$

Expand each capped local factor as

$$g(r'_e) = 1 + \sum_{x=3}^{R_0} \Delta_x \mathbf{1}_{\{r'_e \geq x\}}. \quad (9.10)$$

If an edge is selected in a fixed even subgraph, use instead

$$\mathbf{1}_{\{r'_e \geq 2\}} g(r'_e) = \mathbf{1}_{\{r'_e \geq 2\}} + \sum_{x=3}^{R_0} \Delta_x \mathbf{1}_{\{r'_e \geq x\}}. \quad (9.11)$$

In the product expansion, a selected cycle-space edge contributes either the base demand $\mathbf{1}_{\{r'_e \geq 2\}}$ or one increment term $\Delta_x \mathbf{1}_{\{r'_e \geq x\}}$, but not both in the same monomial. Thus a cell of multiplicity at least three is not charged simultaneously as a separate double-cell demand and as a higher-threshold local-factor demand. After fixing an even subgraph and expanding these alternatives, apply Lemma 6.2 jointly to all demanded cells. Every coefficient and event indicator is nonnegative, so the residual cap may be removed only after this joint bound is taken; doing so enlarges the event and preserves the stated upper-bound direction. This gives

$$\mathcal{A}(\mathcal{M}, j) \leq e^{\Lambda_{\text{loc}}} \sum_{F \in C_{\text{even}}(\mathcal{M}, R)} \prod_{e \in F \setminus \mathcal{M}} q_e, \quad \Lambda_{\text{loc}} = \sum_e \lambda_e^{\text{loc}}. \quad (9.12)$$

Here R is the finite potential residual support graph (all residual row–column cells), and $C_{\text{even}}(\mathcal{M}, R)$ is the finite family of edge subsets $F \subseteq E(\mathcal{M} \cup R)$ having even degree at every vertex; the edges of $\mathcal{M}$ have weight one in the displayed product.

By (9.9),

$$\sum_{a,b} \theta_{ab}^3 = \frac{e^3}{m_0^3} \left(\sum_a d_a^3 \right) \left(\sum_b (d'_b)^3 \right).$$

Every residual degree is at most $u_{\max}$, and both degree sums equal m_0 . Therefore $\sum_a d_a^3 \leq u_{\max}^2 m_0$ and $\sum_b (d'_b)^3 \leq u_{\max}^2 m_0$, which gives

$$\Lambda_{\text{loc}} \leq \frac{C}{m_0^3} \left(\sum_a d_a^3 \right) \left(\sum_b (d'_b)^3 \right) \leq C \frac{u_{\max}^4}{m_0}, \quad (9.13)$$

For a fixed row a , the quadratic bound in (9.9) also gives

$$\sum_b \theta_{ab}^2 = \frac{e^2 d_a^2}{m_0^2} \sum_b (d'_b)^2 \leq C \frac{u_{\max}^3}{m_0},$$

and the same calculation holds for every column. Hence

$$\max_a \sum_b q_{ab} \vee \max_b \sum_a q_{ab} \leq C \frac{u_{\max}^3}{m_0} =: \tau = o(1). \quad (9.14)$$

Cycle-kernel estimate. Every even edge set has a deterministic decomposition into edge-disjoint simple cycles, and the map from an even edge set to that chosen cycle family is injective because the original edge set is recovered by taking the union. We now enlarge the image of this map to all finite families of simple cycles, allowing cycles in the family to share edges. Since all cycle weights are nonnegative, this relaxation can only increase the sum. If $w(C) = \prod_{e \in C \setminus \mathcal{M}} q_e$, then, with $w(F) := \prod_{e \in F \setminus \mathcal{M}} q_e$,

$$\sum_{F \text{ even}} w(F) \leq \prod_C (1 + w(C)) \leq \exp \left(\sum_C w(C) \right),$$

which is precisely

$$\sum_{F \text{ even}} \prod_{e \in F \setminus \mathcal{M}} q_e \leq \exp \left\{ \sum_{C \text{ simple cycle}} \prod_{e \in C \setminus \mathcal{M}} q_e \right\}. \quad (9.15)$$

For cycles disjoint from $\mathcal{M}$, orient the cycle and distinguish one row vertex. We obtain an upper bound by enlarging the family of simple cycles to alternating walks: repeated vertices are permitted and the final return-to-start constraint is omitted. The resulting walks still have at least four residual edges, and successive row- and column-sum bounds by τ give the geometric series in (9.16):

$$\sum_{C: C \cap \mathcal{M} = \emptyset} \prod_{e \in C} q_e \leq \frac{n\tau^4}{1 - \tau^2}. \quad (9.16)$$

For a cycle meeting $\mathcal{M}$, distinguish and orient one matching edge. There are at most $2h$ choices. Cutting at every matching edge leaves nonempty residual walk segments. The residual-walk kernel has row-sum norm at most

$$b = \sum_{\ell \geq 1} \tau^\ell = \frac{\tau}{1 - \tau}, \quad (9.17)$$

while traversal of a matching edge is a partial permutation of norm one. After the initial distinguished matching edge there is no further factor h . Thus, if the relaxed cycle uses r matching edges, its total remaining weight is at most b^r . Therefore

$$\sum_{C: C \cap \mathcal{M} \neq \emptyset} \prod_{e \in C \setminus \mathcal{M}} q_e \leq 2h \sum_{r \geq 1} b^r \leq Ch\tau. \quad (9.18)$$

Every high cell uses more than $u_{\max}/2$ stubs, so $h < 2n/u_{\max}$. At the cutoff $m_0 \geq n/(\ln n)^6$,

$$\tau = O((\ln n)^9/n), \quad h\tau = O((\ln n)^8), \quad \frac{u_{\max}^4}{m_0} + n\tau^4 = o(1).$$

Combining (9.12)–(9.18),

$$\mathcal{A}(\mathcal{M}, j) \leq \exp \left[C \left\{ \frac{u_{\max}^4}{m_0} + n\tau^4 + h\tau \right\} \right] \leq \exp(C'(\ln n)^8) \quad (9.19)$$

at the cutoff $m_0 \geq n/(\ln n)^6$.

Proof: small residual degree

Suppose $m_0 < n/(\ln n)^6$. Since $\mathcal{M}$ is a matching, it is a forest, and adding a residual support edge increases cycle rank by at most one. Hence

$$\beta(\mathcal{M} \cup H_{\text{res}}) \leq |E(H_{\text{res}})| \leq m_0/2. \quad (9.20)$$

Also, pointwise,

$$\sum_e \binom{r'_e}{2} \leq \frac{u_{\max} - 1}{2} \sum_e r'_e = \frac{u_{\max} - 1}{2} m_0. \quad (9.21)$$

Equations (6.5), (9.20), and (9.21) bound the integrand in (9.1) by $2^{u_{\max} m_0/2}$. Therefore

$$\mathcal{A}(\mathcal{M}, j) \leq \exp(O(u_{\max} m_0)) \leq \exp(Cn/(\ln n)^5). \quad (9.22)$$

Both exponents in (9.19) and (9.22) are $o(n/(\ln n)^4)$. $\square$

Proposition 9.2 (normalized signed second moment)

For the exact midpoint profile,

$$\Lambda_n := \ln \frac{\mathbb{E}[(Z_{\mathbf{k}}^{\text{sgn}})^2]}{(\mathbb{E}Z_{\mathbf{k}}^{\text{sgn}})^2} = o(n/(\ln n)^4). \quad (9.23)$$

Proof

Equation (9.2) is an exact canonical decomposition. Write

$$w(\mathcal{M}, j) := \pi(\mathcal{M}, j) \prod_{e \in \mathcal{M}} g(j_e).$$

Lemma 8.3 sums all bare high skeletons by $\exp(o(n/(\ln n)^4))$, including the empty skeleton, all dense endpoint tables, every near-containment decoration, and the two middle-strip regimes. Lemma 9.1 bounds the conditional residual factor by the same exponential scale uniformly over every skeleton. Thus (9.2) gives the explicit estimate

$$\frac{\mathbb{E}[(Z_{\mathbf{k}}^{\text{sgn}})^2]}{(\mathbb{E}Z_{\mathbf{k}}^{\text{sgn}})^2} \leq \left(\sup_{(\mathcal{M}, j)} \mathcal{A}(\mathcal{M}, j) \right) \sum_{(\mathcal{M}, j)} w(\mathcal{M}, j) = \exp\{o(n/(\ln n)^4)\}.$$

This proves the upper bound in (9.23). The normalized second moment is at least one by variance, so its logarithm is nonnegative and (9.23) follows. $\square$

X. Rare-event amplification

The proof of Heckel (2025, Theorem 1) also begins with a rare second-moment seed and then uses vertex-exposure concentration of $\zeta(G_n)$ to obtain a high-probability statement. We use the same seed-to-typical strategic principle, but not that theorem as a black box: Lemma 10.2 proves the quantitative implication needed here for an arbitrary seed exponent Λ_n , and Lemma 10.1 supplies the simultaneous leftover colouring that controls the added parts.

Proposition 9.2 and (1.4) give the seed

$$\Pr\{Z_{\mathbf{k}}^{\text{sgn}} > 0\} \geq e^{-\Lambda_n}. \quad (10.1)$$

The ordinary-colouring concentration argument motivating this amplification was communicated by Noga Alon and is presented by Scott (2017, Theorem 1). Lemmas 10.1 and 10.2 prove the precise simultaneous-leftover and rare-seed forms required here.

By the observation after (5.19), this implies

$$\Pr\{\zeta(G_n) \leq k_{\text{co}}\} \geq e^{-\Lambda_n}. \quad (10.2)$$

We now prove the amplification needed to turn this possibly rare event into a typical one.

Lemma 10.1 (simultaneous leftover colouring)

There is an absolute C such that, with probability $1 - o(1)$, every $S \subseteq [n]$ satisfies

$$\chi(G_n[S]) \leq C \frac{|S|}{\ln n} + n^{1/3}. \quad (10.3)$$

Proof

Let H be the complement of G_n . Put $u_0 = \lceil n^{1/4} \rceil$. For any fixed u_0 -set, (1.5) gives probability $\exp(-\Omega(u_0^2))$ that its H -edge density is below $1/4$. There are at most $\binom{n}{u_0}$ such sets, and for an absolute $c > 0$,

$$\binom{n}{u_0} e^{-cu_0^2} \leq \exp\{u_0 \ln(en/u_0) - cu_0^2\} = o(1).$$

Thus a union bound shows that, with probability $1 - o(1)$, every u_0 -set has density at least $1/4$. By averaging over its u_0 -subsets, the same is then true of every larger set.

Start with any vertex set of size at least $n^{1/3}$. While its current size is at least u_0 , choose a vertex of maximum H -degree and replace the current set by its H -neighbourhood there. Density at least $1/4$ implies that this neighbourhood has size at least one quarter of the current size, up to an additive one. If s_t is the current size after t choices, then

$$s_{t+1} \geq (s_t - 1)/4, \quad s_t \geq 4^{-t} s_0 - 1/3. \quad (10.3a)$$

The chosen vertices form a clique in H . Starting from any $\lceil n^{1/3} \rceil$ -subset, (10.3a) stays above $n^{1/4}$ for at least $\lfloor \ln n / (13 \ln 4) \rfloor$ steps. Thus it constructs at least $c \ln n$ vertices for an absolute $c > 0$. They are an independent set in G_n .

For an arbitrary S , repeatedly remove such independent sets and give each a new colour until fewer than $n^{1/3}$ vertices remain; colour the rest singly. Every removed set has at least $c \ln n$ vertices, so the total number of colours is at most

$$\left\lceil \frac{|S|}{c \ln n} \right\rceil + n^{1/3}.$$

After enlarging the absolute constant, this is (10.3), simultaneously for every S . $\square$

Lemma 10.2 (amplification from a seed)

Suppose deterministic $k_n, \Lambda_n \geq 0$ satisfy

$$\Pr\{\zeta(G_n) \leq k_n\} \geq e^{-\Lambda_n}. \quad (10.4)$$

There is a deterministic $\varepsilon_n = o(1)$, independent of k_n, Λ_n, r , such that, uniformly for every deterministic choice $r = r(n) > 0$,

$$\Pr\left\{\zeta(G_n) > k_n + C\left(\frac{\sqrt{n\Lambda_n} + \sqrt{nr}}{\ln n} + n^{1/3} + 1\right)\right\} \leq e^{-r} + \varepsilon_n. \quad (10.5)$$

Proof

For an induced set W , let $\zeta(W) = \zeta(G_n[W])$, and define

$$S_k = \max\{|W| : \zeta(W) \leq k_n\}. \quad (10.6)$$

Expose the random graph in $n - 1$ independent vertex blocks, where block v contains the edges from v to earlier vertices. Changing one block changes S_k by at most one: after deletion of the affected vertex, every feasible induced set loses at most one vertex and the two configurations agree on the remainder. Therefore (1.3) applies.

Since $S_k = n$ exactly when $\zeta(G_n) \leq k_n$, (10.4) and the upper one-sided bounded-differences tail give

$$e^{-\Lambda_n} \leq \Pr\{S_k - \mathbb{E}S_k \geq n - \mathbb{E}S_k\} \leq \exp\left\{-\frac{2(n - \mathbb{E}S_k)^2}{n - 1}\right\}.$$

Taking logarithms and rearranging imply

$$n - \mathbb{E}S_k \leq \sqrt{(n - 1)\Lambda_n/2}. \quad (10.7)$$

The lower tail with radius $\sqrt{(n - 1)r/2}$ gives, except with probability e^{-r} ,

$$n - S_k \leq \sqrt{(n - 1)\Lambda_n/2} + \sqrt{(n - 1)r/2}. \quad (10.8)$$

Choose a maximizing set W and put $V_{\text{left}} = [n] \setminus W$. Concatenating a k_n -part cocolouring of $G[W]$ with an ordinary colouring of $G[V_{\text{left}}]$ gives

$$\zeta(G_n) \leq k_n + \chi(G_n[V_{\text{left}}]). \quad (10.9)$$

Intersect (10.8) with the simultaneous event in Lemma 10.1 and apply (10.3). Since $|V_{\text{left}}| = n - S_k$, equations (10.3), (10.8), and (10.9) give exactly the number of additional parts displayed in (10.5). The simultaneous event's failure probability is the same deterministic $\varepsilon_n = o(1)$ for every choice of the three deterministic parameters. This proves (10.5). $\square$

Apply Lemma 10.2 to (10.2). Put

$$B_n = \frac{n}{(\ln n)^4}, \quad r_n = \sqrt{B_n}. \quad (10.10)$$

Then $r_n \rightarrow \infty$, and Proposition 9.2 gives

$$\frac{\sqrt{n\Lambda_n}}{\ln n} = o(n/(\ln n)^3), \quad \frac{\sqrt{nr_n}}{\ln n} = o(n/(\ln n)^3), \quad n^{1/3} = o(n/(\ln n)^3). \quad (10.11)$$

Define the deterministic sequence

$$a_n = C \left(\frac{\sqrt{n\Lambda_n} + \sqrt{nr_n}}{\ln n} + n^{1/3} + 1 \right). \quad (10.12)$$

Then (10.11) and Lemma 10.2 give

$$a_n = o(n/(\ln n)^3), \quad \Pr\{\zeta(G_n) > k_{\text{co}} + a_n\} \leq e^{-r_n} + \varepsilon_n \rightarrow 0. \quad (10.13)$$

XI. Completion of the proof

Intersect $\{\chi(G_n) > k_{\chi}^-\}$, whose probability tends to one by (4.5), with $\{\zeta(G_n) \leq k_{\text{co}} + a_n\}$, whose probability tends to one by (10.13). No independence is required; a union bound suffices. Equation (5.20) then gives, with high probability,

$$\chi(G_n) - \zeta(G_n) \geq \left(\frac{(\ln 2)^2 \gamma_4}{16} - o(1) \right) \frac{n}{(\ln n)^3}. \quad (11.1)$$

For every sufficiently large n , the right side is at least

$$c_* \frac{n}{(\ln n)^3}, \quad c_* := \frac{(\ln 2)^2 \gamma_4}{32} = \frac{(\ln 2)^2}{32} \ln \frac{200}{153} > 0. \quad (11.2)$$

This proves (0.1). Since $n/(\ln n)^3$ tends to infinity, for every fixed K ,

$$\Pr\{\chi(G_n) - \zeta(G_n) \geq K\} \rightarrow 1. \quad (11.3)$$

Every estimate used above is uniform for $\delta \in [0, 1)$ and for the exact tangent-rounded finite- n profile. The conclusion therefore holds along all sufficiently large integers, including sequences approaching either side of every independence-number jump. It is a high-probability statement, not a density-subset or subsequence result. $\blacksquare$

XII. Dependency and provenance note

The proof is logically ordered as follows. The elementary phase expansion and exact first-moment variational calculation construct the midpoint profile and prove its complete signed first-moment margin. That margin and elementary independence-set estimates prove the full partial-diagonal sum. The partial-diagonal sum feeds the dense four-type transportation estimate. Independently, the exact sign sum and the configuration model prove the uniform residual-attachment estimate. These two estimates combine only after both are established, yielding the normalized second moment. Paley–Zygmund and the proved leftover amplification are applied last. Thus there is no circular use of a second-moment conclusion in the profile construction. In particular, no ordinary-colouring second-moment theorem is invoked at the signed midpoint, where the ordinary first moment is exponentially small.

The manuscript was consolidated from a larger research dossier containing separate derivations and adversarial checks of the exceptional regime, the four-size profile, overlap enumeration, residual attachments, and concentration amplification. No step of the proof above depends on those internal cross-references. The cited historical sources confirm the definition, the original divergence question, and the standard first-order chromatic asymptotic; none supplies a step of the proof above.

The two Heckel papers have a different role. Heckel’s 2024 theorem gives the anti-concentration obstruction and states the conjectural $n/(\ln n)^3$ scale (Heckel, 2024, Theorem 1 and Conjecture 4). Heckel’s 2025 paper proves the gap on the phase window displayed in the Introduction and supplies the signed 2^k identity, the overlap-sensitive sign bound, and the rare-seed amplification strategy (Heckel, 2025, Theorem 1 and Propositions 5–6). Those are essential strategic antecedents. No numbered theorem from either paper is nevertheless used as an unproved implication in the candidate proof above: the 2^k identity is rederived in Section V, the pair-sign inequality is replaced by the exact component identity in Section VI, the phase-restricted tame-profile estimates are replaced by Sections VII–IX, and the amplification is proved in Section X. Thus the claims of full-phase uniformity and the explicit constant are claims of this manuscript, not claims attributed to the cited papers.

XIII. Status, reproducibility, and responsible AI use

Candidate status. This manuscript presents a candidate proof and has not undergone external peer review. Its appearance as a preprint or in a public repository should not be read as publication, priority verification, community acceptance, or independent confirmation of the theorem. The proof is written to be audited line by line, and any remaining error would affect the claimed conclusion.

Research artifacts and formalization. The public research dossier, source code, independent computational checks, and release materials are available at <https://github.com/SamPetkov/Erdos>. The accompanying Lean 4 development is available at <https://github.com/SamPetkov/Erdos/tree/main/625/formalization>. As of 12 July 2026 the Lean development is ongoing and partial: it certifies a growing collection of definitions and proof components, but it is not yet a machine-checked proof of the complete theorem stated here. Numerical and symbolic checks in the dossier are validation evidence, not substitutes for either the written argument or

a complete formal proof.

Ethics and AI assistance. The proof and manuscript were completed with immense assistance from ChatGPT Sol Ultra 5.6. That assistance included proof development, counterexample-oriented auditing, symbolic and numerical checks, exposition, and preparation of the manuscript. Independent verification runs were performed with ChatGPT 5.6 Sol Pro. Samuil Petkov directed the work, selected the claims and arguments to include, reviewed the resulting manuscript, and takes full responsibility for the accuracy, integrity, citations, and final submission. Aristotle, the automated theorem-proving system of Achim et al. (2025), was used only for quarantined atomic Lean proof search. Every returned candidate was locally ported, reviewed, and rebuilt before acceptance into the formalization; Aristotle was not treated as having proved this paper or its main theorem. The AI systems are not listed as authors.

Declarations. No external funding was received for this work. The author declares that there are no competing interests. No new empirical dataset is introduced; reproducibility materials are linked above.

XIV. References

- Tudor Achim, Alex Best, Alberto Bietti, Kevin Der, Mathis Fédérico, Sergei Gukov, Daniel Halpern-Leistner, Kirsten Henningsgard, Yury Kudryashov, Alexander Meiburg, Martin Michelsen, Riley Patterson, Eric Rodriguez, Laura Scharff, Vikram Shanker, Vladmir Sicca, Hari Sowrirajan, Aidan Swope, Matyas Tamas, Vlad Tenev, Jonathan Thomm, Harold Williams, and Lawrence Wu. Aristotle: IMO-level automated theorem proving, 2025. URL <https://doi.org/10.48550/arXiv.2510.01346>. arXiv:2510.01346; version 2, revised 10 October 2025; DOI 10.48550/arXiv.2510.01346.
- Thomas F. Bloom. Erdős problem #625. <https://www.erdosproblems.com/625>, 2026. Accessed 14 July 2026.
- Béla Bollobás. The chromatic number of random graphs. *Combinatorica*, 8(1):49–55, 1988. doi: 10.1007/BF02122551. URL <https://doi.org/10.1007/BF02122551>.
- Paul Erdős and John Gimbel. Some problems and results in cochromatic theory. In John Gimbel, John W. Kennedy, and Louis V. Quintas, editors, *Quo Vadis, Graph Theory?*, volume 55 of *Annals of Discrete Mathematics*, pages 261–264. North-Holland/Elsevier, 1993. doi: 10.1016/S0167-5060(08)70393-5. URL [https://doi.org/10.1016/S0167-5060\(08\)70393-5](https://doi.org/10.1016/S0167-5060(08)70393-5).
- John Gimbel. Some of my favorite coloring problems for graphs and digraphs. In Ralucca Gera, Stephen T. Hedetniemi, and Craig Larson, editors, *Graph Theory: Favorite Conjectures and Open Problems – 1*, Problem Books in Mathematics, pages 95–108. Springer, Cham, 2016. ISBN 978-3-319-31940-7. doi: 10.1007/978-3-319-31940-7_7. URL https://doi.org/10.1007/978-3-319-31940-7_7.
- Annika Heckel. On a question of erdős and gimbel on the cochromatic number. *The Electronic Journal of Combinatorics*, 31(4):P4.72, 2024. doi: 10.37236/13346. URL <https://doi.org/10.37236/13346>.

Annika Heckel. The difference between the chromatic and the cochromatic number of a random graph, 2025. URL <https://arxiv.org/abs/2409.17614>. Version 2, revised 19 February 2025; first submitted in 2024.

Svante Janson, Tomasz Łuczak, and Andrzej Ruciński. *Random Graphs*. John Wiley & Sons, New York, 2000. ISBN 0-471-17541-2.

Colin McDiarmid. On the method of bounded differences. In Johannes Siemons, editor, *Surveys in Combinatorics, 1989*, volume 141 of *London Mathematical Society Lecture Note Series*, pages 148–188. Cambridge University Press, 1989. doi: 10.1017/CBO9781107359949.008. URL <https://doi.org/10.1017/CBO9781107359949.008>.

Alex Scott. On the concentration of the chromatic number of random graphs, 2017. URL <https://arxiv.org/abs/0806.0178>. Version 2; first submitted in 2008.

Raphael Steiner. On the difference between the chromatic and cochromatic number, 2024. URL <https://arxiv.org/abs/2408.02400>. Version 2, revised 20 August 2024.