

OBLIGATORY TRIPLE SYSTEMS: A SELF-CONTAINED PROOF OF ERDŐS PROBLEM 593

SAMUIL PEKOV

ABSTRACT. We prove the exact characterisation of finite obligatory triple systems. After isolated vertices are removed, obligatoriness is equivalent both to membership in the class generated by private-vertex expansions of finite bipartite graphs under finite disjoint unions and one-point amalgamations, and to the intrinsic conjunction of linearity, one incident Levi bridge at every hyperedge-node, and evenness of every Berge cycle. The argument is valid in ZFC.

Bibliographic note. A contemporaneous preprint by Eric Li claims resolutions of Erdős Problems 593 and 1177 [1]. It is cited solely for bibliographic completeness; no claim from that preprint is used as a premise or presented here as independently verified.

MAIN THEOREM

Theorem A: exact characterisation.

For every finite triple system F , the following are equivalent:

1. F is obligatory.
2. $F \in \mathcal{B}$.
3. F° is linear, every hyperedge-node of $I(F^\circ)$ is incident with a bridge, and every Berge cycle of F° has even length.

Here F° is obtained by deleting all isolated vertices, and $\mathcal{B}$ is the smallest class containing every private-vertex expansion J^+ of a finite bipartite graph, every finite edgeless triple system, and closed under finite disjoint unions and one-point amalgamations.

The proof is arranged as

$$(2) \implies (1), \quad (2) \iff (3), \quad \neg(3) \implies \neg(1).$$

All arguments are in ZFC, and all embeddings are non-induced.

Proof dependencies. The only non-elementary external results used as black boxes are the de Bruijn–Erdős compactness theorem for finite graph colourings [2] and the pair case of the Erdős–Rado partition theorem [3], namely

$$(2^{\aleph_0})^+ \longrightarrow (\aleph_1)_{\aleph_0}^2.$$

Every other ingredient required for the classification is proved below.

Background. The problem was posed by Erdős and is catalogued as Erdős Problem 593 [4, 5]. Background on obligatory subsystems, large-chromatic set systems, and high-girth constructions may be found in [6–14].

1. PRELIMINARY LEMMAS

Lemma 1.1: isolated-vertex reduction.

Let F be a finite triple system and let F° be obtained by deleting all isolated vertices. Then

$$F \text{ is obligatory} \iff F^\circ \text{ is obligatory,}$$

and

$$F \in \mathcal{B} \iff F^\circ \in \mathcal{B}.$$

Proof. If F° is obligatory and H has uncountable chromatic number, then H has infinitely many vertices. After embedding the finite system F° , map the isolated vertices of F injectively to unused host vertices. Containment is non-induced, so additional host edges are irrelevant. The converse follows because F° is a subhypergraph of F .

For membership in $\mathcal{B}$, the implication $F^\circ \in \mathcal{B} \Rightarrow F \in \mathcal{B}$ follows by taking the disjoint union with a finite edgeless system.

For the converse, use structural induction on a construction of F in $\mathcal{B}$. If $F = J^+$, delete the isolated graph vertices of J to obtain J_0 ; then $(J^+)^\circ = J_0^+ \in \mathcal{B}$. If F is edgeless, its reduction is the empty edgeless system. Reduction commutes with disjoint union.

Suppose finally that F is a one-point amalgamation of F_0 and F_1 , identifying $x_i \in V(F_i)$. By induction $F_0^\circ, F_1^\circ \in \mathcal{B}$. If both x_i are nonisolated in their factors, then F° is their one-point amalgamation at those vertices. If exactly one x_i is nonisolated, the isolated side disappears at the identified point and F° is the disjoint union of the two reductions. The same disjoint-union description applies when both x_i are isolated, because the identified point is deleted. Hence $F^\circ \in \mathcal{B}$ in every case. $\square$

Thus isolated vertices may be suppressed throughout the substantive argument and restored at the end.

Lemma 1.2: finite deletion.

If $\chi(H) > \aleph_0$ and $S \subseteq V(H)$ is finite, then

$$\chi(H - S) > \aleph_0.$$

Proof. Otherwise a countable colouring of $H - S$, together with finitely many fresh colours for S , would colour H countably. $\square$

Lemma 1.3: two elementary colouring facts.

The following facts will be used repeatedly.

1. If $E(H) = E_1 \cup \dots \cup E_r$ and every $(V(H), E_i)$ is countably chromatic, then H is countably chromatic.
2. If a graph G admits an orientation in which every vertex has outdegree at most $d < \omega$, then $\chi(G) \leq 2d + 1$.

Proof. For the first assertion, take the product of the r countable colourings.

For the second, every finite subgraph has at most $d|V|$ edges and therefore a vertex of degree at most $2d$. It is $(2d + 1)$ -colourable by greedy deletion. The de Bruijn–Erdős compactness theorem [2] then gives a $(2d + 1)$ -colouring of all of G . $\square$

Lemma 1.4: closure-chain lemma.

Let κ be an infinite cardinal and let

$$\Phi : [\kappa]^r \longrightarrow [\kappa]^{<\omega}$$

have uniformly finite values. There is an increasing continuous chain

$$\emptyset = M_0 \subseteq M_1 \subseteq \dots \subseteq M_i \subseteq \dots \quad (i < \text{cf } \kappa)$$

such that

$$\bigcup_{i < \text{cf } \kappa} M_i = \kappa, \quad |M_i| < \kappa,$$

and every M_i is Φ -closed:

$$a \in [M_i]^r \implies \Phi(a) \subseteq M_i.$$

Proof. Choose sets $A_i \subseteq \kappa$, each of cardinality below κ , whose union is κ . At a successor stage close $M_i \cup A_i$ under Φ , iterating the operation ω times. Since Φ is finitary and finite-valued, this does not increase an infinite cardinal below κ . At a limit stage take the union. A union of an increasing chain of sets closed under a finitary operation remains closed; and because the stage is below $\text{cf } \kappa$, its cardinality remains below κ . $\square$

Write

$$I_i = M_{i+1} \setminus M_i.$$

Then $(I_i)_{i < \text{cf } \kappa}$ partitions κ , and $M_i = \bigcup_{j < i} I_j$.

2. THE GRAPH LEMMA NEEDED FOR THE EXPANSION ATOM

Lemma 2.1: uncountable chromatic number forces complete bipartite graphs.

For every positive integer n , every graph of uncountable chromatic number contains $K_{n,n}$.

Proof. Suppose otherwise, and choose a $K_{n,n}$ -free graph G of uncountable chromatic number whose vertex cardinality κ is minimal.

For $A \in [\kappa]^n$, let

$$N(A) = \{v \in \kappa : v \text{ is adjacent to every member of } A\}.$$

Because G is $K_{n,n}$ -free, $|N(A)| < n$. Apply Lemma 1.4 to $A \mapsto N(A)$, and let M_i, I_i be the resulting chain and layers.

By minimality of κ , every induced graph $G[I_i]$ is countably chromatic. Let $G^\times$ consist of the edges joining distinct layers, and orient each such edge from the later endpoint toward the earlier endpoint. If $v \in I_i$ had n earlier neighbours, those neighbours would form an n -set $A \subseteq M_i$, and closure would give

$$v \in N(A) \subseteq M_i,$$

contrary to $v \in I_i$. Thus the orientation of $G^\times$ has outdegree at most $n - 1$. By Lemma 1.3, $G^\times$ is finitely colourable.

Colour each $G[I_i]$ with colours in ω , reusing the same palette on all layers, and combine this with a finite proper colouring of $G^\times$. The resulting countable product colouring is proper on all of G , a contradiction. $\square$

3. PRIVATE-VERTEX EXPANSIONS OF BIPARTITE GRAPHS ARE OBLIGATORY

Lemma 3.1: rainbow bipartite submatrices.

For all positive integers n, t , there exists $q = q(n, t)$ with the following property. Suppose the edges of $K_{q,q}$ are coloured, with an arbitrary colour set, so that at every vertex each individual colour occurs on fewer than t incident edges. Then there are n -element subsets X' and Y' of the two vertex classes such that all n^2 edges between X' and Y' have distinct colours.

Proof. Choose independent uniformly random n -subsets X' and Y' .

Let m_γ be the number of edges of colour γ . The local bound gives

$$m_\gamma \leq (t-1)q.$$

The number of pairs of same-coloured edges sharing a vertex is at most $(t-1)q^2$. The total number of pairs of same-coloured edges is at most

$$\sum_\gamma \binom{m_\gamma}{2} \leq \frac{1}{2} (\max_\gamma m_\gamma) \sum_\gamma m_\gamma \leq \frac{1}{2} (t-1)q^3.$$

For $q \geq 2n$, a fixed adjacent pair of edges is selected with probability at most $2n^3/q^3$, while a fixed disjoint pair is selected with probability at most $4n^4/q^4$. Hence the expected number of repeated colours in the selected $K_{n,n}$ is at most

$$\frac{2(t-1)n^3}{q} + \frac{2(t-1)n^4}{q}.$$

This is below 1 for sufficiently large q . Some choice of X', Y' therefore has no repeated colour.

□

Proposition 3.2: the complete bipartite expansion atom.

For every positive integer n , the triple system $K_{n,n}^+$ is obligatory.

Proof. Suppose not. Choose a $K_{n,n}^+$ -free triple system

$$H = (\kappa, E)$$

with $\chi(H) > \aleph_0$ and with κ minimal.

For a pair xy , write

$$d_H(xy) = |\{z : \{x, y, z\} \in E\}|.$$

Set $t = 3n^2 + 1$ and define a graph R on κ by

$$xy \in E(R) \iff d_H(xy) \geq t.$$

Claim 3.2.1: R is countably chromatic. Otherwise Lemma 2.1 gives a $K_{n,n}$ in R , with core vertex set C , $|C| = 2n$. For each of its n^2 graph edges xy , choose a third vertex p_{xy} with

$$\{x, y, p_{xy}\} \in E,$$

avoiding C and all previously chosen private vertices. At every stage fewer than $2n + n^2 < t$ vertices are forbidden, while xy has at least t possible third vertices. Thus the p_{xy} can be chosen distinct and outside C , producing a copy of $K_{n,n}^+$, a contradiction. □

Let

$$E_1 = \{e \in E : \text{some pair contained in } e \text{ is an edge of } R\}, \quad E_2 = E \setminus E_1.$$

A proper colouring of R is a proper colouring of (κ, E_1) , since every E_1 -edge contains an R -edge. Hence $\chi(\kappa, E_1) \leq \aleph_0$, and therefore

$$\chi(\kappa, E_2) > \aleph_0. \tag{3.1}$$

For a pair xy , define

$$\Phi(xy) = \begin{cases} \{z : \{x, y, z\} \in E\}, & d_H(xy) < t, \\ \emptyset, & d_H(xy) \geq t. \end{cases}$$

Apply Lemma 1.4 to Φ , obtaining M_i, I_i .

Every $e \in E_2$ has at least two vertices in its highest layer. Indeed, suppose the highest layer met e only in $z \in I_i$. Then the other pair $xy = e \setminus \{z\}$ lies in M_i . Since $e \in E_2$, $xy \notin R$, so $d_H(xy) < t$. Closure gives $z \in \Phi(xy) \subseteq M_i$, a contradiction.

Thus every E_2 -edge is either contained in one layer, or has exactly two vertices in its highest layer I_i and its third vertex in M_i . Let E_i^* be the E_2 -edges contained in I_i . Since $|I_i| < \kappa$ and H is $K_{n,n}^+$ -free, minimality of κ gives $\chi(I_i, E_i^*) \leq \aleph_0$. Consequently

$$E^* = \bigcup_i E_i^*$$

is countably chromatic. By (3.1), the remaining crossing-edge system

$$E^{**} = E_2 \setminus E^*$$

has uncountable chromatic number.

For each i , define a graph G_i on I_i by declaring $xy \in E(G_i)$ when there is a $\beta \in M_i$ such that $\{x, y, \beta\} \in E^{**}$. If every G_i were countably chromatic, colour each I_i properly in G_i , using the same countable palette on every layer. Every edge of E^{**} would then have differently coloured top-layer vertices, giving a countable colouring of (κ, E^{**}) , a contradiction. Hence some G_i is uncountably chromatic.

Choose $q = q(n, t)$ from Lemma 3.1. Lemma 2.1 gives a $K_{q,q}$ in G_i , with sides X, Y . For every $xy \in X \times Y$, choose

$$\beta_{xy} \in M_i$$

such that $\{x, y, \beta_{xy}\} \in E^{**}$, and regard β_{xy} as the colour of the edge xy .

This edge-colouring is locally $(t-1)$ -bounded. For example, if for fixed x and fixed β there were t different vertices y with $\beta_{xy} = \beta$, then the pair $\{x, \beta\}$ would have codegree at least t . It would belong to R , forcing the triples $\{x, y, \beta\}$ into E_1 , contrary to their membership in $E^{**} \subseteq E_2$. The same argument applies at vertices of Y .

Lemma 3.1 gives n -sets $X' \subseteq X$, $Y' \subseteq Y$ for which the β_{xy} , $x \in X'$, $y \in Y'$, are all distinct. They lie in M_i , whereas $X' \cup Y' \subseteq I_i$, so none is a core vertex. The triples

$$\{x, y, \beta_{xy}\}, \quad x \in X', \quad y \in Y',$$

form $K_{n,n}^+$, the final contradiction. $\square$

Corollary 3.3: every finite bipartite expansion is obligatory.

If J is any finite bipartite graph, then J^+ is obligatory.

Proof. Embed the two vertex classes of J in the two classes of some $K_{n,n}$. Then J^+ is a subhypergraph of $K_{n,n}^+$. Obligatoriness is downward closed under non-induced containment: a copy of the larger finite system contains a copy of the smaller one. Apply Proposition 3.2. $\square$

4. CLOSURE OF OBLIGATORINESS UNDER THE OPERATIONS DEFINING $\mathcal{B}$

Lemma 4.1: disjoint-union closure.

The class of finite obligatory triple systems is closed under finite disjoint unions.

Proof. Let F_1, F_2 be obligatory and let H be uncountably chromatic. First find a copy of F_1 . Delete its finite vertex set. By Lemma 1.2 the remaining triple system is still uncountably chromatic, so it contains F_2 . The two copies are disjoint. Iteration proves the finite case. $\square$

Definition 4.2: rooted abundance.

Let (F, r) be a finite rooted triple system with $|V(F)| > 1$. For a triple system H and $m \geq 1$, let $R_m(F, r; H)$ be the set of vertices v for which there are m rooted copies of F , all sending r to v , whose off-root vertex sets are pairwise disjoint.

Lemma 4.3: rooted-abundance lemma.

If (F, r) is finite and F is obligatory, and if $\chi(H) > \aleph_0$, then

$$\chi(H[R_m(F, r; H)]) > \aleph_0.$$

Proof. Put

$$B = V(H) \setminus R_m(F, r; H).$$

For each $v \in B$, consider the family of off-root vertex sets of all copies of F rooted at v . Its matching number is below m . Choose a maximal pairwise disjoint subfamily and let S_v be the union of its members. Then

$$|S_v| \leq (m-1)(|V(F)|-1) =: D,$$

and S_v meets the off-root part of every rooted copy at v .

Form a graph D_B on B by joining v to each member of $S_v \cap B$. If an edge is generated in both directions, assign it to one of its generators, and orient it away from that generator. Every vertex has outdegree at most D , so Lemma 1.3 gives a finite proper colouring of D_B .

Let A be one colour class. Then

$$S_v \cap A = \emptyset \quad (v \in A).$$

Consequently $H[A]$ contains no copy of F . Indeed, if φ were such a copy and $v = \varphi(r)$, its off-root part would have to meet S_v although both are subsets of A . Since F is obligatory, every F -free triple system is countably chromatic. Thus each of the finitely many colour classes of D_B induces a countably chromatic subsystem, and $H[B]$ is countably chromatic.

If $H[R_m(F, r; H)]$ were countably chromatic as well, tagging the two sides would give a countable colouring of H . This contradicts $\chi(H) > \aleph_0$. $\square$

Proposition 4.4: one-point amalgamation closure.

The class of finite obligatory triple systems is closed under one-point amalgamations.

Proof. Let F_1, F_2 be obligatory, with selected vertices r_1, r_2 , and let F be their one-point amalgamation. If one factor consists only of its selected vertex, the assertion is immediate. Otherwise set

$$m = |V(F_2)|.$$

By Lemma 4.3, the set

$$R = R_m(F_1, r_1; H)$$

induces an uncountably chromatic subsystem of every uncountably chromatic H . Hence $H[R]$ contains a copy of F_2 . Let v be the image of r_2 .

At v there are m rooted copies of F_1 with pairwise disjoint off-root sets. The copy of F_2 has only $m-1$ vertices other than v , so at most $m-1$ of those petals meet it. One rooted F_1 -copy avoids the F_2 -copy outside v . Their union is the required one-point amalgamation. $\square$

Corollary 3.3, Lemma 4.1, Proposition 4.4, and the trivial edgeless case prove

$$(2) \implies (1). \tag{4.1}$$

5. EQUIVALENCE OF THE CONSTRUCTIVE AND INTRINSIC DESCRIPTIONS

We prove (2) $\iff$ (3). Isolated vertices may be ignored by Lemma 1.1.

Proposition 5.1: the intrinsic conditions are preserved by the generators and operations.

Every generator of $\mathcal{B}$ satisfies the intrinsic conditions, and the conditions are preserved under finite disjoint unions and one-point amalgamations.

Proof. For a bipartite graph J , the expansion J^+ is linear. Every hyperedge-node is joined to its private point by a bridge. A Berge cycle cannot use a private point, so the Berge cycles of J^+ are precisely the ordinary graph cycles of J , with the same lengths; these are even. Edgeless systems satisfy the conditions vacuously, and disjoint union plainly preserves them.

Consider a one-point amalgamation. Hyperedges belonging to different factors intersect only in the identified point, so linearity is preserved. Its Levi graph is the vertex-sum of the two Levi graphs at a point-node. Every simple cycle in a vertex-sum lies wholly in one factor: a cycle crossing into the other factor would have to pass through the cut vertex twice. Consequently bridges in the factors remain bridges, and every Berge cycle lies in one factor and retains its parity.

□

Thus

$$(2) \implies (3). \quad (5.1)$$

Proposition 5.2: bridge-block decomposition.

Suppose F is finite, has no isolated vertices, and satisfies the three intrinsic conditions. Then $F \in \mathcal{B}$.

Proof. Put $L = I(F)$. Delete all bridges of L , and let $\mathcal{C}$ be the set of connected components of the resulting graph.

Consider a hyperedge-node e . Since e is incident with a bridge and has degree three, it has at most two nonbridge incidences. It cannot have exactly one: a nonbridge incidence lies on a cycle, and that cycle must leave e through a second nonbridge incidence. Therefore every hyperedge-node has either no nonbridge incidences or exactly two.

The nontrivial components. Let $C \in \mathcal{C}$ contain a nonbridge incidence. Every hyperedge-node $e \in C$ has exactly two point-neighbours in C . Contract each such e to an ordinary graph edge between those two point-neighbours. This produces a finite graph J_C .

The graph J_C is simple: parallel edges would correspond to two hyperedges of F sharing two points, contrary to linearity. Every cycle in J_C gives a Berge cycle of the same length in F . Hence J_C has no odd cycle and is bipartite.

Each hyperedge-node $e \in C$ has a third point-neighbour p_e across a bridge. These third points satisfy:

1. $p_e \notin C$; otherwise a path in C from p_e to e , together with the incidence ep_e , would place that incidence on a cycle.
2. If $e \neq f$ lie in C , then $p_e \neq p_f$; otherwise a path in C from e to f , together with ep_e and p_ff , would place those bridge incidences on a cycle.

Thus the hyperedges belonging to C , together with their third points p_e , form exactly the private-vertex expansion J_C^+ .

Components consisting of one hyperedge-node. If a hyperedge-node e has no nonbridge incidence, it is an isolated component after the bridges are deleted. Its single triple is the expansion of a one-edge bipartite graph. Thus every hyperedge of F belongs to a piece P_C isomorphic to J^+ for a finite bipartite graph J .

Quotient forest. Contract every component of L after bridge deletion. The original bridges become edges of a graph T . This graph is a forest. Indeed, a cycle in the quotient would lift, by paths within the contracted components, to a cycle in L containing one of the purported bridges.

Components of T that contain hyperedge-nodes will be called active. For an active component C , the piece P_C consists of the hyperedges represented in C , all point-nodes of C , and the point endpoints of bridge incidences leaving hyperedge-nodes in C .

Claim 5.2.1: running intersection.

Root each component of the forest T , and order the active components by nondecreasing depth, with arbitrary order among equal-depth components. When an active component C is added, its piece P_C meets the union of the earlier pieces in at most one point.

Proof of the claim. Every point of P_C is either a point-node belonging to C , or the point endpoint of a quotient-tree edge incident with C whose hyperedge endpoint lies in C .

Suppose $p \in P_C$ also lies in an earlier piece P_D . In the Levi graph, the occurrences of the same point-node p in the two pieces are linked through the component of L containing p . Consequently the unique path in T from C to D passes through the quotient component containing p , and the first edge on that path leaving C is incident with p .

Because D is earlier, it is not strictly inside a child subtree of C . Hence the path from C to D must leave C through the parent direction, except when C and D are siblings attached to the same inactive point component; in that case the same parent point is used. The quotient is a forest, so C has only one parent edge and two distinct parent incidences cannot join the same pair of quotient components. Thus every point shared with an earlier piece is the same point p_C determined by the parent direction. Therefore

$$P_C \cap \bigcup_{D \text{ earlier}} P_D \subseteq \{p_C\}.$$

For a root active component the intersection is empty. $\square$

Starting with a root piece, add the active pieces in this order. If the new intersection is empty, use a disjoint union. If it is the singleton $\{p_C\}$, use a one-point amalgamation at that point. Several earlier pieces may already contain the same inactive attachment point, but the new step still identifies only that one existing vertex. Different components of T are combined by disjoint union. Hence $F \in \mathcal{B}$. $\square$

This proves

$$(3) \implies (2). \tag{5.2}$$

Combining (5.1) and (5.2),

$$(2) \iff (3). \tag{5.3}$$

6. THE SEQUENCE LIFT USED IN THE AVOIDANCE CONSTRUCTIONS

Let $G = (X, A)$ be an arbitrary graph, where $A = E(G)$ is also regarded as an alphabet. Let

$$T = A^{<\omega_1}$$

be the tree of all sequences of graph edges of countable ordinal length. For $s \in T$ and $x \in X$, write $x_s = (s, x)$.

Definition 6.1: the one-apex lift.

Define the triple system $\mathcal{L}(G)$ by

$$V(\mathcal{L}(G)) = T \times X.$$

For $s \in T$, an edge $a = \{x, y\} \in A$, a sequence $t \in T$ extending $s \frown a$, and $z \in X$, put

$$\{x_s, y_s, z_t\} \in E(\mathcal{L}(G)). \quad (6.1)$$

The two vertices x_s, y_s form the base pair and z_t is the apex. The base node is uniquely recoverable from a hyperedge, so the system is simple and 3-uniform.

Lemma 6.2: chromatic lower bound.

If $\chi(G) > \aleph_0$, then

$$\chi(\mathcal{L}(G)) > \aleph_0.$$

Proof. Suppose $c : T \times X \rightarrow \omega$ is proper. For every $s \in T$, the restriction $x \mapsto c(x_s)$ is a countable colouring of G . Hence there is an edge

$$a_s = \{x_s^0, x_s^1\} \in E(G)$$

whose two vertices at node s have a common colour k_s .

Construct $s_\alpha \in A^\alpha$ for $\alpha < \omega_1$ by

$$s_0 = \emptyset, \quad s_{\alpha+1} = s_\alpha \widehat{a}_{s_\alpha},$$

and at limit ordinals take unions.

If $\alpha < \beta$, then s_β extends $s_\alpha \widehat{a}_{s_\alpha}$. If $k_{s_\alpha} = k_{s_\beta}$, take either endpoint z of the monochromatic edge at s_β . By (6.1),

$$\{(s_\alpha, x_{s_\alpha}^0), (s_\alpha, x_{s_\alpha}^1), (s_\beta, z)\}$$

is a monochromatic hyperedge. Therefore the colours k_{s_α} , $\alpha < \omega_1$, are pairwise distinct, impossible in ω . $\square$

Theorem 6.3: exact finite linear trace theorem.

Every finite linear subhypergraph K of $\mathcal{L}(G)$ is obtainable, by finite disjoint unions and one-point amalgamations, from systems J^+ where J is a finite subgraph of G .

Proof. Every hyperedge of $\mathcal{L}(G)$ has a unique base node s : exactly two of its vertices have sequence coordinate s , while the apex has a strictly longer sequence coordinate. Group the hyperedges of K by their base nodes, and let K_s consist of the hyperedges based at s .

For a fixed graph edge $a = \{x, y\} \in E(G)$, at most one hyperedge of K_s can have base pair $\{x_s, y_s\}$, since two such hyperedges would share two vertices and violate linearity. Hyperedges of K_s associated with distinct graph edges have distinct apices: if z_t were the apex for graph edges a and a' , the coordinate of t immediately after s would have to be both a and a' . It follows that K_s is precisely J_s^+ , where J_s is the finite subgraph of G consisting of the graph edges used at s .

Now consider intersections between different K_s . If a point z_t lies in both K_s and K_u , then both s and u are prefixes of t , so they are comparable. Suppose $s \subsetneq u$. A common point cannot be a core point of K_s ; it is an apex of K_s . The graph edge used by its hyperedge in K_s is the coordinate of u immediately following s . Hence all points in $K_s \cap K_u$ would be apices of the same base edge of K_s . There is at most one such apex. Therefore

$$|V(K_s) \cap V(K_u)| \leq 1. \quad (6.2)$$

Construct a bipartite incidence graph Q whose one class is the set of active base nodes s , and whose other class consists of points lying in at least two of the K_s ; join s to p when $p \in K_s$.

We claim that Q is a forest. Suppose it has a cycle, and choose on that cycle a base node s of minimum length. Its two neighbouring base nodes on the cycle are proper descendants of s . The two intervening shared points are distinct, so they are distinct apices in K_s , arising from two distinct graph edges $a \neq a'$. Thus the two neighbouring base nodes lie in different immediate branches below s , one beginning with a and the other with a' .

On the remainder of the cycle, consecutive base nodes share a point and are therefore comparable. Every base node there has length at least that of s . Comparable descendants of s lie in the same immediate branch below s , so the remainder of the cycle cannot move from the a -branch to the a' -branch without passing through a base node shorter than s . This contradicts the choice of s . Hence Q is acyclic.

Root each component of Q . Start with the factor at a root base node, and attach every child factor at the unique shared point represented by the intervening point-node of Q . By (6.2) and acyclicity, every step is a one-point amalgamation. Different components are combined by disjoint union. $\square$

Corollary 6.4: restrictions on finite linear traces.

Every finite linear $K \subseteq \mathcal{L}(G)$ satisfies:

1. every hyperedge-node of $I(K)$ is incident with a bridge;
2. every Berge cycle of K is contained in a single factor J_s^+ ;
3. a Berge cycle of length ℓ in K yields an ordinary cycle of length ℓ in G .

Proof. In every J_s^+ the incidence with the private vertex is a bridge. Bridgehood is preserved by one-point amalgamation. A simple Levi cycle cannot cross a one-point cut, so every Berge cycle lies in one factor. Berge cycles in J_s^+ are exactly ordinary graph cycles in J_s . $\square$

7. UNCOUNTABLY CHROMATIC GRAPHS OF LARGE ODD GIRTH

For a positive integer r and an infinite cardinal κ , define the directed shift graph $D_r(\kappa)$ as follows:

- its vertices are increasing r -tuples

$$(\alpha_0, \dots, \alpha_{r-1}), \quad \alpha_0 < \dots < \alpha_{r-1} < \kappa;$$

- it has an arc

$$(\alpha_0, \dots, \alpha_{r-1}) \longrightarrow (\alpha_1, \dots, \alpha_r)$$

whenever $\alpha_0 < \dots < \alpha_r$.

Let $S_r(\kappa)$ be its underlying graph. Write

$$\beth_0(\aleph_0) = \aleph_0, \quad \beth_{j+1}(\aleph_0) = 2^{\beth_j(\aleph_0)}.$$

Lemma 7.1: chromatic number of shift graphs.

If

$$\kappa > \beth_{r-1}(\aleph_0),$$

then

$$\chi(S_r(\kappa)) > \aleph_0.$$

Proof. The case $r = 1$ is immediate because $S_1(\kappa) = K_\kappa$. Assume $r \geq 2$. Suppose $S_r(\kappa)$ has a proper colouring c with λ colours. For every increasing $(r-1)$ -tuple u , define

$$C(u) = \{c(u \frown \beta) : \beta > \max u\} \subseteq \lambda.$$

If

$$u = (\alpha_0, \dots, \alpha_{r-2}), \quad v = (\alpha_1, \dots, \alpha_{r-1})$$

are adjacent in $S_{r-1}(\kappa)$, then

$$c(\alpha_0, \dots, \alpha_{r-1}) \in C(u).$$

It cannot belong to $C(v)$, since that would give some $\beta > \alpha_{r-1}$ for which the adjacent r -tuples

$$(\alpha_0, \dots, \alpha_{r-1}), \quad (\alpha_1, \dots, \alpha_{r-1}, \beta)$$

have the same colour. Hence $C(u) \neq C(v)$. Thus a λ -colouring of $S_r(\kappa)$ yields a 2^λ -colouring of $S_{r-1}(\kappa)$.

Starting with a countable colouring of $S_r(\kappa)$ and iterating $r - 1$ times would give a colouring of

$$S_1(\kappa) = K_\kappa$$

with at most $\beth_{r-1}(\aleph_0)$ colours, impossible when κ is larger. $\square$

Lemma 7.2: odd girth of shift graphs.

The odd girth of $S_r(\kappa)$ is at least

$$2r + 1.$$

Proof. For a digraph D , let δD be its arc digraph: the vertices of δD are the arcs of D , and

$$(u, v) \longrightarrow (v, w)$$

is an arc of δD . We have

$$D_{r+1}(\kappa) \cong \delta D_r(\kappa).$$

We use the following fact: if D is acyclic and the underlying graph of D has odd girth g , then the underlying graph of δD has odd girth at least $g + 2$.

Take an odd cycle $a_0, \dots, a_{m-1}$ in the underlying graph of δD . Record, around the cycle, whether each adjacency is oriented forward or backward in δD . Since δD is acyclic, the cyclic sign sequence is not constant. It therefore has at least two sign changes, and the number of sign changes is even.

For each adjacency a_{i-1}, a_i , take their common endpoint in D . Consecutive such endpoints are joined by the arc a_i when the two adjacent signs agree, and are equal when the signs change. Deleting the stationary steps gives an odd closed walk in the underlying graph of D , of length at most $m - 2$. Every odd closed walk contains an odd cycle, so $m \geq g + 2$.

Now $D_1(\kappa)$ is the transitive tournament; it is acyclic and its underlying graph K_κ has odd girth 3. Iterating the preceding observation gives odd girth at least

$$3 + 2(r - 1) = 2r + 1.$$

$\square$

Corollary 7.3: ZFC graphs with uncountable chromatic number and prescribed odd girth.

For every finite m there is a graph G with $\chi(G) > \aleph_0$ and no odd cycle of length at most m .

Proof. Choose r with $2r + 1 > m$, and let

$$G = S_r(\beth_{r-1}(\aleph_0)^+).$$

Apply Lemmas 7.1 and 7.2. $\square$

8. THE AVOIDANCE HOSTS

We prove

$$\neg(3) \implies \neg(1).$$

Proposition 8.1: avoidance of every nonlinear finite triple system.

There is an uncountably chromatic linear triple system. Consequently, every finite nonlinear triple system is non-obligatory.

Proof. We use the standard Erdős–Rado relation [3]

$$(2^{\aleph_0})^+ \longrightarrow (\aleph_1)_{\aleph_0}^2. \quad (8.1)$$

Let $\kappa = (2^{\aleph_0})^+$. Define a triple system T_κ whose vertices are the pairs in $[\kappa]^2$, and whose hyperedges are

$$\{\{\alpha, \beta\}, \{\alpha, \gamma\}, \{\beta, \gamma\}\}, \quad \alpha < \beta < \gamma < \kappa.$$

This triple system is linear: two distinct triangles of a complete graph cannot share two graph edges, because two graph edges of a triangle determine the third edge and the three underlying vertices.

It is uncountably chromatic. A countable vertex-colouring of T_κ is a countable edge-colouring of K_κ . By (8.1) there is a monochromatic set of type $\aleph_1$, and in particular a monochromatic graph triangle. This is a monochromatic hyperedge of T_κ .

If a finite triple system F is nonlinear, it has two distinct edges sharing at least two vertices. No injective embedding of those edges can exist in the linear host T_κ . Thus T_κ is F -free. $\square$

Proposition 8.2: avoidance when the bridge condition fails.

Let F be finite and linear, and suppose some hyperedge-node of $I(F^\circ)$ is incident with no bridge. Then F is non-obligatory.

Proof. By Lemma 1.1 it is enough to avoid F° , so assume that F has no isolated vertices. Take $G = K_{\omega_1}$. Then $\chi(G) > \aleph_0$, and Lemma 6.2 gives

$$\chi(\mathcal{L}(G)) > \aleph_0.$$

If F appeared in $\mathcal{L}(G)$, retain exactly the host hyperedges corresponding to the edges of F . Because the embedding is injective and F is linear, these selected host hyperedges form a finite linear subhypergraph isomorphic to F . Corollary 6.4 says that every hyperedge-node of such a trace is incident with a bridge, contradicting the chosen hyperedge of F . Therefore $\mathcal{L}(K_{\omega_1})$ is an uncountably chromatic F -free host. $\square$

Proposition 8.3: avoidance of an odd Berge cycle.

Let F be finite and linear, and suppose F° has an odd Berge cycle. Then F is non-obligatory.

Proof. Again suppress isolated vertices. Put $m = |E(F)|$. Choose r with $2r + 1 > m$, and set

$$G = S_r(\beth_{r-1}(\aleph_0)^+).$$

By Lemmas 7.1 and 7.2, $\chi(G) > \aleph_0$, and G has no odd cycle of length at most m . Lemma 6.2 gives

$$\chi(\mathcal{L}(G)) > \aleph_0.$$

If F appeared in $\mathcal{L}(G)$, the selected image hyperedges would form a finite linear trace. By Corollary 6.4, every Berge cycle of that trace lies in one factor J_s^+ and induces an ordinary graph cycle of the same length in $J_s \subseteq G$.

The odd Berge cycle of F has length at most m , so G would contain an odd cycle of length at most m , a contradiction. Thus $\mathcal{L}(G)$ is an uncountably chromatic F -free host. $\square$

The three propositions cover every failure of the intrinsic conditions, and prove

$$\neg(3) \implies \neg(1). \quad (8.2)$$

9. CONCLUSION

We have established:

- (2) $\implies$ (1) from the direct bipartite-expansion atom and the two closure arguments;
- (2) $\iff$ (3) from the Levi bridge-block decomposition and running-intersection reconstruction;
- $\neg(3) \implies \neg(1)$ from the explicit nonlinear host and the sequence-lift avoidance constructions.

Therefore

$$F \text{ is obligatory} \iff F \in \mathcal{B},$$

$$F^\circ \text{ is linear,}$$

$$F \in \mathcal{B} \iff \begin{array}{l} \text{every hyperedge-node of } I(F^\circ) \text{ is incident with a bridge,} \\ \text{every Berge cycle of } F^\circ \text{ has even length.} \end{array}$$

All embeddings are non-induced: additional host hyperedges on the image vertices are irrelevant throughout. The proof uses only ZFC, the displayed successor cardinals, graph-colouring compactness, and the standard Erdős–Rado relation (8.1). It assumes neither CH nor GCH, and no forcing axiom or large-cardinal hypothesis.

REFERENCES

- [1] E. Li. *A Resolution of Erdős Problems 593 and 1177: Obligatory Triple Systems and Exact Spectra*. Version 1. Preprint cited solely for bibliographic completeness. It claims resolutions of Erdős Problems 593 and 1177; no claim from it is used as a premise or independently verified in this manuscript. June 2026. DOI: 10.48550/arXiv.2606.24882. arXiv: 2606.24882 [math.CO]. URL: <https://arxiv.org/abs/2606.24882>.
- [2] N. G. de Bruijn and P. Erdős. “A Colour Problem for Infinite Graphs and a Problem in the Theory of Relations”. In: *Nederl. Akad. Wetensch. Proc. Ser. A = Indagationes Mathematicae* 54 = 13 (1951), pp. 369–373.
- [3] P. Erdős and R. Rado. “A Partition Calculus in Set Theory”. In: *Bulletin of the American Mathematical Society* 62 (1956), pp. 427–489. DOI: 10.1090/S0002-9904-1956-10036-0.
- [4] P. Erdős. “On Some Problems in Combinatorial Set Theory”. In: *Publikacije Instituta Matematičkog (Beograd) (N.S.)* 57(71) (1995), pp. 61–65.
- [5] T. F. Bloom. *Erdős Problem 593*. Erdős Problems. URL: <https://www.erdosproblems.com/593> (visited on 07/11/2026).
- [6] P. Komjáth. “Some Remarks on Obligatory Subsystems of Uncountably Chromatic Triple Systems”. In: *Combinatorica* 21.2 (2001), pp. 233–238. DOI: 10.1007/s004930100021.
- [7] A. Hajnal and P. Komjáth. “Obligatory Subsystems of Triple Systems”. In: *Acta Mathematica Hungarica* 119.1–2 (2008), pp. 1–13. DOI: 10.1007/s10474-007-6231-2.
- [8] P. Erdős, A. Hajnal, and B. L. Rothschild. “On Chromatic Number of Graphs and Set-Systems”. In: *Cambridge Summer School in Mathematical Logic (Cambridge, 1971)*. Vol. 337. Lecture Notes in Mathematics. Berlin and New York: Springer, 1973, pp. 531–538.
- [9] P. Erdős and A. Hajnal. “On Chromatic Number of Graphs and Set-Systems”. In: *Acta Mathematica Academiae Scientiarum Hungaricae* 17 (1966), pp. 61–99. DOI: 10.1007/BF02020444.
- [10] P. Erdős, F. Galvin, and A. Hajnal. “On Set-Systems Having Large Chromatic Number and Not Containing Prescribed Subsystems”. In: *Infinite and Finite Sets (Keszthely, 1973)*. Vol. 10. Colloquia Mathematica Societatis János Bolyai. North-Holland, 1975, pp. 425–513.
- [11] P. Komjáth. “An Uncountably Chromatic Triple System”. In: *Acta Mathematica Hungarica* 121.1–2 (2008), pp. 79–92. DOI: 10.1007/s10474-008-7179-6.
- [12] C. Reiher. “Obligatory Hypergraphs”. In: *Proceedings of the American Mathematical Society* (2024). DOI: 10.1090/proc/17021. arXiv: 2403.11223. URL: <https://arxiv.org/abs/2403.11223>.
- [13] C. Reiher. *Graphs of Large Girth*. 2024. arXiv: 2403.13571. URL: <https://arxiv.org/abs/2403.13571>.
- [14] Y. Wang, M. Duan, D. Gerbner, and H. Hama Karim. *On the Largest Chromatic Number of F -Free Hypergraphs*. 2026. arXiv: 2604.21551. URL: <https://arxiv.org/abs/2604.21551>.

ETHICS AND AI-ASSISTANCE STATEMENT

This manuscript was completed with assistance from OpenAI’s ChatGPT-5.6 Pro for proof development, adversarial checking, drafting, editing, bibliographic organisation, and typesetting. Samuil Pekov is the sole author and accepts responsibility for the verification of all mathematical arguments, citations, and conclusions. The AI system is not an author.