

Sam Chong Tay  
Math 341: Real Analysis  
Take-home Final Exam  
December 19, 2012

1. (15 pts) Let  $A$  and  $B$  be non-empty subsets of  $\mathbb{R}$  with  $A \subseteq B$ . Suppose that  $\inf A$ ,  $\inf B$ , and  $\sup B$  all exist.

- (a) Prove that  $\sup A$  exists.

*Proof.* We know  $A$  is non-empty by hypothesis, so we only need to show that  $A$  is bounded above. Considering any  $a \in A$ , we see that  $a \in B$  as well because  $A \subseteq B$ . Thus  $a \leq \sup B$ , and since  $a$  was arbitrary we have shown that  $A$  is bounded above by  $\sup B$ . Hence by Axiom III,  $\sup A$  exists.  $\square$

- (b) Prove that

$$\inf(B) \leq \inf(A) \leq \sup(A) \leq \sup(B).$$

*Proof.* For the first inequality, it suffices to show that  $\inf(B)$  is a lower bound for  $A$ . So, similar to part (a), we note that for any  $a \in A$ ,  $a$  is also in  $B$  and therefore  $\inf(B) \leq a$ . Hence  $\inf(B)$  is a lower bound for  $A$  and since  $\inf(A)$  is the greatest lower bound, we have

$$\inf(B) \leq \inf(A).$$

The next inequality is quite trivial; since  $A$  is non-empty there exists  $a \in A$  and by definition of infimum and supremum we have immediately that

$$\inf(A) \leq a \leq \sup(A).$$

Finally, note that we showed in part (a) that  $\sup(B)$  is an upper bound for  $A$ . Therefore just as in the infimum case, since  $\sup(A)$  is the least upper bound, we have  $\sup(A) \leq \sup(B)$ . Altogether we have shown that

$$\inf(B) \leq \inf(A) \leq \sup(A) \leq \sup(B).$$

$\square$

- (c) Show by giving a counterexample that

$$\inf(B) = \inf(A) \quad \text{and} \quad \sup(A) = \sup(B)$$

need not imply that  $A = B$ .

*Counterexample:* If we define

$$A = \{0, 1\} \quad \text{and} \quad B = [0, 1]$$

we see that

$$\inf(A) = \inf(B) = 0 \quad \text{and} \quad \sup(A) = \sup(B) = 1,$$

but of course  $A \neq B$ .

2. (15 pts) Prove that every non-convergent bounded sequence of real numbers has at least two limit points.

*Proof.* Let  $(s_n)$  be a non-convergent bounded sequence in  $\mathbb{R}$ . Because  $(s_n)$  is bounded we know from Theorem 3.4.9 that  $(s_n)$  has at least one convergent subsequence, and hence at least one limit point  $x$ . Consider an arbitrary subsequence  $(s_{n_k})$  of  $(s_n)$ . This subsequence must also be bounded and thus by the same theorem has a convergent subsequence  $(s_{n_{k_i}})$ . Well  $(s_{n_{k_i}})$  is evidently also a subsequence of  $(s_n)$  in its own right, so we see that if  $x$  was the only limit point, this would imply  $s_{n_{k_i}} \rightarrow x$ . But we chose an arbitrary subsequence, so we have shown that if  $x$  is the only limit point of  $(s_n)$  then every subsequence of  $(s_n)$  has a subsequence converging to  $x$ . By Problem 3.3.6 this would imply  $s_n \rightarrow x$ , which is a contradiction because we are assuming  $(s_n)$  is non-convergent. We have shown that  $(s_n)$  must have a limit point  $x$  and that  $x$  cannot be the only limit point, and therefore the only possibility is that  $(s_n)$  has at least two limit points.  $\square$

3. (20 pts)

- (a) Define  $h : \mathbb{R} \rightarrow \mathbb{R}$  by  $h(x) = x^3 - 2x^2 + 3x - 6$ . Carefully prove that the limit  $\lim_{x \rightarrow 2} h(x)$  exists. (Your solution should also indicate the limit!)

*Proof.* Carefully, and cleverly, we will first prove results about the identity function and constant function. Define the function  $i : \mathbb{R} \rightarrow \mathbb{R}$  by  $i(x) = x$ . First note that we have previously shown every  $x \in \mathbb{R}$  is a limit point of  $\mathbb{R}$ , so we know  $2 \in \mathbb{R}$  is a limit point of the domain  $\mathbb{R}$ . To see that  $\lim_{x \rightarrow 2} i(x) = 2$ , let  $\epsilon > 0$ . Then for  $\delta = \epsilon$ , if  $0 < |x - 2| < \delta$ ,

$$|i(x) - 2| = |x - 2| < \delta = \epsilon.$$

Therefore

$$\lim_{x \rightarrow 2} i(x) = 2.$$

Next fix  $k \in \mathbb{R}$  and define  $c_k : \mathbb{R} \rightarrow \mathbb{R}$  by  $c_k(x) = k$ . Let  $\epsilon > 0$  and pick  $\delta = 64$ . Then for  $0 < |x - 2| < \delta$ , we have

$$|c_k(x) - k| = |k - k| = 0 < \epsilon.$$

Therefore

$$\lim_{x \rightarrow 2} c_k(x) = k.$$

Next recall the function  $h$  from above and observe that

$$\begin{aligned} h(x) &= x^3 - 2x^2 + 3x - 6 \\ &= (i(x))^3 + c_{-2}(x)(i(x))^2 + c_3(x)i(x) + c_{-6}(x). \end{aligned}$$

Apply Theorem 5.3.1 parts (1) and (3) a total of eight times (if you want me to be *really* careful, we apply part (1) three times and part (3) five times) to obtain

$$\begin{aligned} \lim_{x \rightarrow 2} h(x) &= \lim_{x \rightarrow 2} \left( (i(x))^3 + c_{-2}(x)(i(x))^2 + c_3(x)i(x) + c_{-6}(x) \right) \\ &= (2)^3 + (-2)(2)^2 + (3)(2) + (-6) \\ &= 8 - 8 + 6 - 6 \\ &= 0. \end{aligned}$$

Of course applying Theorem 5.3.1 guarantees that the limit does exist, and that it is equal to 0.  $\square$

- (b) Carefully prove that the limit  $\lim_{x \rightarrow 0} \cos \frac{1}{x}$  fails to exist.

*Proof.* We need to be quite careful indeed; the problem is actually stated quite ambiguously because it is not clear over which domain we are to consider this function. I think it is safe to assume that we are considering a function  $f : \mathbb{R}^* \rightarrow \mathbb{R}$  defined by  $f(x) = \cos \frac{1}{x}$ , where  $0 \in \mathbb{R}$  is a limit point of the subset  $\mathbb{R}^*$ . To show  $\lim_{x \rightarrow 0} f(x)$  does not exist, we will show that it is not equal to any candidate  $L \in \mathbb{R}$ . First consider  $L = 0$  and pick  $\epsilon = \frac{1}{2}$ . Let  $\delta > 0$  and choose  $n \in \mathbb{N}$  such that  $n > \frac{1}{\delta}$ . Then

$$0 < \left| \frac{1}{2n\pi} - 0 \right| < \frac{1}{n} < \delta,$$

yet

$$\left| f\left(\frac{1}{2n\pi}\right) - L \right| = \left| \cos(2n\pi) - 0 \right| = |1 - 0| = 1 > \epsilon.$$

This shows that  $\lim_{x \rightarrow 0} f(x) \neq 0$ . Now consider any  $L \neq 0$ , and pick  $\epsilon = |L|$ . As before, for any  $\delta > 0$  choose  $n \in \mathbb{N}$  such that  $n > \frac{1}{\delta}$ . Then

$$0 < \left| \frac{1}{2n\pi + \frac{\pi}{2}} - 0 \right| < \frac{1}{n} < \delta,$$

yet

$$\left| f\left(\frac{1}{2n\pi + \frac{\pi}{2}}\right) - L \right| = \left| \cos\left(2n\pi + \frac{\pi}{2}\right) - L \right| = |0 - L| = |L| = \epsilon.$$

Therefore  $\lim_{x \rightarrow 0} f(x) \neq L$ . So in considering the cases when  $L = 0$  and  $L \neq 0$ , we have shown that the limit of  $f$  as  $x$  goes to 0 is not  $L$  for any  $L$  in the codomain  $\mathbb{R}$ . Therefore the limit does not exist.  $\square$

4. (15 pts) Let  $(X, d)$  be a metric space. Show that the following statements about  $X$  are equivalent.

- i.  $(X, d)$  is discrete.
- ii. All functions on  $X$  are continuous. (The range can be *any* metric space!)

*Proof.* ( $\implies$ ) Suppose  $(X, d)$  is discrete. Consider any  $a \in X$ . Since  $\{a\}$  is open, there exists  $r > 0$  such that  $B_r(a) \subseteq \{a\}$ , from which it follows that  $B_r(a) = \{a\}$ . Then obviously  $B_r(a)$  does not contain infinitely many points of  $X$ , so by the characterization in Theorem 3.5.1.3 we conclude that  $a$  is not a limit point of  $X$ . Thus for any metric space  $Y$  and any function  $f : X \rightarrow Y$ ,  $f$  is continuous at  $a$  by definition. Since  $a$  was arbitrary we conclude that all functions on  $X$  are continuous.

( $\impliedby$ ) Conversely suppose that all functions on  $X$  are continuous. Let  $a \in X$  and define the function  $f_a : X \rightarrow \mathbb{R}$  by

$$f_a(x) = \begin{cases} 64 & \text{if } x = a \\ 0 & \text{if } x \neq a \end{cases}$$

Note that the open interval  $(63, 65) = B_1(64)$  is an open subset of  $\mathbb{R}$ . Since  $f_a$  is continuous by hypothesis, Theorem 4.3.5 guarantees that the inverse image of  $(63, 65)$  is open in  $X$ . By the construction above we have

$$f_a^{-1}((63, 65)) = \{a\},$$

so the previous sentence implies that  $\{a\}$  is open in  $X$ . Since  $a$  was arbitrary, we conclude  $X$  is a discrete metric space.  $\square$

5. (15 pts) Let  $(a_n)$  and  $(b_n)$  be Cauchy sequences in a metric space  $(X, d)$ . Show that the sequence  $(d(a_n, b_n))$  converges. (Note that  $X$  need not be complete, but  $\mathbb{R}$  is complete!)

*Proof.* As hinted above, we will show that  $(d(a_n, b_n))$  is a Cauchy sequence and invoke the completeness of  $\mathbb{R}$  to complete the proof. Let  $\epsilon > 0$ . Since  $(a_n)$  and  $(b_n)$  are Cauchy sequences, we can choose  $N_1 \in \mathbb{N}$  and  $N_2 \in \mathbb{N}$  so that

$$d(a_n, a_m) < \frac{\epsilon}{2} \quad \text{for all } n, m > N_1$$

$$d(b_n, b_m) < \frac{\epsilon}{2} \quad \text{for all } n, m > N_2.$$

Let  $N = \max\{N_1, N_2\}$  and  $n, m > N$ . Supposing without loss of generality that  $d(a_n, b_n) \geq d(a_m, b_m)$ ,

$$\begin{aligned}
 |d(a_n, b_n) - d(a_m, b_m)| &= d(a_n, b_n) - d(a_m, b_m) \\
 &\leq d(a_n, a_m) + d(a_m, b_n) - d(a_m, b_m) \\
 &\leq d(a_n, a_m) + d(b_n, b_m) + d(b_m, a_m) - d(a_m, b_m) \\
 &= d(a_n, a_m) + d(b_n, b_m) \\
 &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.
 \end{aligned}$$

Therefore  $(d(a_n, b_n))$  is a Cauchy sequence in  $\mathbb{R}$  and since  $\mathbb{R}$  is complete,  $(d(a_n, b_n))$  converges.  $\square$

6. (20 pts)

- (a) Let  $(f_n)$  be a sequence of real-valued continuous functions on a metric space  $X$ . Suppose that  $(f_n)$  converges uniformly to a function  $f$ . Prove that if all terms (i.e. all functions) of the sequence  $(f_n)$  are bounded, then  $f$  is bounded.

*Proof.* Let  $X$  be a metric space and  $f_n : X \rightarrow \mathbb{R}$  be a sequence of bounded functions that converges uniformly to  $f$ . To show that  $f$  is bounded we need to show that the range  $f(X) = \{f(x) : x \in X\}$  is bounded in  $\mathbb{R}$ , and to do so we will use the characterization given in Corollary 3.1.14. Explicitly, we will show that there exists  $M \in \mathbb{R}$  such that  $|f(x)| \leq M$  for all elements  $f(x) \in f(X)$ . We will need to use the uniform convergence assumption, so let  $\epsilon = 64$ . Since  $(f_n)$  converges uniformly to  $f$ , we can choose  $N \in \mathbb{N}$  so that for all  $n > N$  and all  $x \in X$ ,  $|f_n(x) - f(x)| < 64$ . Now fix an  $n > N$ . Since  $f_n$  is bounded, by Corollary 3.1.14 there exists  $K \in \mathbb{R}$  such that  $|f_n(x)| \leq K$  for all  $x \in X$ . Then for any  $x \in X$ ,

$$\begin{aligned}
 |f(x)| &= |f(x) - 0| \leq |f(x) - f_n(x)| + |f_n(x) - 0| \\
 &< 64 + |f_n(x)| \\
 &\leq 64 + K.
 \end{aligned}$$

Therefore  $|f(x)| \leq 64 + K$  for all  $x \in X$ , so  $f(X)$  is bounded in  $\mathbb{R}$ . By definition,  $f$  is bounded.  $\square$

- (b) Recall that a function  $f : X \rightarrow Y$  is said to be **bounded** if its range  $f(X)$  is a bounded subset of  $Y$ . Give an example to show that the pointwise limit of real-valued bounded functions on a metric space  $X$  need not be a bounded function. Hint: consider piecewise defined functions on  $\mathbb{R}$ .

*Example:* Consider  $\mathbb{N}$  under the Euclidean metric and the sequence of functions  $f_n : \mathbb{N} \rightarrow \mathbb{R}$  defined by

$$f_n(m) = \begin{cases} m & \text{if } m \leq n \\ 0 & \text{if } m > n. \end{cases}$$

Now each  $f_n$  is bounded by  $n$ :

$$\begin{aligned} f_n(\mathbb{N}) &= \{1, 2, 3, \dots, n, 0, 0, 0, \dots\} \\ &= \{0, 1, 2, \dots, n\}. \end{aligned}$$

However, it is clear that  $(f_n)$  converges pointwise to the function  $f(m) = m$ . To show this rigorously, let  $m \in \mathbb{N}$  (as a point in the domain!) and  $\epsilon > 0$ . Then for all  $n > m$  (as a sequence index!),

$$|f_n(m) - f(m)| = |m - m| = 0 < \epsilon.$$

Therefore  $f_n(m) \rightarrow f(m)$  at the point  $m$ , and since  $m$  was arbitrary we have shown that  $(f_n)$  converges pointwise to  $f$ . But obviously  $f(\mathbb{N}) = \mathbb{N}$  is unbounded! Therefore uniform convergence is in fact a necessary condition for part (a).

Of course, using the same idea I could have constructed  $f_n : \mathbb{R} \rightarrow \mathbb{R}$  as

$$f_n(x) = \begin{cases} |x| & \text{if } x \leq n \\ 0 & \text{if } x > n \end{cases}$$

and gotten the same result. This would have even avoided confusion with regards to choosing  $m \in \mathbb{N}$  in the domain and shortly thereafter using  $N = m$  for the sequence index that we need in showing a sequence converges. However, I like the way I did it because  $\mathbb{N}$  is 1-separated under the Euclidean metric and thus has no limit points. So the sequence functions  $f_n$ , while piecewise, are actually continuous.