

on each wing while the hindwings are ochre background with black border and few black spots, two dark brown wavy broad cross strips. Wingspan is about 10 cm. The thorax bears a prominent Death's head mark. Abdomen is ochraceous in general hue with dark-brown cross bands.

4. Leaf hopper, *Orosius albicinctus*, Cicadellidae, Hemiptera

Symptom of damage: Phyllody.

Nature of damage: Vector of sesamum phyllody.

Adult: Light brown coloured hoppers.

5. Aphid, *Aphis gossypii*, Aphididae, Hemiptera

6. Stink bugs, *Eusarcocoris ventralis*, Pentatomidae, Hemiptera

7. Stink bugs, *Nezara viridula*, Pentatomidae, Hemiptera

8. Stink bugs, *Dolycoris indicus*, Pentatomidae, Hemiptera

9. Lygaeid bug, *Aphanus sordidus*, Lygaeidae, Hemiptera

10. Lygaeid bug, *Nysius inconspicuus*, Lygaeidae, Hemiptera

SUNFLOWER

1. Capitulum borer, *Helicoverpa armigera*, Noctuidae, Lepidoptera

Nature of damage: Feeds on capitulum and leaves.

2. Tobacco caterpillar, *Spodoptera litura*, Noctuidae, Lepidoptera

3. Black hairy caterpillar, *Estigmene lactinea*, Arctiidae, Lepidoptera

4. Bihar hairy caterpillar, *Spilosoma oblique*, Arctiidae, Lepidoptera

5. Ash weevil, *Myllocerus* spp., Curculionidae, Coleoptera

6. Stink bug, *Nezara viridula*, Pentatomidae, Hemiptera

7. Stink bug, *Dolycoris indicus*, Pentatomidae, Hemiptera

8. Peach leaf curl aphid, *Brachycaudus helichrysi*, Aphididae, Hemiptera

9. Compositae tingid, *Cadmilos retiarus*, Tingidae, Hemiptera