

Deep Learning for Source Classification

Edward J. Kim^{1*}, Robert J. Brunner^{2,3,4}

¹*Department of Physics, University of Illinois, Urbana, IL 61801 USA*

²*Department of Astronomy, University of Illinois, Urbana, IL 61801 USA*

³*Department of Statistics, University of Illinois, Champaign, IL 61820 USA*

⁴*National Center for Supercomputing Applications, Urbana, IL 61801 USA*

26 October 2015

ABSTRACT

Deep learning.

Key words: methods: data analysis – methods: statistical – surveys – stars: statistics – galaxies: statistics.

1 INTRODUCTION

2 METHODS

3 DATA

4 RESULTS AND DISCUSSION

5 CONCLUSIONS

ACKNOWLEDGEMENTS

We thank XXXX.

Funding for SDSS-III has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, and the U.S. Department of Energy Office of Science. The SDSS-III web site is <http://www.sdss3.org/>.

SDSS-III is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS-III Collaboration including the University of Arizona, the Brazilian Participation Group, Brookhaven National Laboratory, Carnegie Mellon University, University of Florida, the French Participation Group, the German Participation Group, Harvard University, the Instituto de Astrofísica de Canarias, the Michigan State/Notre Dame/JINA Participation Group, Johns Hopkins University, Lawrence Berkeley National Laboratory, Max Planck Institute for Astrophysics, Max Planck Institute for Extraterrestrial Physics, New Mexico State University, New York University, Ohio State University, Pennsylvania State University, University of Portsmouth, Princeton University, the Spanish Participation Group, University of Tokyo, University of Utah, Vanderbilt University, University of Virginia, University of Washington, and Yale University.

This paper uses data from the VIMOS Public Extragalactic Redshift Survey (VIPERS). VIPERS has been performed using the ESO Very Large Telescope, under the "Large Programme" 182.A-0886. The participating institutions and funding agencies are listed at <http://vipers.inaf.it/>.

This research uses data from the VIMOS VLT Deep Survey, obtained from the VVDS database operated by Cesam, Laboratoire d'Astrophysique de Marseille, France.

This paper has been typeset from a \TeX / \LaTeX file prepared by the author.

* jkim575@illinois.edu