

## Genome sequence of *Atlantibacter subterranea* DSM 16208 Isolated from Uranium-contaminated Sediment

Sofia Amaral<sup>a</sup>, Luis Garcia<sup>a</sup>, Adrianna Gonzalez<sup>a</sup>, Samantha Prince<sup>a</sup>, Komal Takhar<sup>a</sup>, Tricia Van Laar<sup>a#</sup>

a. Department of Biology, California State University, Stanislaus, California, USA

Genome Sequence of *Atlantibacter subterranea*

#Address correspondence to Tricia Van Laar, [tvanlaar@csustan.edu](mailto:tvanlaar@csustan.edu)

All authors contributed equally to this work. Author order was determined by alphabetical order.

### Abstract

We report the whole genome sequence of an antibiotic-resistant strain of *Atlantibacter subterranea* isolated from Uranium-contaminated sediment in Tennessee. The genome sequence of this strain was 4,717,064 bp in length, contained 34 contigs and 29 scaffolds, and with a GC content of 55.17%.

### Announcement

*Atlantibacter subterranea* is a rod-shaped, gram-negative bacterium, found in aquatic areas in North America (1). It is facultatively anaerobic, motile, and may spread in animals by consuming contaminated food (2). The bacterium can reduce hexavalent uranium to tetravalent uranium, later


precipitating to mineral uraninite, effectively immobilizing uranium from radioactive waste (3). This strain was isolated from Uranium (VI)-contaminated subsurface sediment in 2001 in Tennessee, USA using sediment sampling, enrichment, and serial dilutions plated on aerobic agar containing acetate (3). Analysis of *A. subterranea* aids in identifying similarities in specific extracellular polymeric substance secretions (EPS) across bacterial species related to its significance in uranium reduction (3). It was taxonomically identified before genome sequencing and the 16S rRNA gene sequence is found in the NCBI database (accession number AY373829) (3).

**Details on organism growth and DNA isolation to be provided by DSMZ.** Sequencing was performed at JGI using Illumina HiSeq 2000 platform by creating an Illumina std shotgun library, TSPS, with a read type of 2x150 bp. It resulted in 2,999,762 raw sequence reads and 450 mb DNA sequences. Reads were filtered using DUK to remove known contaminants including Illumina sequencing and library preparation artifacts (4). Genome assembly initially used Velvet (v1.02.07) (parameters: contig length 500, coverage cutoff 10) (5) followed by Allpaths-LG (v46652) (parameters: PHRED 64, PLOIDY=1, COVERAGE=125) (6). The genome was annotated using the JGI Microbial Genome Annotation Pipeline (v5.0.0) (7). CheckM2 (v5.5.2) determined 100% completion and 0.11% contamination (8).

<b>Table 1 - Genomic features of <i>Atlantibacter subterranea</i> DSM 16208</b>	
<b><u>Feature</u></b>	<b><u>Finding</u></b>
<u>length (bp)</u>	<u>4,717,064</u>
<u>status</u>	<u>complete</u>
<u>No. of contigs</u>	<u>34</u>

<b>Table 1 - Genomic features of <i>Atlantibacter subterranea</i> DSM 16208</b>	
<b>Feature</b>	<b>Finding</b>
<u>length (bp)</u>	<u>4,717,064</u>
<u>GC content (%)</u>	<u>55.17%</u>
<u>No. of scaffolds</u>	<u>29</u>
<u>Scaffold N50 (bp)</u>	<u>434618</u>
<u>Average fold coverage</u>	<u>0.635938</u>
<u>No. of rRNAs</u>	<u>17</u>
<u>No. of tRNAs</u>	<u>67</u>
<u>No. of genes</u>	<u>4504</u>
<u>No. of coding sequences</u>	<u>4351</u>

38



39

40 **FIG 1** Whole-genome-based phylogenetic classification of *Atlantibacter subterranea*. The  
41 genome BLAST distance phylogeny (GBDP) tree was generated with the Type Strain Genome  
42 Server accessed 14 March 2025. The tree was inferred with FastME 2.1.6.1 (9) from GBDP  
43 distances calculated from genome sequences. The branch lengths are scaled in terms of the  
44 GBDP distance formula  $d_5$ . The numbers at the nodes are GBDP pseudobootstrap support values  
45 of >60% from 100 replications. The average branch support was 93.7%. The tree was midpoint  
46 rooted (10).

The genome of *A. subterranea* consists of one chromosome measuring 4,717,064 bp with a G+C content of 55.17% (Table 1). Whole-genome-based phylogenetic classification, generated using the Type Strain Genome Server and inferred with FastME 2.1.6.1, shows that *A. subterranea* is most closely related to *A. hermannii* NBRC 105704T, with close proximity to *Franconibacter helvetivus* LMG 23732 (Figure 1). The probability of *A. subterranea* being a human pathogen is 0.702, as determined by PathogenFinder v1.1 (11). Using the Comprehensive Antibiotic Resistance Database (CARD 4.0.0), we confirmed that *A. subterranea* is resistant to various types of antibiotics, including cephalosporin, betalactam, fluoroquinolone, and macrolide (12). CRISPR-Cas Finder version 1.1.2 -I2BC identified 1 CRISPR region (13). AntiSMASH 7.0 software identified the production of secondary metabolites including arylpolyene, NRP metallophore, terpene, thiopeptide, butyrolactone and RiPP like (14).

## Data availability statement

This Whole Genome Shotgun project has been deposited in DDBJ/ENA/GenBank under the accession no. [LC126283](#). The version described in this paper is the first version, [LC126283.1](#). The data was deposited under the BioProject accession no. [PRJDB2388](#), the BioSample accession no. [SAMMD00010876](#), and the Sequence Read Archive accession no. [DRR015979](#).

## Acknowledgments

The work (proposal DOI[:xxx]) conducted by the U.S. Department of Energy Joint Genome Institute (<https://ror.org/04xm1d337> [Links to an external site.](#)), a DOE Office of Science User Facility, is supported by the Office of Science of the U.S. Department of Energy operated under contract no. [DE-AC02-05CH11231](#) [Links to an external site.](#)

## 68    **References**

- 69    1.    Reimer LC, Sarda Carbasse J, Schober I, Koblit J, Podstawka A, Overmann J. 2024. *Salmonella*  
70        *subterranea* Shelobolina et al. 2005 (9.2). DSMZ.
  
- 71    2.    Hata H, Natori T, Mizuno T, Kanazawa I, Eldesouky I, Hayashi M, Miyata M, Fukunaga H, Ohji S,  
72        Hosoyama A, Aono E, Yamazoe A, Tsuchikane K, Fujita N, Ezaki T. 2016. Phylogenetics of family  
73        Enterobacteriaceae and proposal to reclassify *Escherichia hermannii* and *Salmonella subterranea* as  
74        *Atlantibacter hermannii* and *Atlantibacter subterranea* gen. nov., comb. nov. *Microbiol Immunol*  
75        60:303–311.
  
- 76    3.    Shelobolina ES, Sullivan SA, O'Neill KR, Nevin KP, Lovley DR. 2004. Isolation, characterization,  
77        and U(VI)-reducing potential of a facultatively anaerobic, acid-resistant Bacterium from Low-pH,  
78        nitrate- and U(VI)-contaminated subsurface sediment and description of *Salmonella subterranea* sp.  
79        nov. *Appl Environ Microbiol* 70:2959–2965.
  
- 80    4.    Sharp CE, Smirnova AV, Kalyuzhnaya MG, Bringel F, Hirayama H, Jetten MSM, Khmelenina VN,  
81        Klotz MG, Knief C, Kyrpides N, Camp HJMO den, Reshetnikov AS, Sakai Y, Shapiro N,  
82        Trotsenko YA, Vuilleumier S, Woyke T, Dunfield PF. 2015. Draft Genome Sequence of the  
83        Moderately Halophilic Methanotroph *Methylohalobius crimeensis* Strain 10Ki. *Genome*  
84        Announcements 3:e00644.
  
- 85    5.    Zerbino DR, Birney E. 2008. Velvet: Algorithms for de novo short read assembly using de Bruijn  
86        graphs. *Genome Res* 18:821–829.
  
- 87    6.    Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea  
88        TP, Sykes S, Berlin AM, Aird D, Costello M, Daza R, Williams L, Nicol R, Gnirke A, Nusbaum C,  
89        Lander ES, Jaffe DB. 2011. High-quality draft assemblies of mammalian genomes from massively  
90        parallel sequence data. *Proceedings of the National Academy of Sciences* 108:1513–1518.

91 7. Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Palaniappan K, Szeto E,  
92 Pillay M, Chen I-MA, Pati A, Nielsen T, Markowitz VM, Kyrpides NC. 2015. The standard  
93 operating procedure of the DOE-JGI Microbial Genome Annotation Pipeline (MGAP v.4).  
94 Standards in Genomic Sciences 10:86.

95 8. Chklovski A, Parks DH, Woodcroft BJ, Tyson GW. 2023. CheckM2: a rapid, scalable and accurate  
96 tool for assessing microbial genome quality using machine learning. Nat Methods 20:1203–1212.

97 9. Lefort V, Desper R, Gascuel O. 2015. FastME 2.0: A Comprehensive, Accurate, and Fast Distance-  
98 Based Phylogeny Inference Program. Molecular Biology and Evolution 32:2798.

99 10. Farris JS. 1972. Estimating Phylogenetic Trees from Distance Matrices. The American Naturalist  
100 106:645–668.

101 11. Cosentino S, Larsen MV, Aarestrup FM, Lund O. 2013. PathogenFinder - Distinguishing Friend  
102 from Foe Using Bacterial Whole Genome Sequence Data. PLOS ONE 8:e77302.

103 12. Alcock BP, Huynh W, Chalil R, Smith KW, Raphenya AR, Wlodarski MA, Edalatmand A, Petkau  
104 A, Syed SA, Tsang KK, Baker SJC, Dave M, McCarthy MC, Mukiri KM, Nasir JA, Golbon B,  
105 Imtiaz H, Jiang X, Kaur K, Kwong M, Liang ZC, Niu KC, Shan P, Yang JYJ, Gray KL, Hoad GR,  
106 Jia B, Bhando T, Carfrae LA, Farha MA, French S, Gordzevich R, Rachwalski K, Tu MM,  
107 Bordeleau E, Dooley D, Griffiths E, Zubyk HL, Brown ED, Maguire F, Beiko RG, Hsiao WWL,  
108 Brinkman FSL, Van Domselaar G, McArthur AG. 2023. CARD 2023: expanded curation, support  
109 for machine learning, and resistome prediction at the Comprehensive Antibiotic Resistance  
110 Database. Nucleic Acids Res 51:D690–D699.

111 13. Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Néron B, Rocha EPC,  
112 Vergnaud G, Gautheret D, Pourcel C. CRISPRCasFinder, an update of CRISRFinder, includes a  
113 portable version, enhanced performance and integrates search for Cas proteins.

114 14. Blin K, Shaw S, Augustijn HE, Reitz ZL, Biermann F, Alanjary M, Fetter A, Terlouw BR, Metcalf  
115 WW, Helfrich EJN, van Wezel GP, Medema MH, Weber T. 2023. antiSMASH 7.0: new and  
116 improved predictions for detection, regulation, chemical structures and visualisation. *Nucleic Acids*  
117 *Research* 51:W46–W50.

118