

Recursive structure of primes reloaded. Now with mod 2, too.*

Carolin Zöbelein[†]

March 21, 2016

Abstract.

Contents

1	Introduction	1
2	Road map	1
3	What we already know ...	2
4	Intersection	3

1 Introduction

In this working document I will make a revision of my work [1]. Now for mod 2 and not mod 6 which makes all so easy. Sometimes I'm so stupid. wtf!

2 Road map

Since the main steps will be the same like in [1] I will only make a few comments. If you need more information please read the arXiv paper at first.

To future comments: Yes, I know it's no 'professional' math, but why I should use abstract math if it's also possible to explain issues with easy tools. This here is to be only a

***Keywords:** integer divisible numbers, prime numbers, recursive structure, mod 2; **License:** CC BY-ND 3.0 DE (see also LICENSE), No warranty for mistakes!

[†]*E-mail address:* contact@carolin-zoebelein.de, *PGP Fingerprint:* D4A7 35E8 D47F 801F 2CF6 2BA7 927A FD3C DE47 E13B; *URL:* <http://www.carolin-zoebelein.de>

first impression. Not the solution of all! After this you can construct your buildings, too!

Personal road map for this document:

1. Formulation of the important points for mod 2.
2. Consequences from this.
3. ...

3 What we already know ...

Short repetition of the properties of primes which we already know and we will use for our work.

1. Q: What are prime numbers \mathbb{P} ?
A: A prime number $p \in \mathbb{P}$ is a integer number larger than one which has no positive integer divisors apart from 1 and itself.
 \Rightarrow More mathematical: $p \in \mathbb{Z} : \gcd(n, p) = 1, \forall n \neq p \in \mathbb{N}$
2. The set of primes: $\mathbb{P} := \{2, 3, 5, 7, 11, 13, \dots\}$
3. The number 2 is the only even prime.

In the following work we will always ignore the prime 2!

\Rightarrow Hence, for all primes $p \in \mathbb{P} \setminus \{2\}$ we know that we are able to write them as $p_n := 2n + 1, n \in \mathbb{N}_0$. Q: For which n we receive primes?

Now if we look at the set of all odd integer divisible numbers $p_{nn'}$ $\in \mathbb{N}$. We recognise

$$\begin{aligned} p_{nn'} &:= (2n + 1)(2n' + 1) \\ &= 4nn' + 2n + 2n' + 1 \\ &= 2 \left(\underbrace{2nn' + n + n'}_{=: N_{n,n'}} \right) + 1, \end{aligned} \tag{1}$$

$\forall n, n' \in \mathbb{N}_0$. Since the case $a \cdot 1 = 1 \cdot a = a$ isn't interesting for use we will change our domain of definition from $n, n' \in \mathbb{N}_0$ to $n, n' \in \mathbb{N}$.

Additionally we will see that it make sense to expand p_n to p_z with the set \mathbb{Z} . In this case we will write

$$\begin{aligned} p_{z,z'} &:= (2z + 1)(2z' + 1) \\ &= 4zz' + 2z + 2z' + 1 \\ &= 2 \left(\underbrace{2zz' + z + z'}_{=: Z_{z,z'}} \right) + 1, \end{aligned} \tag{2}$$

$\forall z, z' \in \mathbb{Z}$. Here we are also not interested in the cases $a \cdot 1 = 1 \cdot a = a$ and $a \cdot (-1) = (-1) \cdot a = -a$. So we use $z, z' \in \mathbb{Z} \setminus \{-1, 0\}$.

4 Intersection

Ok. Now we have an equation to describe all integer divisible numbers of \mathbb{Z} . So we also write

$$\begin{aligned} Z_{z,z'} &= 2zz' + z + z' \\ &= (2z + 1)z' + z \\ &= (2z' + 1)z + z'. \end{aligned} \tag{3}$$

For example, the second line can be interpreted as the equation which gives us all $Z_{z,z'}$ which belongs to numbers which are the product of $(2z + 1)V$, $V \in \mathbb{Z}$.

We use this to write the following for all numbers which are not integer divisible by $(2z + 1)$:

$$Z_{z,z'} = (2z + 1)z' + z + \chi_z, \tag{4}$$

$\chi_z \in [1, (2z + 1) - 1]$. For the intersection of two of this equations follows

$$\begin{aligned} 0 &= Z_{z_1,z_1'} - Z_{z_2,z_2'} \\ &= (2z_1 + 1)z_1' - (2z_2 + 1)z_2' + \underbrace{z_1 + \chi_{z_1}}_{=: \kappa_{z_1}} - \left(\underbrace{z_2 + \chi_{z_2}}_{=: \kappa_{z_2}} \right) \\ &= (2z_1 + 1)z_1' - (2z_2 + 1)z_2' + \kappa_{z_1} - \kappa_{z_2}. \end{aligned} \tag{5}$$

References

- [1] ZÖBELEIN, C.: The recursive structure of the distribution of primes. In: *ArXiv e-prints*, 1411.2824 [*math.GM*] (2014), November