

PRIVATE RESEARCH PROJECT

The recursively calculation of prime numbers.

Draft/(Working) paper

*Carolin Zöbelein*¹

Available at

<https://github.com/Samdney/primescalc>

License: CC BY-ND 3.0 DE (see also LICENSE)

State: April 26, 2017

Keywords: Prime numbers, Primes, Recursive, Number Theory

Subjclass: 2010 *Mathematics Subject Classification*. Primary XX.

¹*E-mail address:* contact@carolin-zoebelein.de, *PGP Fingerprint:* D4A7 35E8 D47F 801F 2CF6
2BA7 927A FD3C DE47 E13B; *URL:* <http://www.carolin-zoebelein.de>

Abstract.

Roadmap

- ...

Contents

| | | |
|----------|---|-----------|
| 1 | Introduction | 5 |
| 2 | Odd-Divisible Numbers | 6 |
| 2.1 | Basic description: Odd-Numbers | 6 |
| 2.2 | Basic description: Odd-Divisible Numbers | 7 |
| 2.3 | Odd-Divisible Numbers: Different perspectives | 8 |
| 3 | Odd-Not-Divisible Numbers | 9 |
| 3.1 | Representation: Odd-Divisible Numbers | 9 |
| 3.2 | Representation: Odd-Not-Divisible Numbers | 10 |
| 3.3 | Odd-Not-Divisible Numbers: Intersection | 11 |
| 4 | The recursive calculation | 12 |
| 4.1 | Recursion step: $n^{(0)} = 0$ | 12 |

1 Introduction

In the following paper, I will show that prime numbers can be calculated recursively. I will start with the suggestion of descriptions itself, over different perspectives on this problem, until the final explanation of calculating prime numbers in the most efficient way, as a result from this considerations.

Let's start with the definition of prime numbers itself.

Definition 1.0.1 (Prime numbers) *Every natural number greater than one which has no positive integer divisors apart from one and itself is called Prime Number or just only Prime.*

Be \mathcal{P} the set of all prime numbers p . So we can write

$$\mathcal{P} := \{p \in \mathbb{N}_{>1} \mid \forall n \in \mathbb{N}_{>1} \setminus \{p\} : n \nmid p\}.$$

Hence, the first prime numbers are $\mathcal{P} := \{2, 3, 5, 7, 11, 13, 17, 19, 23, \dots\}$.

2 Odd-Divisible Numbers

Contents

| | | |
|------------|--|----------|
| 2.1 | Basic description: Odd-Numbers | 6 |
| 2.2 | Basic description: Odd-Divisible Numbers | 7 |
| 2.3 | Odd-Divisible Numbers: Different perspectives | 8 |

At first, for the description of prime numbers, we have to look at the set of divisible numbers. Since, apart from 2, all prime numbers are odd, we will only analyze this numbers. In the whole paper, we will ignore the prime number 2, because we will see, this makes a lot easier.

2.1 Basic description: Odd-Numbers

Be given the set of all odd natural numbers $y \in \mathbb{N}_{>1}$ through

$$y_i(x_i) := 2x_i + 1, \quad (2.1)$$

with $x, i \in \mathbb{N}$. If we expand the definition set of x to \mathbb{Z} , we also know

$$y(0) = 1 \quad (2.2)$$

$$\begin{aligned} \text{and } y(-x) &= 2(-x) + 1 \\ &= -(2x - 1) \\ &= -(2(x - 1) + 1) \\ &= -y(x - 1). \end{aligned} \quad (2.3)$$

Later, we will see that this properties can be very useful.

2.2 Basic description: Odd-Divisible Numbers

Next, we look at all odd-divisible numbers. We know, they can't have a factor which is a multiple of 2. Hence, we get an equation which describes all odd-divisible numbers by

$$\begin{aligned}
 y_{i,j}(x_i, x_j) &= y_i(x_i) \cdot y_j(x_j) \\
 &= (2x_i + 1)(2x_j + 1) \\
 &= 2^2 x_i x_j + 2x_i + 2x_j + 1 \\
 &= 2 \left(\underbrace{2x_i x_j + x_i + x_j}_{=: x_{i,j}} \right) + 1 \\
 &= y_{i,j}(x_{i,j}).
 \end{aligned} \tag{2.4}$$

If we expand again our sets to \mathbb{Z} , we receive additional cases. At first, assume at one factor is $y(0) = 1$. We see directly

$$\begin{aligned}
 y_{0,j}(0, x_j) &= y_0(0) \cdot y_j(x_j) \\
 &= 1 \cdot (2x_j + 1) \\
 &= 2x_j + 1 \\
 &= y_j(x_j)
 \end{aligned} \tag{2.5}$$

$$\text{respectively } y_{i,0}(x_i, 0) = y_i(x_i). \tag{2.6}$$

Next, assume we have one factor with $y(-x)$.

$$\begin{aligned}
 y_{i,j}(-x_i, x_j) &= y_i(-x_i) \cdot y_j(x_j) \\
 &= (2(-x_i) + 1)(2x_j + 1) \\
 &= -2^2 x_i x_j - 2x_i + 2x_j + 1 \\
 &= -(2(2x_i x_j + x_i - x_j - 1) + 1) \\
 &= -(2(2x_i x_j + x_i - 2x_j + x_j - 1) + 1) \\
 &= -(2(2(x_i - 1)x_j + (x_i - 1) + x_j) + 1) \\
 &= -y_i(x_i - 1) \cdot y_j(x_j)
 \end{aligned} \tag{2.7}$$

$$\text{respectively } y_{i,j}(x_i, -x_j) = -y_i(x_i) \cdot y_j(x_j - 1) \tag{2.8}$$

In the case of two negative factors, we have

$$\begin{aligned}
y_{i,j}(-x_i, -x_j) &= y_i(-x_i) \cdot y_j(-x_j) \\
&= (2(-x_i) + 1)(2(-x_j) + 1) \\
&= (2x_i - 1)(2x_j - 1) \\
&= 2^2 x_i x_j - 2x_i - 2x_j + 1 \\
&= 2(2x_i x_j - x_i - x_j) + 1 \\
&= (2x_i - 2 + 1)(2x_j - 2 + 1) \\
&= (2(x_i - 1) + 1)(2(x_j - 1) + 1) \\
&= (-1)y_i(x_i - 1)(-1)y_j(x_j - 1) \\
&= (-1)^2 y_{i,j}(x_i - 1, x_j - 1). \tag{2.9}
\end{aligned}$$

2.3 Odd-Divisible Numbers: Different perspectives

Finally, we see the different possible perspectives for odd-divisible numbers.

$$\begin{aligned}
y_{i,j}(x_i, x_j) &= 2(2x_i x_j + x_i + x_j) + 1 \\
&= 2((2x_i + 1)x_j + x_i) + 1 \tag{2.10}
\end{aligned}$$

$$\text{respectively} \quad = 2((2x_j + 1)x_i + x_j) + 1 \tag{2.11}$$

We will use (2.10) respectively (2.11) in the next step, for the description of odd numbers which are not divisible by a particular other odd number.

3 Odd-Not-Divisible Numbers

Contents

| | |
|--|-----------|
| 3.1 Representation: Odd-Divisible Numbers | 9 |
| 3.2 Representation: Odd-Not-Divisible Numbers | 10 |
| 3.3 Odd-Not-Divisible Numbers: Intersection | 11 |

After we spent time with the set of all odd-divisible numbers, now, we switch to the set of all odd numbers which are not divisible by a particular other odd number.

3.1 Representation: Odd-Divisible Numbers

Let's look again at (2.10)

$$y_{i,j}(x_i, x_j) = 2((2x_i + 1)x_j + x_i) + 1,$$

and its belonging values.

- Be $x_i = 1$:

$$y_{1,j}(1, x_j) = 2(3x_j + 1) + 1, \quad x_{1,j} = 3x_j + 1 \quad (3.1)$$

Table 3.1: The first ten values for (3.1).

| x_j | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|-----------|---|----|----|----|----|----|----|----|----|----|
| $x_{1,j}$ | 4 | 7 | 10 | 13 | 16 | 19 | 22 | 25 | 28 | 31 |
| $y_{1,j}$ | 9 | 15 | 21 | 27 | 33 | 39 | 45 | 51 | 57 | 63 |

- Be $x_i = 2$:

$$y_{2,j}(2, x_j) = 2(5x_j + 2) + 1, \quad x_{2,j} = 5x_j + 2 \quad (3.2)$$

Table 3.2: The first ten values for (3.2).

| x_j | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|-----------|----|----|----|----|----|----|----|----|----|-----|
| $x_{2,j}$ | 7 | 12 | 17 | 23 | 28 | 33 | 38 | 43 | 48 | 53 |
| $y_{2,j}$ | 15 | 25 | 35 | 47 | 57 | 67 | 77 | 87 | 97 | 107 |

- Be $x_i = 3$:

$$y_{3,j}(3, x_j) = 2(7x_j + 3) + 1, \quad x_{3,j} = 7x_j + 3 \quad (3.3)$$

Table 3.3: The first ten values for (3.3).

| x_j | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 |
|-----------|----|----|----|----|----|----|-----|-----|-----|-----|
| $x_{3,j}$ | 10 | 17 | 24 | 31 | 38 | 45 | 52 | 59 | 66 | 73 |
| $y_{3,j}$ | 21 | 35 | 49 | 63 | 77 | 91 | 105 | 119 | 133 | 147 |

- Be $x_i = \dots: \dots$

3.2 Representation: Odd-Not-Divisible Numbers

Now, we take again $y_{i,j}(x_i, x_j) = 2((2x_i + 1)x_j + x_i) + 1$ and rephrase it into an equation which describes all odd numbers which are not divisible by $2x_i + 1$.

That's not very hard. We can write

$$y_{i,j}(x_i, x_j) = 2((2x_i + 1)x_j + x_i - \mu(x_i)) + 1, \quad (3.4)$$

with

$$\mu(x_i) = 1, \dots, 2x_i, \quad \mu(x_i) \in \mathbb{N}. \quad (3.5)$$

Let's have a short look at the first values for $x_i = 1, 2, 3$.

- Be $x_i = 1$:

$$y_{1,j}(1, x_j) = 2(3x_j + 1 - \mu(1)) + 1, \quad \mu(1) = 1, 2, \quad x_{1,j} = 3x_j + 1 \quad (3.6)$$

Table 3.4: The first values for (3.6).

| x_j | 1 | 2 | 3 | 4 | 5 |
|-----------|---|---|----|----|----|
| $\mu(1)$ | 1 | 2 | 1 | 2 | 1 |
| $x_{1,j}$ | 2 | 3 | 5 | 6 | 8 |
| $y_{1,j}$ | 5 | 7 | 11 | 13 | 17 |

- Be $x_i = 2$:

$$y_{2,j}(2, x_j) = 2(5x_j + 2 - \mu(2)) + 1, \quad \mu(2) = 1, \dots, 4, \quad x_{2,j} = 5x_j + 1 \quad (3.7)$$

Table 3.5: The first values for (3.7).

| x_j | 1 | 2 | 3 | 4 | 5 |
|-----------|---|---|----|----|----|
| $\mu(2)$ | 1 | 2 | 3 | 4 | 1 |
| $x_{2,j}$ | 3 | 4 | 5 | 6 | 8 |
| $y_{2,j}$ | 7 | 9 | 11 | 13 | 17 |

- Be $x_i = 3$:

$$y_{3,j}(3, x_j) = 2(7x_j + 1 - \mu(3)) + 1, \quad \mu(3) = 1, \dots, 6 \quad x_{3,j} = 7x_j + 1 \quad (3.8)$$

Table 3.6: The first values for (3.8).

| | | | | | | | | | | |
|-----------|---|----|----|----|----|----|----|----|----|----|
| x_j | 1 | | | | | | | | | |
| $\mu(1)$ | 1 | 2 | 3 | 4 | 5 | 6 | 1 | 2 | 3 | 4 |
| $x_{3,j}$ | 4 | 5 | 6 | 7 | 8 | 9 | 11 | 12 | 13 | 14 |
| $y_{3,j}$ | 9 | 11 | 13 | 15 | 17 | 19 | 23 | 25 | 17 | 29 |

- Be $x_i = \dots: \dots$

Remark 3.2.1 (Value set) *You can see, the valid value set start not till $x_{i,j} = x_i + 1$.*

3.3 Odd-Not-Divisible Numbers: Intersection

Now we look at the intersection of two equations of the type (3.4) with (3.5). Hence, we start with

$$\begin{aligned} y_{i,j}^{(1)}(x_i^{(1)}, x_j^{(1)}) &= 2\left(\left(2x_i^{(1)} + 1\right)x_j^{(1)} + x_i^{(1)} - \mu\left(x_i^{(1)}\right)\right) + 1 \\ \mu\left(x_i^{(1)}\right) &= 1, \dots, 2x_i^{(1)} \end{aligned} \quad (3.9)$$

$$\begin{aligned} \text{and } y_{i,j}^{(2)}(x_i^{(2)}, x_j^{(2)}) &= 2\left(\left(2x_i^{(2)} + 1\right)x_j^{(2)} + x_i^{(2)} - \mu\left(x_i^{(2)}\right)\right) + 1 \\ \mu\left(x_i^{(2)}\right) &= 1, \dots, 2x_i^{(2)}. \end{aligned} \quad (3.10)$$

We do the intersection:

$$0 = \left(2x_i^{(1)} + 1\right)x_j^{(1)} - \left(2x_i^{(2)} + 1\right)x_j^{(2)} + x_i^{(1)} - x_i^{(2)} - \mu\left(x_i^{(1)}\right) + \mu\left(x_i^{(2)}\right) \quad (3.11)$$

$$\Leftrightarrow 0 = \left(2x_i^{(1)} + 1\right)\left(x_j^{(1)} - x_j^{(2)}\right) - \left(2x_j^{(2)} + 1\right)\Delta x_i^{(1,2)} - \mu\left(x_i^{(1)}\right) + \mu\left(x_i^{(2)}\right) \quad (3.12)$$

For the second one, we used $x_i^{(2)} = x_i^{(1)} + \Delta x_i^{(1,2)}$, $x_i^{(2)} > x_i^{(1)}$ and $\Delta x_i^{(1,2)} \in \mathbb{N}$.

To solve (3.11) respectively (3.12), we recognize that we have the boundary constraint, that $\left(2x_i^{(1)} + 1\right)$ and $\left(2x_i^{(2)} + 1\right)$ must not have any common factors.

4 The recursive calculation

Contents

| | |
|---|-----------|
| 4.1 Recursion step: $n^{(0)} = 0$ | 12 |
|---|-----------|

In this section, now, we do the final recursive calculation. To understand the deep structure we will do this by discussing the first steps by manually calculation. We will see, it exists different ways how you can look at each situation/problem in each step, hence this will mainly a discussion of this different ways.

4.1 Recursion step: $n^{(0)} = 0$

We start our calculation with an easy consideration. We assume, we know $3 = 2 \cdot 1 + 1$ and $5 = 2 \cdot 2 + 1$ are prime numbers. Hence, we know

$$\begin{aligned} y_{1,j}^{(1)} \left(1, x_j^{(1)} \right) &= 2 \left(3x_j^{(1)} + 1 - \mu(1) \right) + 1 \\ \mu(1) &= 1, 2 \end{aligned} \tag{4.1}$$

$$\begin{aligned} \text{and } y_{2,j}^{(2)} \left(2, x_j^{(2)} \right) &= 2 \left(5x_j^{(2)} + 2 - \mu(2) \right) + 1 \\ \mu(1) &= 1, 2, 3, 4 \end{aligned} \tag{4.2}$$

and want to calculate the intersection

$$3x_j^{(1)} + 1 - \mu(1) = 5x_j^{(2)} + 2 - \mu(2). \tag{4.3}$$

Now, we can choose between different angle of views, how we solve this intersection and how we handle the whole situation for our next recursion step.

Version 1:

The way to go ...

1. We fix: The whole set of valid $\mu(1) - \mu(2)$ values is allowed.
2. We determine the solutions for $x_j^{(1)}$ respectively $x_j^{(2)}$ without consideration of any definition ranges for x_j 's and the final range of interest.
3. ...
4. ...

The calculation ...

1. Trivial: $\mu(1) - \mu(2) = \{1, 2\} - \{1, 2, 3, 4\} = \{-3, \dots, 1\}$

- 2.

$$\begin{aligned} 3x_j^{(1)} - 5x_j^{(2)} &= 1 + \mu(1) - \mu(2) \\ x_j^{(1)} - x_j^{(2)} &= \frac{1}{3} \left(2x_j^{(2)} + 1 + \mu(1) - \mu(2) \right) \end{aligned} \quad (4.4)$$

The right side of (4.4) has to be an integer. We can directly see one valid solution. Be

$$x_j^{(2)} = 1 + \mu(1) - \mu(2) \quad (4.5)$$

which we put into (4.4) and receive

$$x_j^{(1)} = 2(1 + \mu(1) - \mu(2)). \quad (4.6)$$

Finally, we have all possible solutions for

$$x_j^{(1)} = 5z^{(1,2)} + 2(1 + \mu(1) - \mu(2)) \quad (4.7)$$

$$x_j^{(2)} = 3z^{(1,2)} + 1(1 + \mu(1) - \mu(2)), \quad (4.8)$$

$$z^{(1,2)} \in \mathbb{Z}.$$

3. ...

4. ...

List of Figures

List of Tables

| | | |
|-----|---|----|
| 3.1 | The first ten values for (3.1). | 9 |
| 3.2 | The first ten values for (3.2). | 9 |
| 3.3 | The first ten values for (3.3). | 10 |
| 3.4 | The first values for (3.6). | 10 |
| 3.5 | The first values for (3.7). | 10 |
| 3.6 | The first values for (3.8). | 11 |

Listings

Bibliography

Changelog

