

Tracing Baryons in the Warm Hot Intergalactic Medium using Broad Lyman- α Absorbers

Mid-Term 2

Sameer Patidar
SC19B161

Dual Degree (Astronomy & Astrophysics)
Indian Institute of Space Science and Technology

Supervisors : Dr. Vikram Khaire and Dr. Anand Narayanan

Thesis Phase I : Recap

Recap

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey

Recap

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey

Ref. : Fukugita et al. (1998)

Recap


- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey

Ref. : Fukugita et al. (1998)

Recap

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey

Ref. : Fukugita et al. (1998)
Shull et al. (2012)



Figures/Mid-term/Baryon_distrib

Figure 1: Baryon budget at $z \sim 0$.
Shull et al. (2012)

How to detect WHIM ?

How to detect WHIM ?

- ▶ Quasars as backlight

How to detect WHIM ?

- ▶ Quasars as backlight

Figures/Mid-term/spec slice.png

Figure 2: Slice of spectrum of quasar HE0153-4520 ($z_{em} = 0.4510$).
Danforth et al. (2016)

How to detect WHIM ?

- ▶ Quasars as backlight
 - O VI-VIII, Ne IX-X, N VII, etc.

Ref. : Tepper-García et al. (2013)
Savage et al. (2014)


Figures/Mid-term/spec slice.png

Figure 2: Slice of spectrum of quasar HE0153-4520 ($z_{em} = 0.4510$).
Danforth et al. (2016)

How to detect WHIM ?

- ▶ Quasars as backlight
 - O VI-VIII, Ne IX-X, N VII, etc.
 - **BLAs**

Ref. : Tepper-García et al. (2013)
Savage et al. (2014)



Figures/Mid-term/spec slice.png

Figure 2: Slice of spectrum of quasar HE0153-4520 ($z_{em} = 0.4510$).
Danforth et al. (2016)

How to detect WHIM ?

- ▶ Quasars as backlight
 - O VI-VIII, Ne IX-X, N VII, etc.
 - **BLAs**

Ref. : Tepper-García et al. (2013)
Savage et al. (2014)



Figures/Mid-term/BLA-individual.png

Figure 3: A BLA towards the LOS of quasar H 1821+643 ($z_{em} = 0.297$)
Philipp Richter (2005)

How to detect WHIM ?

- ▶ Quasars as backlight
 - O VI-VIII, Ne IX-X, N VII, etc.
 - **BLAs**

Ref. : Tepper-García et al. (2013)
Savage et al. (2014)

Figures/Mid-term/BLA.png

Figure 4: A BLA blended with other Ly α absorption lines towards the LOS of quasar PG1116+215 ($z_{em} = 0.176$). Philipp Richter (2020)

Data

HST/COS Data

HST/COS Data

- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)

HST/COS Data

- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)
- ▶ $z_{AGN} < 0.85$

HST/COS Data

- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)
- ▶ $z_{AGN} < 0.85$
- ▶ High S/N $\gtrsim 15$

HST/COS Data

- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)
- ▶ $z_{AGN} < 0.85$
- ▶ High S/N $\gtrsim 15$
- ▶ FUV channel : G130M and G160M gratings

HST/COS Data

- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)
- ▶ $z_{AGN} < 0.85$
- ▶ High S/N $\gtrsim 15$
- ▶ FUV channel : G130M and G160M gratings
- ▶ Coverage : 1130-1790 Å

HST/COS Data


- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)
- ▶ $z_{AGN} < 0.85$
- ▶ High S/N $\gtrsim 15$
- ▶ FUV channel : G130M and G160M gratings
- ▶ Coverage : 1130-1790 Å
- ▶ $R \sim 17,000 \approx 17 \text{ km s}^{-1}$

Phase I

Absorber system towards PG0003+158

Absorber system

- ▶ Quasar at $z_{em} = 0.45089$
- ▶ $z_{abs} \sim 0.347$
- ▶ 3 component system



Figures/Mid-term/Component_structure2.png

Voigt profile fitting

Figures/Mid-term/PG0003+158- $z=0.347579$ -sys-plot.png


Figure 5: System plot of the absorber system. Velocity is taken zero at $z = 0.347579$

Voigt profile fitting

- ▶ H I : 3 components
- ▶ O VI : 2 components
- ▶ C II, C III, Si II, Si III : 1 component

Figures/Mid-term/param.png

CLOUDY



Figures/Mid-term/cloudy-transparent.png

Figure 6: Schematic diagram of CLOUDY simulations.

Ionization Modelling

- ▶ Component I : -
- ▶ Component II : Hybrid - Collisional + Photo-ionization
- ▶ Component III : Photo-ionization (PI)

Component III : PI

Component III : PI

- ▶ Grid of CLOUDY models : Density and Metallicity
Ref. : Acharya and Khaire (2021)

Component III : PI

- ▶ Grid of CLOUDY models : Density and Metallicity
- ▶ $\log(R_{\text{eff}}/\text{cm}) = 5$ and 10 (step 2011, 02)

Component III : PI

- ▶ Grid of CLOUDY models : Density and Metallicity
- ▶ $\log(R_{\text{H}}/a_{\text{H}}^3)$: -5 to 1 in steps of 0.02
Ref: Anahaya and Khatri (2011)
- ▶ $\log(Z/Z_{\odot})$: -3 to 2 in steps of 0.05

Component III : PI

- ▶ Grid of CLOUDY models : Density and Metallicity
- ▶ $\log (N_{\text{H}} / \text{cm}^{-2})$: -5 to 0 in steps of 0.02
- ▶ $\log (Z / Z_{\odot})$: -3 to 2 in steps of 0.05
- ▶ Solution : Model that best matches the observed column densities





Figure 7: Modelled and observed column densities for the component III based on photoionization modelling


Component II : Hybrid

Component II : Hybrid



Figures/Mid-term/b3.png


Component II : Hybrid



Figures/Mid-term/b3.png

► $T = 10^{5.29^{+0.07}_{-0.08}} \text{ K}$


Component II : Hybrid



Figures/Mid-term/b3.png

- ▶ $T = 10^{5.29^{+0.07}_{-0.08}} \text{ K}$
- ▶ Constant temperature CLOUDY models

Component II : Hybrid



Figures/Mid-term/b3.png

- ▶ $T = 10^{5.29^{+0.07}_{-0.08}}$ K
- ▶ Constant temperature CLOUDY models
- ▶ O VI and size as constraining factors

Figures/Mid-term/physical-params.png

The Survey

Hunt for BLAs

Hunt for BLAs

- ▶ Broad $\text{Ly}\alpha$:

Hunt for BLAs

- ▶ Broad Ly α :

$$b \geq 45 \text{ km s}^{-1}$$

Hunt for BLAs

- ▶ Broad Ly α :

$$b \geq 45 \text{ km s}^{-1} \Rightarrow 568 \text{ systems}$$

Hunt for BLAs

- ▶ Broad Ly α :

$$b \geq 45 \text{ km s}^{-1} \Rightarrow 568 \text{ systems}$$

- ▶ Ionisation modelling :

Hunt for BLAs

- ▶ Broad Ly α :

$$b \geq 45 \text{ km s}^{-1} \Rightarrow 568 \text{ systems}$$

- ▶ Ionisation modelling :

$$\text{metal ions} \geq 3$$

Hunt for BLAs

- ▶ Broad Ly α :

$$b \geq 45 \text{ km s}^{-1} \Rightarrow 568 \text{ systems}$$

- ▶ Ionisation modelling :

$$\text{metal ions} \geq 3 \Rightarrow 28 \text{ systems}$$

Hunt for BLAs


- ▶ Broad Ly α :

$$b \geq 45 \text{ km s}^{-1} \Rightarrow 568 \text{ systems}$$

- ▶ Ionisation modelling :

$$\text{metal ions} \geq 3 \Rightarrow 28 \text{ systems}$$

28 BLA candidates



Figures/Mid-term/metal-ions.png

Figure 8: No. of different metal ions in all the 28 candidate BLAs

References

Acharya A., Khaire V., 2021, MNRAS, 509, 5559

Cen R., Ostriker J. P., 1999, ApJ, 514, 1

Danforth C. W., et al., 2016, ApJ, 817, 111

Fukugita M., Hogan C. J., Peebles P. J. E., 1998, ApJ, 503, 518

Savage B. D., Kim T. S., Wakker B. P., Keeney B., Shull J. M., Stocke J. T., Green J. C., 2014, ApJS, 212, 8

Shull J. M., Smith B. D., Danforth C. W., 2012, ApJ, 759, 23

Tepper-García T., Richter P., Schaye J., 2013, MNRAS, 436, 2063