

Tracing Baryons in the Warm Hot Intergalactic Medium using Broad Lyman- α Absorbers

Mid-Term 2

Sameer Patidar
SC19B161

Dual Degree (Astronomy & Astrophysics)
Indian Institute of Space Science and Technology

Supervisors : Dr. Vikram Khaire and Dr. Anand Narayanan

Thesis Phase I : Recap

Recap

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey : 28 BLA candidates

Figures/Baryon_distribution.png

Figure 1: Baryon budget at $z \sim 0$.
Shull et al. (2012)

Ionisation Modelling

Method

Method

- ▶ Grid of PI CLOUDY models : Density and Metallicity

Ref. : Acharya and Khaire (2021)

Method

- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶ $\log (n_{\text{H}}/\text{cm}^{-3})$: -5 to 1 in steps of 0.02

Ref. : Acharya and Khaire (2021)

Method

- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶ $\log (n_{\text{H}}/\text{cm}^{-3})$: -5 to 1 in steps of 0.02
- ▶ $\log (Z/Z_{\odot})$: -3 to 2 in steps of 0.05

Ref. : Acharya and Khaire (2021)

Method

- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶ $\log (n_{\text{H}}/\text{cm}^{-3})$: -5 to 1 in steps of 0.02
- ▶ $\log (Z/Z_{\odot})$: -3 to 2 in steps of 0.05
- ▶ Solution : Model that best predicts the observed column densities

Ref. : Acharya and Khaire (2021)

Scaling approximation

Scaling approximation

- ▶ 2d CLOUDY models : computationally expensive

Ref. : Acharya and Khaire (2021)

Scaling approximation

- ▶ 2d CLOUDY models : computationally expensive
- ▶ Scale column density with metallicity

Ref. : Acharya and Khaire (2021)

Scaling approximation


- ▶ 2d CLOUDY models : computationally expensive
- ▶ Scale column density with metallicity
- ▶ $N(n_H, Z) = N(n_H, Z_0) + \log(Z/Z_0)$

Ref. : Acharya and Khaire (2021)

Scaling approximation

- ▶ 2d CLOUDY models : computationally expensive
- ▶ Scale column density with metallicity
- ▶ $N(n_H, Z) = N(n_H, Z_0) + \log(Z/Z_0)$
- ▶ $Z_0 = 0.1Z_\odot$

Ref. : Acharya and Khaire (2021)



Figures/BLA-individual.png

Figure 2: A BLA towards the LOS of quasar H 1821+643 ($z_{em} = 0.297$)
Philipp Richter (2005)

How to detect WHIM ?

- ▶ Quasars as backlight
 - O VI-VIII, Ne IX-X, N VII, etc.
 - **BLAs**

Figures/BLA.png

Ref. : Tepper-García et al. (2013)
Savage et al. (2014)

Figure 3: A BLA blended with other Ly α absorption lines towards the LOS of quasar PG1116+215 ($z_{em} = 0.176$). Philipp Richter (2020)

Data

HST/COS Data

HST/COS Data

- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)

HST/COS Data

- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)
- ▶ $z_{AGN} < 0.85$

HST/COS Data

- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)
- ▶ $z_{AGN} < 0.85$
- ▶ High S/N $\gtrsim 15$

HST/COS Data

- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)
- ▶ $z_{AGN} < 0.85$
- ▶ High S/N $\gtrsim 15$
- ▶ FUV channel : G130M and G160M gratings

HST/COS Data

- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)
- ▶ $z_{AGN} < 0.85$
- ▶ High S/N $\gtrsim 15$
- ▶ FUV channel : G130M and G160M gratings
- ▶ Coverage : 1130-1790 Å

HST/COS Data

- ▶ 82 UV-bright AGNs chosen by Danforth et al. (2016)
- ▶ $z_{AGN} < 0.85$
- ▶ High S/N $\gtrsim 15$
- ▶ FUV channel : G130M and G160M gratings
- ▶ Coverage : 1130-1790 Å
- ▶ $R \sim 17,000 \approx 17 \text{ km s}^{-1}$

Phase I

Absorber system towards PG0003+158

Absorber system

- ▶ Quasar at $z_{em} = 0.45089$
- ▶ $z_{abs} \sim 0.347$
- ▶ 3 component system

Figures/Component_structure2.png

Voigt profile fitting

Figures/PG0003+158-z=0.347579-sys-plot.png

Figure 4: System plot of the absorber system. Velocity is taken zero at $z = 0.347579$

Voigt profile fitting

- ▶ H I : 3 components
- ▶ O VI : 2 components
- ▶ C II, C III, Si II, Si III : 1 component

Figures/param.png

CLOUDY


The image shows a large, empty rectangular box with a thin black border. Inside the box, the text 'Figures/cloudy-transparent.png' is written in a monospaced font. This likely represents a placeholder for a schematic diagram of CLOUDY simulations.

Figure 5: Schematic diagram of CLOUDY simulations.

Ionization Modelling

- ▶ Component I : -
- ▶ Component II : Hybrid - Collisional + Photo-ionization
- ▶ Component III : Photo-ionization (PI)

Component III : PI

Component III : PI

- ▶ Grid of CLOUDY models : Density and Metallicity

Ref. : Acharya and Khaire (2021)

Component III : PI

- ▶ Grid of CLOUDY models : Density and Metallicity
- ▶ $\log (n_{\text{H}}/\text{cm}^{-3})$: -5 to 1 in steps of 0.02

Ref. : Acharya and Khaire (2021)

Component III : PI

- ▶ Grid of CLOUDY models : Density and Metallicity
- ▶ $\log (n_{\text{H}}/\text{cm}^{-3})$: -5 to 1 in steps of 0.02
- ▶ $\log (Z/Z_{\odot})$: -3 to 2 in steps of 0.05

Ref. : Acharya and Khaire (2021)

Component III : PI

- ▶ Grid of CLOUDY models : Density and Metallicity
- ▶ $\log (n_{\text{H}}/\text{cm}^{-3})$: -5 to 1 in steps of 0.02
- ▶ $\log (Z/Z_{\odot})$: -3 to 2 in steps of 0.05
- ▶ Solution : Model that best matches the observed column densities

Ref. : Acharya and Khaire (2021)





Figure 6: Modelled and observed column densities for the component III based on photoionization modelling


Component II : Hybrid

Component II : Hybrid



Figures/b3.png


Component II : Hybrid



Figures/b3.png

► $T = 10^{5.29^{+0.07}_{-0.08}} \text{ K}$


Component II : Hybrid



Figures/b3.png

- ▶ $T = 10^{5.29^{+0.07}_{-0.08}} \text{ K}$
- ▶ Constant temperature CLOUDY models

Component II : Hybrid



Figures/b3.png

- ▶ $T = 10^{5.29^{+0.07}_{-0.08}} \text{ K}$
- ▶ Constant temperature CLOUDY models
- ▶ O VI and size as constraining factors

Figures/physical-params.png

The Survey

Hunt for BLAs


Hunt for BLAs

- ▶ Broad Ly α :

Hunt for BLAs

- ▶ Broad Ly α :


$$b \geq 45 \text{ km s}^{-1}$$



Hunt for BLAs

- ▶ Broad Ly α :

$$b \geq 45 \text{ km s}^{-1} \quad \Rightarrow \quad 568 \text{ systems}$$




Hunt for BLAs

- ▶ Broad Ly α :

$$b \geq 45 \text{ km s}^{-1} \quad \Rightarrow \quad 568 \text{ systems}$$

- ▶ Ionisation modelling :



Hunt for BLAs

- ▶ Broad Ly α :

$$b \geq 45 \text{ km s}^{-1} \Rightarrow 568 \text{ systems}$$

- ▶ Ionisation modelling :

$$\text{metal ions} \geq 3$$

Hunt for BLAs

- ▶ Broad Ly α :

$b \geq 45 \text{ km s}^{-1} \Rightarrow 568 \text{ systems}$

- ▶ Ionisation modelling :

metal ions $\geq 3 \Rightarrow 28 \text{ systems}$

Hunt for BLAs


- ▶ Broad Ly α :

$b \geq 45 \text{ km s}^{-1} \Rightarrow 568 \text{ systems}$

- ▶ Ionisation modelling :

metal ions $\geq 3 \Rightarrow 28 \text{ systems}$

28 BLA candidates



Figures/metal-ions.png

Figure 7: No. of different metal ions in all the 28 candidate BLAs

References

Acharya A., Khaire V., 2021, MNRAS, 509, 5559

Cen R., Ostriker J. P., 1999, ApJ, 514, 1

Danforth C. W., et al., 2016, ApJ, 817, 111

Fukugita M., Hogan C. J., Peebles P. J. E., 1998, ApJ, 503, 518

Savage B. D., Kim T. S., Wakker B. P., Keeney B., Shull J. M., Stocke J. T., Green J. C., 2014, ApJS, 212, 8

Shull J. M., Smith B. D., Danforth C. W., 2012, ApJ, 759, 23

Tepper-García T., Richter P., Schaye J., 2013, MNRAS, 436, 2063