Tracing Baryons in the Warm Hot Intergalactic Medium using Broad Lyman- α Absorbers

Mid-Term 2

Sameer Patidar SC19B161

Dual Degree (Astronomy & Astrophysics)
Indian Institute of Space Science and Technology

Supervisors: Dr. Vikram Khaire and Dr. Anand Narayanan

Thesis Phase I: Recap

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey : 28 BLA candidates




Figure 1: Baryon budget at $z \sim 0$. Shull et al. (2012)

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- Absorber towards PG 0003+158
- ▶ BLA survey : 28 BLA candidates

Figures/BLA-individual.png

Figure 2: A BLA towards the LOS of quasar H 1821+643. Philipp Richter (2005)

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- Absorber towards PG 0003+158
- ▶ BLA survey : 28 BLA candidates

Figures/PG0003+158-z=0.347579-sys-plot.png

Figure 3: System plot of the absorber system towards PG 0003+158. Velocity is taken zero at z = 0.347579

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey : 28 BLA candidates

Figures/metal-ions.png

Figure 4: Distribution of metal ions in all 28 candidate BLAs

The BLA Survey

Survey so far...

Survey so far...

▶ Voigt profile fitting : 16 (O VI) + 6 systems

Survey so far...

- ▶ Voigt profile fitting : 16 (O VI) + 6 systems
- ► Ionisation Modelling : **16 (O VI)** systems

Insights




Figure 5: Distribution of H I column densities and Doppler parameters.

Insights

```
Figures/placeholder.png
```

Figure 6: H I column density vs. Doppler parameter

Insights

```
Figures/placeholder.png
```

Figure 7: b(O VI) vs. b(H I)

Ionisation Modelling

▶ Grid of PI CLOUDY models : Density and Metallicity

- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶ $\log (n_H/cm^{-3})$: -5 to 1 in steps of 0.02

- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶ $log (n_H/cm^{-3})$: -5 to 1 in steps of 0.02
- ▶ $\log (Z/Z \odot)$: -3 to 2 in steps of 0.05

- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶ $\log (n_H/cm^{-3})$: -5 to 1 in steps of 0.02
- ▶ $\log (Z/Z \odot)$: -3 to 2 in steps of 0.05
- Solution : Model that best predicts the observed column densities

2d CLOUDY models : computationally expensive

- 2d CLOUDY models : computationally expensive
- Scale column density with metallcity

- 2d CLOUDY models : computationally expensive
- Scale column density with metallcity
- $N(n_H, Z) = N(n_H, Z_0) + \log(Z/Z_0)$

- 2d CLOUDY models : computationally expensive
- Scale column density with metallcity
- $N(n_H, Z) = N(n_H, Z_0) + \log(Z/Z_0)$
- ► $Z_0 = 0.1Z$ ⊙




Figure 8: Column density of ion vs. Hydrogen densities at various metallcities

Results

```
Figures/placeholder.png
```

Figure 9: Distribution of no. of metal ions in the absorber systems

Results

```
Figures/placeholder.png
```

Figure 10: Distribution of no. of metal ions in the absorber systems

Figures/placeholder.png

Figure 11: System plot of the absorber system towards LOS of




Figure 12: Observed and predicted column densities of the ions.

References

Acharva A., Khaire V., 2021, MNRAS, 509, 5559

Cen R., Ostriker J. P., 1999, ApJ, 514, 1

Danforth C. W., et al., 2016, ApJ, 817, 111

Fukugita M., Hogan C. J., Peebles P. J. E., 1998, ApJ, 503, 518

Savage B. D., Kim T. S., Wakker B. P., Keeney B., Shull J. M., Stocke J. T., Green J. C., 2014, ApJS, 212, 8

Shull J. M., Smith B. D., Danforth C. W., 2012, ApJ, 759, 23

Tepper-García T., Richter P., Schaye J., 2013, MNRAS, 436, 2063