Tracing Baryons in the Warm Hot Intergalactic Medium using Broad Lyman- α Absorbers

Mid-Term 2

Sameer Patidar SC19B161

Dual Degree (Astronomy & Astrophysics)
Indian Institute of Space Science and Technology

Supervisors: Dr. Vikram Khaire and Dr. Anand Narayanan





Thesis Phase I: Recap

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey : 28 BLA candidates




Figure 1: Baryon budget at $z \sim 0$. Shull et al. (2012)

- ▶ The missing baryon problem
- **▶** BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey : 28 BLA candidates




Figure 2: A BLA towards the LOS of quasar H 1821+643. Philipp Richter (2005)

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ► Absorber towards PG 0003+158
- ▶ BLA survey : 28 BLA candidates




Figure 3: System plot of the absorber system towards PG 0003+158. Velocity is taken zero at z = 0.347579

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey : 28 BLA candidates




Figure 4: Distribution of metal ions in all 28 candidate BLAs

The BLA Survey

Survey so far...

Survey so far...

▶ Voigt profile fitting : 16 (O VI) + 6

Survey so far...

- ▶ Voigt profile fitting : 16 (O VI) + 6
- ► Ionisation Modelling : **16 (O VI)**




Figure 5: Distribution of H I column densities and Doppler parameters.




Figure 6: H I column density vs. Doppler parameter






Figure 7: b(O VI) vs. b(H I). Grey filled circles are measurements from Danforth et. al 2016.






Figure 8: Distribution of temperature calculated from Doppler parameters of H I and O VI lines.

Ionisation Modelling



▶ Grid of PI CLOUDY models : Density and Metallicity

- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶ $\log (n_H/cm^{-3})$: -5 to 1 in steps of 0.02



- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶ $\log (n_H/cm^{-3})$: -5 to 1 in steps of 0.02
- ▶ $\log (Z/Z \odot)$: -3 to 2 in steps of 0.05

- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶ $\log (n_H/cm^{-3})$: -5 to 1 in steps of 0.02
- ▶ $\log (Z/Z \odot)$: -3 to 2 in steps of 0.05
- Solution : Model that best predicts the observed column densities





▶ 16 O VI absorbers



- ▶ 16 O VI absorbers
- ▶ 26 components



- ▶ 16 O VI absorbers
- ▶ 26 components
- ▶ Origin of O VI



Solutions




Figure 9: Ionisation modelling solutions (n_H, Z) for all 26 components.



+ve correlation




Figure 10: Variation of N(H I) with n_H



O VI cases




Figure 11: O VI column density predictions.



Ex-1




Figure 12: $N(H I)=16.49 cm^{-2}$

Ex-2




Figure 13: $N(H I)=12.76 \text{ cm}^{-2}$

Ex-3




Figure 14: $N(H I)=13.46 cm^{-2}$

Towards the end

Voigt profile fitting : 6



- ▶ Voigt profile fitting : 6
- ► Ionisation modelling : 12



- ▶ Voigt profile fitting : 6
- ▶ Ionisation modelling : 12
- Exploring the survey results



- ▶ Voigt profile fitting : 6
- ▶ Ionisation modelling : 12
- Exploring the survey results
- *Finally*, calculating $\Omega_b(BLA)$

Summary

- Voigt profile fitting: 22 absorbers Voigt profiles
- ▶ Ionisation modelling : 16 absorbers 26 components
- ▶ O VI couldn't be explained with photoionization models
- BLA survey towards completion

References

Acharya A., Khaire V., 2021, MNRAS, 509, 5559

Danforth C. W., et al., 2016, ApJ, 817, 111

Lehner N., Savage B. D., Richter P., Sembach K. R., Tripp T. M., Wakker B. P., 2007, ApJ, 658, 680

Richter P., Savage B. D., Sembach K. R., Tripp T. M., 2006, A&A, 445, 827

Savage B. D., Kim T. S., Wakker B. P., Keeney B., Shull J. M., Stocke J. T., Green J. C., 2014, ApJS, 212, 8

So much universe, and so little time...