

# Tracing Baryons in the Warm Hot Intergalactic Medium using Broad Lyman- $\alpha$ Absorbers

Mid-Term 2

Sameer Patidar  
SC19B161

Dual Degree (Astronomy & Astrophysics)  
Indian Institute of Space Science and Technology

Supervisors : Dr. Vikram Khaire and Dr. Anand Narayanan



# Thesis Phase I : Recap

# Recap

- ▶ **The missing baryon problem**
- ▶ BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey : 28 BLA candidates





Figure 1: Baryon budget at  $z \sim 0$ .  
Shull et al. (2012)

# Recap

- ▶ The missing baryon problem
- ▶ **BLAs : Way to probe WHIM**
- ▶ Absorber towards PG 0003+158
- ▶ BLA survey : 28 BLA candidates



**Figure 2:** A BLA towards the LOS of quasar H 1821+643.  
Philipp Richter (2005)

# Recap

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ▶ **Absorber towards PG 0003+158**
- ▶ BLA survey : 28 BLA candidates




Figure 3: System plot of the absorber system towards PG 0003+158. Velocity is taken zero at  $z = 0.347579$

# Recap

- ▶ The missing baryon problem
- ▶ BLAs : Way to probe WHIM
- ▶ Absorber towards PG 0003+158
- ▶ **BLA survey : 28 BLA candidates**




Figure 4: Distribution of metal ions in all 28 candidate BLAs

# The BLA Survey

# Survey so far...



# Survey so far...

- ▶ Voigt profile fitting : 16 (O VI) + 6

# Survey so far...

- ▶ Voigt profile fitting : 16 (O VI) + 6
- ▶ Ionisation Modelling : **16 (O VI)**

# Insights




Figure 5: Distribution of H I column densities and Doppler parameters.

# Insights




Figure 6: H I column density vs. Doppler parameter

# Insights

$$b^2 = b_{th}^2 + b_{nt}^2$$

$$b_{th}^2 = \frac{2kT}{m}$$




Figure 7:  $b(\text{O VI})$  vs.  $b(\text{H I})$ . Grey filled circles are measurements from Danforth et. al 2016.

# Insights

$$T = \frac{8m}{15k} (b^2_{\text{HI}} - b^2_{\text{OVI}})$$




Figure 8: Distribution of temperature calculated from Doppler parameters of HI and OVI lines.

# Ionisation Modelling

# Method



# Method

- ▶ Grid of PI CLOUDY models : Density and Metallicity

Ref. : Acharya and Khaire (2021)

# Method

- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶  $\log (n_{\text{H}}/\text{cm}^{-3})$  : -5 to 1 in steps of 0.02

Ref. : Acharya and Khaire (2021)

# Method

- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶  $\log (n_{\text{H}}/\text{cm}^{-3})$  : -5 to 1 in steps of 0.02
- ▶  $\log (Z/Z_{\odot})$  : -3 to 2 in steps of 0.05

Ref. : Acharya and Khaire (2021)

# Method

- ▶ Grid of PI CLOUDY models : Density and Metallicity
- ▶  $\log (n_{\text{H}}/\text{cm}^{-3})$  : -5 to 1 in steps of 0.02
- ▶  $\log (Z/Z_{\odot})$  : -3 to 2 in steps of 0.05
- ▶ Solution : Model that best predicts the observed column densities

Ref. : Acharya and Khaire (2021)

# Results

# Results

- ▶ 16 O VI absorbers

# Results

- ▶ 16 O VI absorbers
- ▶ 26 components

# Results

- ▶ 16 O VI absorbers
- ▶ 26 components
- ▶ Origin of O VI



# Solutions




Figure 9: Ionisation modelling solutions ( $n_H$ ,  $Z$ ) for all 26 components.

## +ve correlation




Figure 10: Variation of  $N(\text{H I})$  with  $n_{\text{H}}$

## O VI cases




Figure 11: O VI column density predictions.

# Ex : Unexplained




Figure 12:  $N(\text{H I}) = 16.49 \text{ cm}^{-2}$

# Ex : Explained





Figure 13:  $N(\text{H I}) = 12.76 \text{ cm}^{-2}$

## Ex : Uncertain

Figure 14:  $N(\text{H I}) = 13.46 \text{ cm}^{-2}$

# Towards *the end*

# Ongoing and Future Work



# Ongoing and Future Work

- ▶ Voigt profile fitting : 6

# Ongoing and Future Work

- ▶ Voigt profile fitting : 6
- ▶ Ionisation modelling : 12

# Ongoing and Future Work

- ▶ Voigt profile fitting : 6
- ▶ Ionisation modelling : 12
- ▶ Exploring the survey results

# Ongoing and Future Work

- ▶ Voigt profile fitting : 6
- ▶ Ionisation modelling : 12
- ▶ Exploring the survey results
- ▶ *Finally*, calculating  $\Omega_b(BLA)$

# Summary

- ▶ Voigt profile fitting : 22 absorbers - 231 Voigt profiles
- ▶ Ionisation modelling : 16 absorbers - 26 components
- ▶ O VI couldn't be explained with photoionization models
- ▶ BLA survey towards completion

# References

Acharya A., Khaire V., 2021, MNRAS, 509, 5559

Danforth C. W., et al., 2016, ApJ, 817, 111

Lehner N., Savage B. D., Richter P., Sembach K. R., Tripp T. M., Wakker B. P., 2007, ApJ, 658, 680

Richter P., Savage B. D., Sembach K. R., Tripp T. M., 2006, A&A, 445, 827

Savage B. D., Kim T. S., Wakker B. P., Keeney B., Shull J. M., Stocke J. T., Green J. C., 2014, ApJS, 212, 8

*So much universe, and so little  
time...*