A Machine-Checked Proof that the Kerr–Newman Spacetime Solves the Einstein–Maxwell Equations

Samen Hossain

October 2026

Abstract

We present a complete formal proof, in the Lean 4 proof assistant with the Mathlib library, that the Kerr–Newman metric and its electromagnetic field satisfy the Einstein–Maxwell equations. The statement is phrased in Kerr–Schild Cartesian coordinates on $\mathbb{R}^4$. It holds for all real mass, spin and charge parameters (m, a, e), at every point where the Kerr–Schild radial function is positive. It has three parts: the Einstein equation $G_{\mu\nu} = 8\pi T_{\mu\nu}$ with the electromagnetic stress–energy tensor, the source-free Maxwell equation $\partial_\mu(\sqrt{|\det g|}\, F^{\mu\nu}) = 0$, and closedness of F. Christoffel symbols, curvature and all derivatives are defined from first principles through Fréchet derivatives. The proof uses no axioms beyond the three standard ones of Lean.

The obstacle is the size of the curvature computation, which computer algebra performs routinely but which is far beyond what can be written by hand in a proof assistant. We overcome it by expressing every quantity as a polynomial in five *atoms*: the radial function r, the spatial components of the Kerr–Schild null covector, and $W = 1/(r^2 + a^2\cos^2\theta)$. The atoms are closed under differentiation and satisfy two polynomial relations. This turns each step of the curvature computation into an identity in a quotient ring. Each identity is certified by cofactors from Gröbner-basis division, computed offline with SymPy and checked in Lean by `linear_combination`, so the computer algebra system is never trusted. The Maxwell equations are proved conceptually: the Kerr–Schild structure reduces the curved equation to the flat one, which follows from the harmonicity of Appell's complex-shifted Coulomb potential. We describe the method, the engineering needed to fit about one megabyte of mostly generated Lean into a 300-second per-file checking budget, and the trusted base of the result. The formalization was produced by an AI coding agent on the Prove2Me platform under the author's direction, and we document that process.

Contents

1 Introduction **2**

2 The formal statement **4**
2.1 Coordinate Lorentzian geometry 4
2.2 The Kerr–Newman solution in Kerr–Schild form 5
2.3 The Lean statement 6

3 Overview of the proof **6**

4 Kerr–Schild algebra in atoms **6**
4.1 Atoms and relations 6
4.2 Derivatives of the atoms 7
4.3 Kerr–Schild identities 7

5 Certified polynomial identities **8**
5.1 Ideal membership and Gröbner cofactors 8
5.2 Replay in Lean 8
5.3 Why the extra atom W 8

6 Derivatives: a small calculus of coordinate partials **9**

7 The Einstein equation **10**

8 The Maxwell equations **11**
8.1 Closedness 11
8.2 From curved to flat space 11
8.3 The flat Maxwell equation 11

9 Engineering a large proof under a time budget **12**
9.1 Platform constraints 12
9.2 Lessons 12

10 Trusted base and validation **13**

11 Related work **14**

12 Conclusion **15**

A The platform definitions **19**

B Platform objects **21**

1 Introduction

In 1963 Kerr found the metric of a rotating black hole [27]. Two years later Newman and collaborators found its electrically charged generalization [37]. The resulting Kerr–Newman family is labelled by a mass m, a spin parameter a (angular momentum per unit mass) and a charge e. It plays a distinguished role because of the black-hole uniqueness theorems [6, 8, 22–24, 32, 42]. Under suitable regularity and global assumptions, these theorems show that Kerr–Newman metrics exhaust the stationary, asymptotically flat black-hole solutions of the Einstein–Maxwell equations. This is the "no-hair" property; see the review [10] for the precise hypotheses and for the questions that remain open.

The most basic fact about an exact solution is that it actually solves the field equations. For Kerr–Newman this is a substantial calculation. One computes forty independent Christoffel symbols, their derivatives and the Ricci tensor, and then compares the result with the electromagnetic stress–energy tensor. Every intermediate quantity is a sizeable rational function of the coordinates. In practice the check is left to computer algebra systems [30]. These are reliable, but they do not produce a proof that can be checked independently. Interactive theorem provers such as Lean [12] reduce trust to a small proof-checking kernel. So far they have been used little for heavy computations in general relativity (Section 11).

Result. We report a complete formal proof of the following statement. The precise Lean statement and all the definitions it uses are reproduced in Section 2 and Appendix A.

Theorem 1.1 (informal)**.** *Let $m, a, e \in \mathbb{R}$. Let g be the Kerr–Newman metric in Kerr–Schild coordinates, and let $F = dA$ be the Kerr–Newman electromagnetic field. Then at every point $p \in \mathbb{R}^4$ where the Kerr–Schild radius satisfies $r(p) > 0$:*

(i) $G_{\mu\nu} = 2\big(F_{\mu\alpha}F_{\nu}{}^{\alpha} - \frac{1}{4}\, g_{\mu\nu}F_{\alpha\beta}F^{\alpha\beta}\big)$*, that is,* $G_{\mu\nu} = 8\pi T_{\mu\nu}$ *in Gaussian geometrized units;*

(ii) $\partial_\mu\big(\sqrt{|\det g|}\, F^{\mu\nu}\big) = 0$ *for every* ν*;*

(iii) $\partial_\lambda F_{\mu\nu} + \partial_\mu F_{\nu\lambda} + \partial_\nu F_{\lambda\mu} = 0$ *for all* λ, μ, ν*.*

The theorem places no restriction on the parameters. It therefore covers black holes ($a^2+e^2 \le m^2$) as well as over-extremal and negative-mass members of the family. It holds everywhere in the Kerr–Schild chart except on the disc spanned by the ring singularity, including inside the horizons. All derivatives are genuine Fréchet derivatives, the inverse metric is the matrix inverse, and curvature follows the textbook coordinate formulas. The kernel accepts the proof with only the axioms `propext`, `Classical.choice` and `Quot.sound`.

Contributions.

- A complete, kernel-checked proof of Theorem 1.1 in Lean 4 and Mathlib [47] (Sections 7 and 8). To our knowledge, no earlier machine-checked proof shows that Kerr–Newman, Kerr or even Schwarzschild satisfies the Einstein equations with the curvature computed from the metric by the prover's own calculus (Section 11).
- A method for curvature computations in a proof assistant that keeps the computer algebra system outside the trusted base (Sections 4 to 6). Every quantity is written in a small set of algebraically constrained *atoms* that is closed under differentiation. Each identity is then a polynomial ideal-membership problem, certified by Gröbner-basis cofactors and replayed by Lean's `linear_combination` tactic.
- A short conceptual proof of the Maxwell equations (Section 8). It rests on two classical facts: the reduction of the curved equation to the flat one for Kerr–Schild fields aligned with k, and Appell's interpretation of the Kerr–Newman potential as a Coulomb potential centred at an imaginary point.
- An account of the engineering needed to make a proof of this size check under a strict time budget (Section 9), together with a discussion of the trusted base (Section 10).

Provenance. The work was carried out on Prove2Me [41], a platform where formal statements are posted and submitted Lean proofs are checked on the server. Theorem 1.1 is the milestone `NoHair.kerrNewman_einsteinMaxwell` of the platform's mission *No-Hair Conjecture: Uniqueness of Stationary Kerr–Newman Black Holes*. The mission authors fixed the formal statement and all the definitions it depends on before this work began. We did not change them. The proofs, the code generators and a first draft of this paper were written by an AI coding agent (Claude Code, by Anthropic) working under the author's direction. The Lean kernel checked every proof both on the platform and locally (Section 10). We say this up front because it bears on how the artefact should be read. The Lean files are large, machine-generated and not intended for line-by-line human reading. Their correctness rests on the kernel, while their *meaning* rests on the definitions, which are short and are reproduced in full in Appendix A.

2 The formal statement

2.1 Coordinate Lorentzian geometry

The platform formalizes Lorentzian geometry in a single global chart. Coordinate space is $\mathbb{R}^4$ (`E4 := EuclideanSpace ℝ (Fin 4)`), coordinate 0 is time, and $\eta = \mathrm{diag}(-1, 1, 1, 1)$. A metric is a matrix-valued function $g : \mathbb{R}^4 \to \mathbb{R}^{4\times 4}$. The following definitions are verbatim transcriptions of Appendix A; the Lean names are in typewriter font.

- Coordinate partial derivative (`coordPartial`): $\partial_k f(p) := Df(p)(e_k)$, the Fréchet derivative of $f : \mathbb{R}^4 \to \mathbb{R}$ at p applied to the k-th basis vector.
- Inverse metric (`invMetric`): $g^{\mu\nu}(p) := (g(p))^{-1}_{\mu\nu}$, the matrix inverse.
- Christoffel symbols (`christoffel`): $\Gamma^i_{jk} := \frac{1}{2}\sum_l g^{il}(\partial_j g_{lk} + \partial_k g_{lj} - \partial_l g_{jk})$.
- Riemann tensor (`riemann`): $R^\rho{}_{\sigma\mu\nu} := \partial_\mu \Gamma^\rho_{\nu\sigma} - \partial_\nu \Gamma^\rho_{\mu\sigma} + \sum_\lambda (\Gamma^\rho_{\mu\lambda}\Gamma^\lambda_{\nu\sigma} - \Gamma^\rho_{\nu\lambda}\Gamma^\lambda_{\mu\sigma})$.
- Ricci tensor, scalar curvature and Einstein tensor:

$$R_{\sigma\nu} := \sum_\rho R^\rho{}_{\sigma\rho\nu}, \qquad R := \sum_{\sigma,\nu} g^{\sigma\nu} R_{\sigma\nu}, \qquad G_{\mu\nu} := R_{\mu\nu} - \tfrac{1}{2} R\, g_{\mu\nu}.$$

  In Lean these are `ricci`, `scalarCurvature` and `einsteinTensor`.

- Electromagnetic source (`raiseBoth`, `maxwellSource`): $F^{\alpha\beta} := g^{\alpha\mu} g^{\beta\nu} F_{\mu\nu}$ and

$$S_{\mu\nu} := 2\Big(\sum_{\alpha,\beta} F_{\mu\alpha}\, g^{\alpha\beta} F_{\nu\beta} - \tfrac{1}{4}\, g_{\mu\nu} \sum_{\alpha,\beta} F_{\alpha\beta} F^{\alpha\beta} \Big).$$

  With $T_{\mu\nu} = \frac{1}{4\pi}(F_{\mu\alpha}F_\nu{}^\alpha - \frac{1}{4} g_{\mu\nu} F_{\alpha\beta}F^{\alpha\beta})$ in Gaussian units with $G = c = 1$, this is $S_{\mu\nu} = 8\pi T_{\mu\nu}$.

- The Einstein–Maxwell system at a point (`EinsteinMaxwellAt`) is the conjunction of (i)–(iii) in Theorem 1.1.

These are the conventions of Misner, Thorne and Wheeler [34]: signature $(-,+,+,+)$, Riemann tensor with the sign that gives round spheres positive scalar curvature, and $G_{\mu\nu} = +8\pi T_{\mu\nu}$. Wald [52] places the indices of the Riemann tensor differently, but his Ricci tensor, and hence R and $G_{\mu\nu}$, agree with the ones here. As a sanity check on the definitions themselves, we evaluated them symbolically on round spheres (positive scalar curvature) and on the Reissner–Nordström solution with $-g_{tt} = 1 - 2M/r + Q^2/r^2$, which satisfies the platform's Einstein equation exactly with positive energy density.

Remark 2.1 (Total functions). Lean's mathematical functions are total. The Fréchet derivative of a non-differentiable function is 0, the inverse of a singular matrix is 0, and $\sqrt{x} = 0$ for $x < 0$. A statement about derivatives could therefore hold for degenerate reasons. This does not happen here. On the region $r > 0$ we prove $\det g = -1$ and identify the inverse explicitly (Lemma 4.2), and every denominator is shown to be nonzero. Every spatial derivative used in the proof comes with an explicit proof of differentiability (Section 6). The only exceptions are time derivatives, which vanish by translation invariance whether or not the function is differentiable (Lemma 6.1). Every function involved is in any case smooth on the region, so all quantities in the formal statement have their intended meaning.

2.2 The Kerr–Newman solution in Kerr–Schild form

Write a point of $\mathbb{R}^4$ as $p = (t, x, y, z)$ and let $\rho^2 = x^2 + y^2 + z^2$. (In the Lean code the point itself is called `y` and its coordinates are `y 0`, …, `y 3`.) The Kerr–Schild radius (`ksRadius`) is

$$r := \sqrt{\tfrac{1}{2}\Big(\rho^2 - a^2 + \sqrt{(\rho^2 - a^2)^2 + 4a^2 z^2}\Big)}\,. \tag{1}$$

It depends on a only through a^2. Where $r > 0$ it is the unique positive root of

$$\frac{x^2 + y^2}{r^2 + a^2} + \frac{z^2}{r^2} = 1, \qquad \text{equivalently} \qquad r^4 - (\rho^2 - a^2)\, r^2 - a^2 z^2 = 0. \tag{2}$$

The Kerr–Schild covector (`ksNull`), profile (`ksProfile`), metric (`kerrNewmanMetric`), potential (`kerrNewmanPotential`) and field (`kerrNewmanField`) are

$$k_\mu = \Big(1,\ \frac{rx + ay}{r^2 + a^2},\ \frac{ry - ax}{r^2 + a^2},\ \frac{z}{r}\Big), \qquad f = \frac{r^2(2mr - e^2)}{r^4 + a^2 z^2}, \tag{3}$$

$$g_{\mu\nu} = \eta_{\mu\nu} + f\, k_\mu k_\nu, \qquad A_\mu = \frac{e\, r^3}{r^4 + a^2 z^2}\, k_\mu, \qquad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu. \tag{4}$$

With $z = r\cos\theta$ we have $r^4 + a^2 z^2 = r^2 \Sigma$, where $\Sigma = r^2 + a^2 \cos^2\theta$. Then $f = (2mr - e^2)/\Sigma$ and $A = (er/\Sigma)\, k$, which is the familiar Kerr–Schild form of Kerr–Newman [1, 13, 51]. Setting $e = 0$ gives the Kerr metric. Setting $a = e = 0$ gives Schwarzschild in ingoing Kerr–Schild (Eddington–Finkelstein-type) coordinates.

The region $U = \{r > 0\}$. By (1), $r = 0$ exactly when $z = 0$ and $x^2 + y^2 \le a^2$. Hence, for $a \neq 0$, U is the complement of $\mathbb{R} \times D$, where D is the closed disc whose boundary is the ring singularity. For $a = 0$, U is the complement of the world line $x = y = z = 0$. In either case U is open, and on U the function r is real-analytic. Removing the disc is not an artefact of the formalization: $k_3 = z/r$ changes sign across the open disc, so for $e \neq 0$ the metric is discontinuous there, and for $e = 0$, $m \neq 0$ it is continuous but not C^1. The theorem concerns the $r > 0$ sheet of the Kerr–Schild chart; the analytic extension through the disc to $r < 0$ is not considered. Since nothing is assumed about (m, a, e), U includes the exterior, the ergoregion, the horizons themselves and the region inside the inner horizon. For $a \neq 0$ and either $e \neq 0$ or $m < 0$, the region inside the inner horizon contains closed timelike curves, which the pointwise statement does not see.

Orientation and signs. The chart is the ingoing one: for $a = e = 0$ it is Schwarzschild in ingoing Eddington–Finkelstein form, $g = \eta + (2m/r)(dt + dr)^2$, regular across the future horizon. The angular momentum is $J = +ma\,\hat{z}$; replacing a by $-a$ amounts to the reflection $y \mapsto -y$. With ∂_t future-directed and the usual identification $E_i = F_{i0}$, the electric field is $\vec{E} = \nabla H \approx -e\,\vec{x}/r^3$ at large distance, so the physical charge is $-e$; the magnetic moment is $-ea\,\hat{z}$, giving gyromagnetic ratio 2. Since the metric depends on e only through e^2 and the equations are invariant under $F \mapsto -F$, we simply call e the charge parameter. Charges are in Gaussian geometrized units: $\Delta = r^2 - 2mr + a^2 + e^2$ in Boyer–Lindquist form, and e^2 corresponds to $GQ^2/(4\pi\varepsilon_0 c^4)$ in SI units. Under the standard transformation to Boyer–Lindquist coordinates the metric becomes the textbook Kerr–Newman metric and $A \simeq (er/\Sigma)(dt - a\sin^2\theta\, d\phi)$; we verified this symbolically.




Figure 1: Dependency structure of the formal proof. An arrow $A \to B$ means that the proof of B uses A. Every node is a theorem or definition block checked on the platform. The intermediate theorems `KNE.thm_*` exist only so that each file checks within the time limit (Section 9).

2.3 The Lean statement

```
theorem kerrNewman_einsteinMaxwell (m a e : ℝ) (y : E4) (hy : 0 < ksRadius a y) :
    EinsteinMaxwellAt (kerrNewmanMetric m a e) (kerrNewmanField a e) y
```

The three conjuncts were posted as separate platform theorems, and the milestone follows from them in one line:

```
theorem solution (m a e : ℝ) (y : E4) (hy : 0 < ksRadius a y) :
    EinsteinMaxwellAt (kerrNewmanMetric m a e) (kerrNewmanField a e) y :=
  ⟨kerrNewman_einstein m a e y hy, kerrNewman_maxwell m a e y hy,
   kerrNewman_field_closed a e y hy⟩
```

3 Overview of the proof

Figure 1 shows how the proof is organized.

- The Bianchi identity (iii) holds for the exterior derivative of any C^2 potential.
- The Maxwell equation (ii) is reduced to its flat-space counterpart by Kerr–Schild algebra (Section 8.2). The flat equation is then proved from two explicit identities for the potential (Section 8.3).
- The Einstein equation (i) is the computational core. It is split into eight intermediate platform theorems, which rest on fifteen definition blocks containing closed forms and certified lemmas. A light final file assembles them.

4 Kerr–Schild algebra in atoms

4.1 Atoms and relations

Fix $p \in U$. We call the following five real numbers the *atoms* at p:

$$r, \qquad k_1,\ k_2,\ k_3 \quad \text{(the spatial components of } k\text{)}, \qquad W := \frac{1}{r^2 + a^2 k_3^2}.$$

Since $k_3 = z/r = \cos\theta$, we have $W = 1/\Sigma$, the reciprocal of the Boyer–Lindquist function.

At every point of U:

Lemma 4.1 (`ks_rel`, `ks_W`, `ks_x1`–`ks_x3`, `profile_eq`, `pot_eq`). *(a) $k_1^2 + k_2^2 + k_3^2 = 1$;*

(b) $W\,(r^2 + a^2 k_3^2) = 1$;

(c) $x = rk_1 - ak_2$, $y = ak_1 + rk_2$ and $z = rk_3$;

(d) $f = (2mr - e^2)\,W$ and $H := er^3/(r^4 + a^2 z^2) = e\,r\,W$, so that $A = H\,k$.

Proof sketch. We have $(rx+ay)^2+(ry-ax)^2 = (r^2+a^2)(x^2+y^2)$. Hence $k_1^2+k_2^2 = (x^2+y^2)/(r^2+a^2)$, and (a) is the first form of (2). Part (c) inverts the linear relations defining k_1, k_2. Part (d) uses $r^4 + a^2z^2 = r^2(r^2 + a^2k_3^2)$. □

4.2 Derivatives of the atoms

All atoms are independent of t. Implicit differentiation of (2) gives $\partial_x r = xr/(2r^2 - \rho^2 + a^2)$. By (2), $2r^2 - \rho^2 + a^2 = r^2 + a^2k_3^2 = 1/W$. Together with Lemma 4.1(c) this yields

$$\partial_1 r = Wr\,(rk_1 - ak_2), \qquad \partial_2 r = Wr\,(ak_1 + rk_2), \qquad \partial_3 r = Wk_3\,(r^2 + a^2). \tag{5}$$

The derivatives of k_1, k_2, k_3 and W are likewise polynomials in the atoms (`dKt`, `dWt`). For instance, $\partial_1 k_1 = Wr(k_2^2 + k_3^2)$ and $\partial_3 k_3 = Wr(1 - k_3^2)$. For W we have $\partial_j W = -W^2\,(2r\,\partial_j r + 2a^2 k_3\,\partial_j k_3)$.

The algebraic content of these formulas is the following. Let

$$\mathcal{P} = \mathbb{Q}[a, m, e][r, k_1, k_2, k_3, W], \qquad p_1 = k_1^2+k_2^2+k_3^2-1, \qquad p_2 = W(r^2+a^2k_3^2)-1, \qquad I = (p_1, p_2).$$

Formulas (5) and their analogues define derivations D_1, D_2, D_3 on $\mathcal{P}$ that preserve I. This was checked symbolically; it is not needed in the formal proof. So $\mathcal{P}/I$ is a differential ring, and evaluation at the atoms of a point $p \in U$ intertwines D_j with ∂_j. The ring $\mathcal{P}/I$ is the localization of $\mathbb{Q}[a, m, e][r, k_1, k_2, k_3]/(p_1)$ at $\Sigma/r^2 = r^2 + a^2k_3^2$; the extra variable W is the Rabinowitsch device for inverting it.

4.3 Kerr–Schild identities

Let $k^\sharp := \eta k = (-1, k_1, k_2, k_3)$ be the η-raised vector. On U we have $\eta^{\mu\nu}k_\mu k_\nu = 0$, $\det g = -1$ and

$$g^{-1} = \eta - f\,k^\sharp\,k^{\sharp\top}.$$

Proof. Nullity is Lemma 4.1(a). The determinant and the inverse are then checked by direct matrix computation; see [19, 46] for the general Kerr–Schild statements. □

Hence every component of g, g^{-1} and ∂g is (the evaluation of) an explicit element of $\mathcal{P}$. Since $\mathcal{P}$ is closed under the D_j, so is every component of Γ, $\partial\Gamma$, the Ricci tensor, F, $F^{\alpha\beta}$ and $S_{\mu\nu}$. No division is ever needed, because the only denominator, $r^2 + a^2k_3^2$, has been absorbed into W. This is what makes the computation tractable in a proof assistant.

5 Certified polynomial identities

5.1 Ideal membership and Gröbner cofactors

Every algebraic step of the proof has the following form. Two explicit polynomials $L, R \in \mathcal{P}$ are given, for example the Christoffel formula applied to the closed forms of g^{-1} and ∂g on one side, and a stored closed form on the other. We must show $L = R$ at all real points satisfying $p_1 = p_2 = 0$. It suffices to find $c_1, c_2 \in \mathcal{P}$ (or in its fraction field) with

$$L - R = c_1\, p_1 + c_2\, p_2. \tag{6}$$

Lemma 5.1. *Order the variables lexicographically by $W > k_1 > r > k_2 > k_3 > a > m > e$. Then $\{p_1, p_2\}$ is a Gröbner basis of I.*

Proof. The leading monomials are $\mathrm{LM}(p_1) = k_1^2$ and $\mathrm{LM}(p_2) = W r^2$, which are coprime. By Buchberger's first criterion [4, 5, 11], the S-polynomial of p_1 and p_2 reduces to zero. □

Consequently, multivariate division of $L - R$ by (p_1, p_2) returns quotients c_1, c_2 and a remainder. The remainder is zero exactly when $L - R \in I$, and in that case the quotients satisfy (6). We compute the division with SymPy's `reduced` [33]. The generator stops with an error if the remainder is nonzero.

5.2 Replay in Lean

In Lean the relations are hypotheses `hrel` : $K_1^2 + K_2^2 + K_3^2 = 1$ and `hW` : $W\,(R^2 + a^2 K_3^2) = 1$ (Lean names `R`, `K1`, …, `W` for the atoms). Each identity is closed by a single call such as

```
theorem trGam_0 (a m e R K1 K2 K3 W : ℝ) (hrel : K1 ^ 2 + K2 ^ 2 + K3 ^ 2 = 1)
    (hW : W * (R ^ 2 + a ^ 2 * K3 ^ 2) = 1) :
    Σ ρ, Gam a m e R K1 K2 K3 W ρ ρ 0 = 0 := by
  simp only [Gam, G000, G101, G202, G303, Matrix.cons_val, Fin.sum_univ_four]
  linear_combination (-2 * R^3 * W^3 * m^2 + 3 * R^2 * W^3 * m * e^2
      + 2 * R * K3^2 * W^3 * a^2 * m^2 - R * W^3 * e^4
      - K3^2 * W^3 * a^2 * m * e^2) * hrel + (0) * hW
```

(The exponent annotations `(n : ℕ)` of the source have been dropped for readability.) The tactic `linear_combination` [15] subtracts the given combination of the hypotheses from the goal. It then closes the resulting equation with `ring1`, Mathlib's normalizer for commutative rings [7, 17], which produces a proof term checked by the kernel. Nothing the computer algebra system computed is trusted. A wrong cofactor makes `ring1` fail and cannot produce a false theorem. This is the "skeptical" combination of a proof assistant with an untrusted oracle [21]. Mathlib's `polyrith` tactic [3] applied the same idea one goal at a time: it sent the goal to SageMath through a web service and printed the resulting `linear_combination` call. In the Mathlib version we use it is disabled because that service was shut down, which is one reason we compute the cofactors offline. Our generators apply it offline and in bulk: the Einstein part contains 166 calls of `linear_combination`.

5.3 Why the extra atom W

An earlier version kept $1/(r^2 + a^2 k_3^2)$ as a denominator. It closed identities with `field_simp` followed by `ring`. On the larger expressions this was far too slow, because clearing denominators multiplies terms of high degree. Making W an independent variable linked to r and k_3 by p_2 turns every

rational identity into a polynomial one. The cost is a second relation, which the Gröbner basis handles at no extra effort. A few base lemmas still use `field_simp` on small expressions, namely the derivative of the radius and the identities in Lemma 4.1.

Each closed form stored in a definition block is chosen, among a few equivalent representatives modulo p_1, to have the fewest monomials (`best` in the generator). It is written in a canonical expanded form. The Base block holds 120 such definitions: 40 Christoffel symbols with up to 23 monomials each, 30 divergence terms $\partial_\rho \Gamma^\rho_{\nu\sigma}$ with up to 101 monomials, 10 Ricci components with up to 13 monomials, the field components and the invariant. There are 162 certificates of the form $c_1 \cdot \texttt{hrel} + c_2 \cdot \texttt{hW}$ in the Einstein part. Twenty-one of them, in the base lemmas, have rational cofactors and are normalized by `field_simp`; 23 are trivial (both cofactors zero, so `ring` alone would do); the remaining 118 have polynomial cofactors with a median of 32 monomials. The largest certificate, for the $(1, 2)$ component of the Ricci tensor, has 356 monomials, total degree 24, and occupies 28 kB of source.

6 Derivatives: a small calculus of coordinate partials

The definitions use $\partial_k f(p) = Df(p)(e_k)$. Rewriting with Mathlib's `fderiv` lemmas on large expressions generates differentiability side goals. It also manipulates continuous linear maps, which is slow. We instead use a one-line predicate whose closure rules compute partial derivatives syntactically.

```
/-- ‘f‘ is differentiable at ‘z‘ with coordinate partials ‘d k‘. -/
structure HasCP (f : E4 → ℝ) (z : E4) (d : Fin 4 → ℝ) : Prop where
  diff : DifferentiableAt ℝ f z
  line : ∀ k, HasDerivAt (fun t : ℝ => f (z + t • coordBasis k)) (d k) 0

theorem HasCP.eq (h : HasCP f z df) (k : Fin 4) : coordPartial f k z = df k
```

The closure rules (`HasCP.cst`, `add`, `sub`, `neg`, `mul`, `div`, `pow`) follow from the one-variable rules for `HasDerivAt` along coordinate lines. For example,

$$\texttt{HasCP.mul}: \quad \mathsf{HasCP}\, f\, z\, d_f \;\wedge\; \mathsf{HasCP}\, g\, z\, d_g \;\Longrightarrow\; \mathsf{HasCP}\,(fg)\, z\,(k \mapsto d_f(k)\, g(z) + f(z)\, d_g(k)).$$

A further rule, `congr_d`, replaces the derivative by any pointwise-equal one.

Base cases. `hasCP_R` differentiates the radius (1) along coordinate lines. It needs $r > 0$ and positivity of the inner square root, and it is matched to (5) by a certificate. `hasCP_K1`–`hasCP_K3`, `hasCP_W`, `hasCP_profile` and `hasCP_H` follow, all with derivatives in atom form.

Generated derivations. Each Christoffel closed form is a polynomial P in the atoms. To obtain ∂_ρ of $p \mapsto P(\text{atoms}(p))$, the generator emits a `HasCP` proof term built from the closure rules. The term follows the same syntax tree that was used to print P: a balanced binary sum of monomials. Lean therefore accepts the term by unfolding definitions, with no rewriting. The derivative it produces is the chain-rule expansion, which `congr_d` together with one certificate identifies with the stored closed form of the divergence term. The Lean statement is `dG112`, for instance, which gives both differentiability and the value of ∂_1 of the component Γ^1_{12}. The thirty-four generated derivations contain 5,286 applications of the closure rules in total; the largest single term is about 4 kB.

For any $\Phi : \mathbb{R}^4 \to \mathbb{R}$, the function $p \mapsto \Phi(r(p), k_1(p), k_2(p), k_3(p))$ has $\partial_0 = 0$ everywhere.

Proof. The function is invariant under $p \mapsto p + te_0$. If it is differentiable at p, its derivative along e_0 is that of a constant. If not, the Fréchet derivative is 0 by convention. □

From local identities to derivatives. The proof shows that the platform's Christoffel symbols equal the closed form `Gam` evaluated at the atoms, at every point of U (`thm_christoffel`). Because U is open, the two functions agree near any $p \in U$, so their derivatives at p agree (`Filter.EventuallyEq.fderiv_eq`). This is how derivatives of the platform's `christoffel` reduce to derivatives of closed forms.

7 The Einstein equation

Fix $p \in U$ and write Γ, $D^\rho_{\nu\sigma} := \partial_\rho \Gamma^\rho_{\nu\sigma}$ (no sum) and so on for the closed forms evaluated at the atoms of p.

Christoffel symbols. By Lemma 4.2, g^{-1} is polynomial in the atoms. With the `HasCP` derivatives of $g = \eta + f\, k \otimes k$ (`cp_metric`), the Christoffel formula becomes a polynomial expression. Forty certificates, one for each Γ^i_{jk} with $j \le k$, identify it with the stored closed forms (`thm_christoffel`).

Ricci tensor. Unfolding the definitions,

$$R_{\sigma\nu} = \sum_\rho \partial_\rho \Gamma^\rho_{\nu\sigma} - \partial_\nu\Big(\sum_\rho \Gamma^\rho_{\rho\sigma}\Big) + \sum_{\rho,\lambda} \Gamma^\rho_{\rho\lambda}\Gamma^\lambda_{\nu\sigma} - \sum_{\rho,\lambda} \Gamma^\rho_{\nu\lambda}\Gamma^\lambda_{\rho\sigma}. \tag{7}$$

The contracted symbol $\sum_\rho \Gamma^\rho_{\rho\sigma} = \frac{1}{2}\partial_\sigma \log|\det g|$ vanishes on U because $\det g = -1$. In the formal proof this is not derived from the determinant. It is checked directly on the closed forms by four certificates (`trGam`). Since the trace vanishes on the open set U and each summand is differentiable (`diff_tr`), the second term of (7) is zero, and so is the third. The first term is given by the divergence closed forms. For $\rho = 0$ it vanishes by Lemma 6.1. For $\rho = 1, 2, 3$ it comes from the thirty generated derivations (`thm_div_one`–`thm_div_three`). The remaining expression is identified with a closed-form Ricci tensor `RicA` by ten certificates, one for each component with $\sigma \le \nu$ (`thm_ricci_a`–`thm_ricci_c`). The other six components follow by symmetry.

Scalar curvature. One certificate shows $\sum_{\sigma,\nu} g^{\sigma\nu} R_{\sigma\nu} = 0$ (`scal_zero`). This reflects the tracelessness of the Maxwell stress tensor in four dimensions. Hence $G_{\mu\nu} = R_{\mu\nu}$ at p.

The source. Using the `HasCP` rules for $A = Hk$, the field $F_{\mu\nu} = \partial_\mu(Hk_\nu) - \partial_\nu(Hk_\mu)$ is put in closed form (`field_closed` in the final file). Further certificates give $F^{\alpha\beta}$, the invariant $F_{\alpha\beta}F^{\alpha\beta}$ and finally $S_{\mu\nu}$ in closed form. The last step identifies $S_{\mu\nu}$ with the same polynomial `RicA` (`thm_maxwell`).

Conclusion. The final file chains these facts:

```
theorem einstein_eq (m a e : ℝ) (y : E4) (hy : 0 < ksRadius a y) (μ ν : Fin 4) :
    einsteinTensor (kerrNewmanMetric m a e) y μ ν
      = RicA a m e (ksRadius a y) (ksNull a y 1) (ksNull a y 2) (ksNull a y 3) (Wf a y)
    ↪ μ ν
theorem maxwell_eq (m a e : ℝ) (y : E4) (hy : 0 < ksRadius a y) (μ ν : Fin 4) :
```

```
maxwellSource (kerrNewmanMetric m a e) (kerrNewmanField a e) y μ ν
  = RicA a m e (ksRadius a y) (ksNull a y 1) (ksNull a y 2) (ksNull a y 3) (Wf a y)
↪  μ ν
```

The Einstein equation (i) is then a single rewrite. Before any Lean code was generated, the same symbolic pipeline verified $G_{\mu\nu} = S_{\mu\nu}$ and $R = 0$ in SymPy, and spot checks were made at random numerical points. These checks guided development but play no role in the formal proof.

8 The Maxwell equations

8.1 Closedness

For any potential A that is C^2 near p, the three terms of $\partial_{[\lambda} F_{\mu\nu]}$ cancel pairwise by symmetry of second derivatives. Mathlib provides this symmetry as `ContDiffAt.isSymmSndFDerivAt`. Smoothness of A on U is discharged by Mathlib's `fun_prop`, given that $r > 0$, that $r^2 + a^2 \neq 0$ and $r^4 + a^2 z^2 \neq 0$, and that the discriminant in (1) is positive. The whole proof (`kerrNewman_field_closed`) is 88 lines.

8.2 From curved to flat space

On U, $\sqrt{|\det g|}\, F^{\mu\nu} = \eta^{\mu\alpha}\eta^{\nu\beta} F_{\alpha\beta}$.

Proof. By Lemma 4.2, $\sqrt{|\det g|} = 1$ and $g^{-1} = P := \eta - f\, k^\sharp k^{\sharp\top}$. Write $F = d(Hk)$.

(i) Contracting $F_{\mu\nu} = \partial_\mu H\, k_\nu - \partial_\nu H\, k_\mu + H(\partial_\mu k_\nu - \partial_\nu k_\mu)$ with $k^{\sharp\nu}$ uses three facts. First, $k_\nu k^{\sharp\nu} = 0$ (nullity). Second, $k^{\sharp\nu}\partial_\mu k_\nu = \frac{1}{2}\partial_\mu(\eta^{\alpha\beta} k_\alpha k_\beta) = 0$. Third, $k^{\sharp\nu}\partial_\nu k_\mu = 0$, i.e. $k^\sharp$ is geodesic for η; this is checked with (5). Together they give the alignment

$$F_{\mu\nu}\, k^{\sharp\nu} = \lambda\, k_\mu, \qquad \lambda = -k^{\sharp\nu}\partial_\nu H.$$

(ii) By antisymmetry, $k^{\sharp\top} F = -\lambda\, k^\top$. Using $\eta k = k^\sharp$, we get $\eta F k^\sharp = \lambda k^\sharp$, $k^{\sharp\top} F \eta = -\lambda k^{\sharp\top}$ and $k^{\sharp\top} F k^\sharp = 0$. Hence

$$\begin{aligned} PFP^\top &= \eta F \eta - f\, k^\sharp (k^{\sharp\top} F \eta) - f\, (\eta F k^\sharp) k^{\sharp\top} + f^2\, k^\sharp (k^{\sharp\top} F k^\sharp) k^{\sharp\top} \\ &= \eta F \eta + f\lambda\, k^\sharp k^{\sharp\top} - f\lambda\, k^\sharp k^{\sharp\top} = \eta F \eta. \end{aligned}$$

□

Since the identity holds on the open set U, the divergences of both sides agree at every $p \in U$. The curved Maxwell equation (ii) is thus equivalent to the flat one, $\partial_\mu(\eta^{\mu\alpha}\eta^{\nu\beta} F_{\alpha\beta}) = 0$. This proof (`kerrNewman_maxwell`, 460 lines) does not use the atom machinery. Proposition 8.1 is closely related to the Kerr–Schild double copy [35], in which the gauge field ϕk built from a stationary Kerr–Schild metric solves the flat Maxwell equations.

8.3 The flat Maxwell equation

All fields are t-independent. Write $A = (H, \vec{A})$ with $\vec{A} = H\,\vec{k}$. The $\nu = 0$ component of the flat equation is $-\Delta H = 0$. The $\nu = i$ components are $\sum_j \partial_j(\partial_j A_i - \partial_i A_j) = -(\nabla \times \nabla \times \vec{A})_i = 0$. We prove two identities on U:

$$\Delta H = 0, \qquad \nabla \times \vec{A} = \nabla\psi, \qquad \psi := \frac{e\, a\, z\, r}{r^4 + a^2 z^2}. \tag{8}$$

The second implies $\nabla \times \nabla \times \vec{A} = \nabla \times \nabla\psi = 0$, by symmetry of the second derivatives of the smooth function ψ. Both identities in (8) are verified componentwise in Lean. Each component is an explicit derivative computation (along coordinate lines, as in Section 6) followed by `field_simp`, `ring` and `linear_combination` with the quartic relation (2). The proof (`kerrNewman_flat_maxwell`) is 720 lines.

Interpretation. The pair (H, ψ) is the real and imaginary part of Appell's potential [2], the Coulomb potential of a charge placed at the imaginary point $(0, 0, ia)$:

$$H + i\psi = \frac{e\,r}{r^2 - iaz} = \frac{e}{r - iaz/r} = \frac{e}{\sqrt{x^2 + y^2 + (z - ia)^2}}.$$

The last equality uses $(r - iaz/r)^2 = x^2 + y^2 + (z - ia)^2$, which follows from (2), and picks the root with positive real part r. This is the classical picture of the Kerr–Newman field as the field of a complex-shifted point charge [1, 29, 36]. In this picture $\Delta H = 0$ is the harmonicity of the real part of a harmonic function. The formal proof does not use complex numbers; it verifies (8) directly.

9 Engineering a large proof under a time budget

9.1 Platform constraints

Prove2Me checks each submission on its server within 300 seconds. This budget includes building any platform modules the submission imports that are not already cached. Submissions may not define new syntax or tactics (no `macro` or `syntax` commands). The platform distinguishes *definition blocks*, which are published modules that other files may import and which may contain lemmas, from *theorems*. A proof may import a theorem's statement and use it as an axiom-free reduction that the platform tracks. That theorem's own proof is checked separately.

Our first complete proof of the Einstein equation was a single file of about 2,650 lines. It needed more than 13 GB of memory and more than 300 seconds on our machine, and the platform ran out of memory while rebuilding its dependencies. The final development (Table 1) splits the work along the natural stages of the computation.

- Fifteen definition blocks hold the closed forms, the `HasCP` base lemmas and the heavy certified lemmas.
- Eight intermediate theorems restate the needed facts over the closed forms. Their proofs are a few lines that `exact` lemmas from the blocks.
- A final file of 251 lines imports only the two light blocks and the eight theorem statements, and assembles the result.

The heaviest block now checks in 76 seconds, and the whole Einstein part in about ten minutes of elaboration time. Generating the Lean source takes about four and a half minutes of SymPy time, the bulk of it in the symbolic curvature computation that doubles as an independent check. The generated fragments are 94% of the Einstein part by volume; the rest is hand-written glue.

9.2 Lessons

Several low-level choices mattered as much as the overall design.

File	Content	Lines	kB	Lemmas	Certif.	Check (s)
Definition blocks						
Base	atoms, `HasCP` calculus, 120 closed forms	798	245	51	–	28
Atoms	`HasCP` for r, k_i, W, f, H; g, g^{-1}, Γ	256	20	12	28	26
Christoffel	40 identities Γ^i_{jk} = closed form	310	96	46	40	74
Div1	∂_0 differentiability; $\partial_1\Gamma^1_{0\sigma}$	142	36	8	4	20
Div2	$\partial_1\Gamma^1_{\nu\sigma}$, $1 \le \nu \le \sigma$	150	82	6	6	73
Div3	$\partial_2\Gamma^2_{\nu\sigma}$	238	85	10	10	76
Div4	$\partial_3\Gamma^3_{\nu\sigma}$	238	63	10	10	49
Ricci1–6	10 identities $R_{\sigma\nu}$ = closed form	168	232	10	10	205
MaxwellF	F, $F^{\alpha\beta}$, $F_{\alpha\beta}F^{\alpha\beta}$ in closed form	223	19	35	33	18
MaxwellT	16 identities $S_{\mu\nu}$ = closed form	118	70	17	16	41
Theorems						
`KNE.thm_*` (8)	restatements over the blocks	193	10	8	–	< 1
`kerrNewman_einstein`	assembly (Section 7)	251	17	28	5	3
`kerrNewman_field_closed`	$dF = 0$	88	4	7	–	1
`kerrNewman_flat_maxwell`	flat Maxwell (Section 8.3)	720	35	79	–	8
`kerrNewman_maxwell`	reduction to flat (Section 8.2)	460	23	48	–	16
`kerrNewman_einsteinMaxwell`	milestone	15	1	1	–	< 1
Total		4,368	1,038	376	162	639

Table 1: The files of the formal proof. "Lemmas" counts top-level theorems and lemmas; "Certif." counts `linear_combination` certificates of the form $c_1 \cdot$ `hrel` $+ c_2 \cdot$ `hW` (the Ricci and Maxwell rows also contain a few hand-written ones). "Check" is the wall-clock time to re-elaborate the file against compiled imports, net of the 7.6 s needed to load Mathlib, minimum of two runs on a 4-core 2.1 GHz machine with 15 GB of memory. Peak memory was 6.4–8.4 GB per file, almost all of it the memory-mapped Mathlib library.

- *Annotate exponents.* Writing `R ^ (4 : ℕ)` instead of `R ^ 4` in generated polynomials avoided slow elaboration of the many `^` occurrences under Lean's default-instance mechanism.
- *One definition per closed form.* A single table holding all forty Christoffel symbols exceeded the heartbeat limit during elaboration; one `def` per entry did not.
- *Print and prove from the same tree.* The `HasCP` terms are generated from the expression tree used to print the definition they differentiate. The two then agree by definitional unfolding, and no rewriting is needed. Mismatched association of sums was the main source of early type errors.
- *Exploit symmetry.* Only components with $j \le k$ of Γ^i_{jk} and $\sigma \le \nu$ of $R_{\sigma\nu}$ are certified. The symmetry lemmas themselves are proved by `rfl` or by unfolding `Matrix.cons_val` with `simp only`.
- *Prefer polynomial identities.* As explained in Section 5, replacing `field_simp` by polynomial certificates was decisive for speed.

10 Trusted base and validation

What is checked. For the final theorem and for the milestone, `#print axioms` reports

```
'NoHair.kerrNewman_einsteinMaxwell' depends on axioms: [propext, Classical.choice, Quot.sound]
```

No `sorry`, `admit`, `native_decide` or `axiom` occurs anywhere in the 4,368 lines of proof, and no heartbeat limit is raised. Every file was checked by the Prove2Me server with the platform's toolchain. On the platform each theorem is checked against `sorry`-stubs of the theorems it imports, so the platform's verdict is compositional. We also rebuilt the whole chain locally in a single Lean environment, with every stub replaced by its accepted proof under the platform name, and ran `#print axioms` on the milestone and on each of its three conjuncts; all four report exactly the three axioms above. The scripts for this check are in the repository.

What must be trusted. To believe Theorem 1.1 one must trust the Lean kernel and the Mathlib definitions involved: `fderiv`, matrix inverse and determinant, `Real.sqrt`. One must also accept that the platform definitions of Sections 2.1 and 2.2 express the intended notions. These are about 150 lines, reproduced in Appendix A. We compared them with the standard references (Section 2.1). The generators, SymPy and the AI agent that wrote the proofs need not be trusted. Neither does the reader need to inspect the roughly one megabyte of generated proof text.

Limitations. The statement is coordinate-based and pointwise. It does not use Mathlib's manifold library, which does not yet provide pseudo-Riemannian curvature. It covers the $r > 0$ sheet of the Kerr–Schild chart. It concerns only the *existence* side of the no-hair story, namely that Kerr–Newman is a solution. The uniqueness theorems, which are the mission's real goal, are far beyond the present work. Another milestone, `NoHair.kerrNewman_isStationaryBlackHole`, which asserts that the Kerr–Newman spacetime satisfies the global hypotheses of the uniqueness theorem, is not addressed here. Finally, the generated proof is optimized for checkability, not readability. The human-readable content is the method described above together with the generators.

11 Related work

Computer algebra in general relativity. Verifying exact solutions is a classical application of computer algebra; see MacCallum's review [30]. It is supported by dedicated systems such as xAct [31], Cadabra [38] and SageManifolds [16], as well as general-purpose systems. The catalogue of Stephani et al. [46] relies on such checks. These tools produce results, not proofs. Our pipeline uses SymPy in exactly this role, but only to find certificates that Lean then checks.

Certificates from untrusted oracles. Combining a proof assistant with an external algebra system that supplies checkable witnesses goes back at least to Harrison and Théry [21]. Later instances include the use of Maple from Coq [14] and certified computer algebra on top of a prover [25]. Mathlib's `polyrith` [3] found `linear_combination` coefficients with an external system. Gröbner-basis certificates have been used for ring identities in Isabelle [9] and HOL Light [20], and Buchberger's algorithm itself has been verified in Coq [48]. Coq's `nsatz` checks certificates produced by external Gröbner computations [40]. Lean has been connected to Mathematica [28], and recent work develops Gröbner-based tactics and Gröbner theory in Lean 4 [18, 44]. Our use of Gröbner-basis cofactors as certificates is standard. What is specific here is the reduction of an entire curvature computation to ideal membership in a small differential ring of atoms.

Formalized physics and relativity. Formal treatments of relativity have so far been mostly axiomatic or kinematic. Stannett and Németi verified results of first-order axiomatic relativity in Isabelle/HOL [45], and Schutz's axioms for Minkowski spacetime have been formalized there as well [43]. In Lean 4, the HepLean project (now Physlib) [39, 49, 50] formalizes parts of high-energy

physics, tensor index notation and special relativity. Recent work proves analytic results on a fixed black-hole background [26], taking the metric as given. We are not aware of an earlier formal proof that the Schwarzschild, Kerr or Kerr–Newman metric satisfies the Einstein equations with the curvature computed from the metric through the prover's own analysis library. Our searches of the literature and of public code repositories (October 2026) found none.

12 Conclusion

We have given a complete machine-checked proof that the Kerr–Newman metric and field solve the Einstein–Maxwell equations. The definitions are elementary and the region is the full $r > 0$ part of the Kerr–Schild chart. The approach combines a choice of atoms that makes the solution's algebra polynomial and closed under differentiation, cheap certificates from an untrusted Gröbner-basis computation, and a small differentiation calculus. It should transfer to other exact solutions whose metric components lie in a finitely generated differential algebra with a few relations. Natural next targets are Kerr–Schild metrics more generally and accelerating or NUT-charged solutions. On the conceptual side, connecting the coordinate statement to a manifold-level Lorentzian geometry library would make the result reusable in the mission's harder milestones.

Acknowledgements and use of AI tools

The Lean proofs, the SymPy generators and a first draft of this manuscript were produced by Claude Code, an AI coding agent developed by Anthropic. The agent worked under the author's direction on the Prove2Me platform. The formal statement and definitions were authored by the Prove2Me mission authors. All formal claims were checked by the Lean kernel. The author takes responsibility for the content of this paper.

Data availability

The complete development is publicly available in the repository

`https://github.com/SamenSHossain/math-solver`

in the directory `prove2me/nohair/`, which holds all Lean files, the Python generators and the definition blocks. The platform objects are listed in Appendix B and can be inspected on `https://prove2.me`. The directory `paper/kerr-newman/` contains the source of this paper, the scripts that reproduce the statistics and the single-environment axiom check, and the audit scripts for the conventions of Section 2.1. The Lean toolchain was version 4.33.1 with Mathlib commit `0df444a3`.

References

[1] Tim Adamo and E. T. Newman. The Kerr–Newman metric: A review. *Scholarpedia*, 9(10): 31791, 2014. doi: 10.4249/scholarpedia.31791.

[2] Paul Appell. Quelques remarques sur la théorie des potentiels multiformes. *Mathematische Annalen*, 30:155–156, 1887.

[3] Dhruv Bhatia, Eric Wieser, Mario Carneiro, and Thomas Zhu. Mathlib.Tactic.Polyrith: the `polyrith` tactic. Mathlib source file, `https://github.com/leanprover-community/mathlib4/blob/master/Mathlib/Tactic/Polyrith.lean`, 2022.

[4] Bruno Buchberger. *Ein Algorithmus zum Auffinden der Basiselemente des Restklassenringes nach einem nulldimensionalen Polynomideal*. PhD thesis, Universität Innsbruck, 1965.

[5] Bruno Buchberger. A criterion for detecting unnecessary reductions in the construction of Gröbner-bases. In Edward W. Ng, editor, *Symbolic and Algebraic Computation (EUROSAM '79)*, volume 72 of *Lecture Notes in Computer Science*, pages 3–21. Springer, 1979. doi: 10.1007/3-540-09519-5_52.

[6] Gary L. Bunting. *Proof of the Uniqueness Conjecture for Black Holes*. PhD thesis, University of New England, Armidale, N.S.W., 1983.

[7] Mario Carneiro, Aurélien Saue, and Anne Baanen. Mathlib.Tactic.Ring: the `ring` tactic. Mathlib source file, `https://github.com/leanprover-community/mathlib4/blob/master/Mathlib/Tactic/Ring/Basic.lean`, 2018.

[8] Brandon Carter. Axisymmetric black hole has only two degrees of freedom. *Physical Review Letters*, 26(6):331–333, 1971. doi: 10.1103/PhysRevLett.26.331.

[9] Amine Chaieb and Makarius Wenzel. Context aware calculation and deduction: ring equalities via Gröbner bases in Isabelle. In Manuel Kauers, Manfred Kerber, Robert Miner, and Wolfgang Windsteiger, editors, *Towards Mechanized Mathematical Assistants (Calculemus 2007 / MKM 2007)*, volume 4573 of *Lecture Notes in Artificial Intelligence*. Springer, 2007. doi: 10.1007/978-3-540-73086-6_3.

[10] Piotr T. Chruściel, João Lopes Costa, and Markus Heusler. Stationary black holes: Uniqueness and beyond. *Living Reviews in Relativity*, 15:7, 2012. doi: 10.12942/lrr-2012-7.

[11] David A. Cox, John Little, and Donal O'Shea. *Ideals, Varieties, and Algorithms: An Introduction to Computational Algebraic Geometry and Commutative Algebra*. Undergraduate Texts in Mathematics. Springer, Cham, 4 edition, 2015. doi: 10.1007/978-3-319-16721-3.

[12] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In André Platzer and Geoff Sutcliffe, editors, *Automated Deduction – CADE 28*, volume 12699 of *Lecture Notes in Computer Science*, pages 625–635, Cham, 2021. Springer. doi: 10.1007/978-3-030-79876-5_37.

[13] G. C. Debney, R. P. Kerr, and A. Schild. Solutions of the Einstein and Einstein–Maxwell equations. *Journal of Mathematical Physics*, 10(10):1842–1854, 1969. doi: 10.1063/1.1664769.

[14] David Delahaye and Micaela Mayero. Dealing with algebraic expressions over a field in Coq using Maple. *Journal of Symbolic Computation*, 39(5):569–592, 2005. doi: 10.1016/j.jsc.2004.12.004.

[15] Abby J. Goldberg, Mario Carneiro, and Heather Macbeth. Mathlib.Tactic.LinearCombination: the `linear_combination` tactic. Mathlib source file, `https://github.com/leanprover-community/mathlib4/blob/master/Mathlib/Tactic/LinearCombination.lean`, 2022.

[16] Eric Gourgoulhon, Michał Bejger, and Marco Mancini. Tensor calculus with open-source software: the SageManifolds project. *Journal of Physics: Conference Series*, 600:012002, 2015.

[17] Benjamin Grégoire and Assia Mahboubi. Proving equalities in a commutative ring done right in Coq. In Joe Hurd and Tom Melham, editors, *Theorem Proving in Higher Order Logics*

(TPHOLs 2005), volume 3603 of *Lecture Notes in Computer Science*, pages 98–113. Springer, 2005. doi: 10.1007/11541868_7.

[18] Junyu Guo, Hao Shen, Junqi Liu, and Lihong Zhi. Formalizing Gröbner basis theory in Lean, 2026.

[19] Metin Gürses and Feza Gürsey. Lorentz covariant treatment of the Kerr–Schild geometry. *Journal of Mathematical Physics*, 16(12):2385–2390, 1975. doi: 10.1063/1.522480.

[20] John Harrison. Automating elementary number-theoretic proofs using Gröbner bases. In Frank Pfenning, editor, *Automated Deduction – CADE-21*, volume 4603 of *Lecture Notes in Computer Science*, pages 51–66. Springer, 2007.

[21] John Harrison and Laurent Théry. A skeptic's approach to combining HOL and Maple. *Journal of Automated Reasoning*, 21(3):279–294, 1998. doi: 10.1023/A:1006023127567.

[22] Stephen W. Hawking. Black holes in general relativity. *Communications in Mathematical Physics*, 25:152–166, 1972. doi: 10.1007/BF01877517.

[23] Werner Israel. Event horizons in static vacuum space-times. *Physical Review*, 164(5):1776–1779, 1967. doi: 10.1103/PhysRev.164.1776.

[24] Werner Israel. Event horizons in static electrovac space-times. *Communications in Mathematical Physics*, 8(3):245–260, 1968. doi: 10.1007/BF01645859.

[25] Cezary Kaliszyk and Freek Wiedijk. Certified computer algebra on top of an interactive theorem prover. In *Towards Mechanized Mathematical Assistants (Calculemus 2007 / MKM 2007)*, volume 4573 of *Lecture Notes in Computer Science*, pages 94–105. Springer, 2007. doi: 10.1007/978-3-540-73086-6_8.

[26] Christoph Kehle. Charged scalar mode stability on Reissner–Nordström (Lean 4 formalization). `https://github.com/christoph-ke/rn-charged-scalar-mode-stability`, 2026.

[27] Roy P. Kerr. Gravitational field of a spinning mass as an example of algebraically special metrics. *Physical Review Letters*, 11(5):237–238, 1963. doi: 10.1103/PhysRevLett.11.237.

[28] Robert Y. Lewis and Minchao Wu. A bi-directional extensible interface between Lean and Mathematica. *Journal of Automated Reasoning*, 66(2):215–238, 2022. doi: 10.1007/s10817-021-09611-1.

[29] Donald Lynden-Bell. A magic electromagnetic field. In Michael J. Thompson and Jørgen Christensen-Dalsgaard, editors, *Stellar Astrophysical Fluid Dynamics*, pages 369–375. Cambridge University Press, Cambridge, 2003.

[30] Malcolm A. H. MacCallum. Computer algebra in gravity research. *Living Reviews in Relativity*, 21:6, 2018. doi: 10.1007/s41114-018-0015-6.

[31] José M. Martín-García. xPerm: fast index canonicalization for tensor computer algebra. *Computer Physics Communications*, 179(8):597–603, 2008. doi: 10.1016/j.cpc.2008.05.009.

[32] Pawel O. Mazur. Proof of uniqueness of the Kerr–Newman black hole solution. *Journal of Physics A: Mathematical and General*, 15(10):3173–3180, 1982. doi: 10.1088/0305-4470/15/10/021.

[33] Aaron Meurer, Christopher P. Smith, Mateusz Paprocki, Ondřej Čertík, Sergey B. Kirpichev, Matthew Rocklin, Amit Kumar, Sergiu Ivanov, Jason K. Moore, Sartaj Singh, Thilina Rathnayake, Sean Vig, Brian E. Granger, Richard P. Muller, Francesco Bonazzi, Harsh Gupta, Shivam Vats, Fredrik Johansson, Fabian Pedregosa, Matthew J. Curry, Andy R. Terrel, Štěpán Roučka, Ashutosh Saboo, Isuru Fernando, Sumith Kulal, Robert Cimrman, and Anthony Scopatz. SymPy: symbolic computing in Python. *PeerJ Computer Science*, 3:e103, 2017. doi: 10.7717/peerj-cs.103.

[34] Charles W. Misner, Kip S. Thorne, and John Archibald Wheeler. *Gravitation*. W. H. Freeman, San Francisco, 1973. ISBN 0716703440.

[35] Ricardo Monteiro, Donal O'Connell, and Chris D. White. Black holes and the double copy. *Journal of High Energy Physics*, 2014(12):056, 2014. doi: 10.1007/JHEP12(2014)056.

[36] E. T. Newman. Maxwell's equations and complex Minkowski space. *Journal of Mathematical Physics*, 14:102–103, 1973.

[37] E. T. Newman, E. Couch, K. Chinnapared, A. Exton, A. Prakash, and R. Torrence. Metric of a rotating, charged mass. *Journal of Mathematical Physics*, 6(6):918–919, 1965. doi: 10.1063/1.1704351.

[38] Kasper Peeters. Cadabra: a field-theory motivated symbolic computer algebra system. *Computer Physics Communications*, 176:550–558, 2007. doi: 10.1016/j.cpc.2007.01.003.

[39] PhysLean/Physlib contributors. Physlib: Digitalizing physics in Lean 4 (formerly HepLean/PhysLean). `https://github.com/leanprover-community/physlib`, commit 5b36d166bcb4ef1b86826e50c8accad46d47bd71, 2026.

[40] Loïc Pottier. Connecting Gröbner bases programs with Coq to do proofs in algebra, geometry and arithmetics, 2010.

[41] Prove2Me. Prove2me: a platform for collaborative formal mathematics in Lean. `https://prove2.me`, 2026. Mission "No-Hair Conjecture: Uniqueness of Stationary Kerr–Newman Black Holes", accessed October 2026.

[42] D. C. Robinson. Uniqueness of the Kerr black hole. *Physical Review Letters*, 34(14):905–906, 1975. doi: 10.1103/PhysRevLett.34.905.

[43] Richard Schmoetten, Jake E. Palmer, and Jacques D. Fleuriot. Towards formalising Schutz' axioms for Minkowski spacetime in Isabelle/HOL, 2021.

[44] Hao Shen, Junyu Guo, Junqi Liu, and Lihong Zhi. Automated tactics for polynomial reasoning in Lean 4, 2026.

[45] Mike Stannett and István Németi. Using Isabelle/HOL to verify first-order relativity theory. *Journal of Automated Reasoning*, 52(4):361–378, 2014. doi: 10.1007/s10817-013-9292-7.

[46] Hans Stephani, Dietrich Kramer, Malcolm MacCallum, Cornelius Hoenselaers, and Eduard Herlt. *Exact Solutions of Einstein's Field Equations*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 2 edition, 2003. ISBN 9780521461368. doi: 10.1017/CBO9780511535185.

[47] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020)*, pages 367–381, New York, NY, USA, 2020. ACM. doi: 10.1145/3372885.3373824.

[48] Laurent Théry. A machine-checked implementation of Buchberger's algorithm. *Journal of Automated Reasoning*, 26(2):107–137, 2001. doi: 10.1023/A:1026518331905.

[49] Joseph Tooby-Smith. Formalization of physics index notation in Lean 4, 2024.

[50] Joseph Tooby-Smith. HepLean: Digitalising high energy physics. *Computer Physics Communications*, 308:109457, 2025. doi: 10.1016/j.cpc.2024.109457.

[51] Matt Visser. The Kerr spacetime: A brief introduction. In David L. Wiltshire, Matt Visser, and Susan M. Scott, editors, *The Kerr Spacetime: Rotating Black Holes in General Relativity*. Cambridge University Press, Cambridge, 2009. ISBN 9780521885126.

[52] Robert M. Wald. *General Relativity*. University of Chicago Press, Chicago, 1984. ISBN 0226870332.

A The platform definitions

These definitions are reproduced verbatim from the Prove2Me definition blocks `NoHair_coordinates` and `NoHair_kerrNewman`; docstrings are shortened. They are everything Theorem 1.1 depends on beyond Mathlib.

```
abbrev E4 := EuclideanSpace ℝ (Fin 4)
def coordBasis (μ : Fin 4) : E4 := EuclideanSpace.single μ 1
def minkowski : Matrix (Fin 4) (Fin 4) ℝ := Matrix.diagonal ![-1, 1, 1, 1]
def coordPartial (f : E4 → ℝ) (k : Fin 4) (y : E4) : ℝ := fderiv ℝ f y (coordBasis k)
def invMetric (G : E4 → Matrix (Fin 4) (Fin 4) ℝ) (y : E4) : Matrix (Fin 4) (Fin 4) ℝ
    ↪ :=
  (G y)⁻¹

def christoffel (G : E4 → Matrix (Fin 4) (Fin 4) ℝ) (y : E4) (i j k : Fin 4) : ℝ :=
  Σ l, (1 / 2 : ℝ) * invMetric G y i l *
    (coordPartial (fun z => G z l k) j y + coordPartial (fun z => G z l j) k y
      - coordPartial (fun z => G z j k) l y)

def riemann (G : E4 → Matrix (Fin 4) (Fin 4) ℝ) (y : E4) (ρ σ μ ν : Fin 4) : ℝ :=
  coordPartial (fun z => christoffel G z ρ ν σ) μ y
    - coordPartial (fun z => christoffel G z ρ μ σ) ν y
    + Σ l, (christoffel G y ρ μ l * christoffel G y l ν σ
      - christoffel G y ρ ν l * christoffel G y l μ σ)

def ricci (G : E4 → Matrix (Fin 4) (Fin 4) ℝ) (y : E4) (σ ν : Fin 4) : ℝ :=
  Σ ρ, riemann G y ρ σ ρ ν

def scalarCurvature (G : E4 → Matrix (Fin 4) (Fin 4) ℝ) (y : E4) : ℝ :=
  Σ σ, Σ ν, invMetric G y σ ν * ricci G y σ ν

def einsteinTensor (G : E4 → Matrix (Fin 4) (Fin 4) ℝ) (y : E4) (μ ν : Fin 4) : ℝ :=
  ricci G y μ ν - (1 / 2 : ℝ) * scalarCurvature G y * G y μ ν
```

```
def raiseBoth (G : E4 → Matrix (Fin 4) (Fin 4) ℝ) (F : E4 → Matrix (Fin 4) (Fin 4) ℝ
    ↪ )
    (y : E4) (α β : Fin 4) : ℝ :=
  Σ μ, Σ ν, invMetric G y α μ * invMetric G y β ν * F y μ ν

def maxwellSource (G : E4 → Matrix (Fin 4) (Fin 4) ℝ) (F : E4 → Matrix (Fin 4) (Fin
    ↪ 4) ℝ)
    (y : E4) (μ ν : Fin 4) : ℝ :=
  2 * ((Σ α, Σ β, F y μ α * invMetric G y α β * F y ν β)
    - (1 / 4 : ℝ) * G y μ ν * Σ α, Σ β, F y α β * raiseBoth G F y α β)

def EinsteinMaxwellAt (G : E4 → Matrix (Fin 4) (Fin 4) ℝ) (F : E4 → Matrix (Fin 4) (
    ↪ Fin 4) ℝ)
    (y : E4) : Prop :=
  (∀ μ ν, einsteinTensor G y μ ν = maxwellSource G F y μ ν) ∧
  (∀ ν, Σ μ, coordPartial
      (fun z => Real.sqrt |(G z).det| * raiseBoth G F z μ ν) μ y = 0) ∧
  (∀ l μ ν, coordPartial (fun z => F z μ ν) l y + coordPartial (fun z => F z ν l) μ y
      + coordPartial (fun z => F z l μ) ν y = 0)

def ksRadius (a : ℝ) (y : E4) : ℝ :=
  let ρ2 := y 1 ^ 2 + y 2 ^ 2 + y 3 ^ 2
  Real.sqrt ((ρ2 - a ^ 2 + Real.sqrt ((ρ2 - a ^ 2) ^ 2 + 4 * a ^ 2 * y 3 ^ 2)) / 2)

def ksNull (a : ℝ) (y : E4) : Fin 4 → ℝ :=
  let r := ksRadius a y
  ![1, (r * y 1 + a * y 2) / (r ^ 2 + a ^ 2), (r * y 2 - a * y 1) / (r ^ 2 + a ^ 2), y
    ↪ 3 / r]

def ksProfile (m a e : ℝ) (y : E4) : ℝ :=
  let r := ksRadius a y
  r ^ 2 * (2 * m * r - e ^ 2) / (r ^ 4 + a ^ 2 * y 3 ^ 2)

def kerrNewmanMetric (m a e : ℝ) (y : E4) : Matrix (Fin 4) (Fin 4) ℝ :=
  fun μ ν => minkowski μ ν + ksProfile m a e y * ksNull a y μ * ksNull a y ν

def kerrNewmanPotential (a e : ℝ) (y : E4) : Fin 4 → ℝ :=
  let r := ksRadius a y
  fun μ => e * r ^ 3 / (r ^ 4 + a ^ 2 * y 3 ^ 2) * ksNull a y μ

def kerrNewmanField (a e : ℝ) (y : E4) : Matrix (Fin 4) (Fin 4) ℝ :=
  fun μ ν => coordPartial (fun z => kerrNewmanPotential a e z ν) μ y
    - coordPartial (fun z => kerrNewmanPotential a e z μ) ν y
```

B Platform objects

Platform theorem (all status *Proved*)	Identifier
`NoHair.kerrNewman_einsteinMaxwell`	`35932774-6f3d-4986-915d-e2e8baec247a`
`NoHair.kerrNewman_einstein`	`29511860-2b59-4995-a0aa-33f3e4ef6f27`
`NoHair.kerrNewman_maxwell`	`c5c86a9f-be51-4ff0-b03c-7af1195b1b3b`
`NoHair.kerrNewman_flat_maxwell`	`2a76a228-69dd-4f40-a775-d8e98e4f6349`
`NoHair.kerrNewman_field_closed`	`9c810d9f-30fa-4607-8131-cb6b04375499`
`NoHair.KNE.thm_christoffel`	`8dcd053d-2b42-4da1-b938-e3b583380372`
`NoHair.KNE.thm_div_one`	`6c168c47-f9a1-4c51-8479-700ca0814cee`
`NoHair.KNE.thm_div_two`	`36d617c4-dee0-4ad2-a404-b36e7ef9f8df`
`NoHair.KNE.thm_div_three`	`05c945f9-3d3d-4c04-b90f-a76537cfef0c`
`NoHair.KNE.thm_ricci_a`	`97f00edb-2452-4ddf-bc82-400a0b9946d5`
`NoHair.KNE.thm_ricci_b`	`eb386bc8-000f-4b46-b331-5014c2f82f08`
`NoHair.KNE.thm_ricci_c`	`e9f7c9ec-a103-4680-885e-9110e2332bfc`
`NoHair.KNE.thm_maxwell`	`a193ccd8-906e-420e-9057-670b9556dc7e`

The fifteen definition blocks are published as `NoHair_KerrNewmanEinstein_`X for X one of Base, Atoms, Christoffel, Div1–Div4, Ricci1–Ricci6, MaxwellF and MaxwellT.