


The slide features a light blue background. In the top right corner, there are several overlapping triangles in various shades of blue and purple. In the bottom right corner, there are overlapping triangles in shades of pink and red. A solid dark blue horizontal bar runs across the bottom of the slide.

Lecture 11: Primes and Greatest Common Divisors

Section 4.3

Section Summary

- Prime Numbers and their Properties
- Conjectures and Open Problems About Primes
- Greatest Common Divisors and Least Common Multiples
- The Euclidean Algorithm
- gcds as Linear Combinations

Primes

Definition : A positive integer p greater than 1 is called *prime* if the only positive factors of p are 1 and p . A positive integer that is greater than 1 and is not prime is called *composite*.

Example : The integer 7 is prime because its only positive factors are 1 and 7, but 9 is composite because it is divisible by 3.

Examples :

-

Finding Prime Numbers


If an integer n is a composite integer,
then it has a prime divisor less than or equal to $\sqrt{n}$.

To see this, note that if $n = ab$, then $a \leq \sqrt{n}$ or $b \leq \sqrt{n}$.

Trial division , a very **inefficient** method of determining if a number n is prime, is to try every integer $i \leq \sqrt{n}$ and see if n is divisible by i .

Sieve of Eratosthenes uses the trial division idea from the other way around.

continued →



Eratosthenes
(276-194 B.C.)

The Sieve of Eratosthenes

The *Sieve of Eratosthenes* can be used to find all primes not exceeding a specified positive integer. For example, begin with the list of integers between 1 and 100.

- a. Delete all the integers, other than 2, divisible by 2.
- b. Delete all the integers, other than 3, divisible by 3.
- c. Next, delete all the integers, other than 5, divisible by 5.
- d. Next, delete all the integers, other than 7, divisible by 7.
- e. Since all the remaining integers are not divisible by any of the previous integers, other than 1, the primes are: {2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47, 53, 59, 61, 67, 71, 73, 79, 83, 89, 97}

continued →



Eratosthenes
(276-194 B.C.)

The Sieve of Eratosthenes

TABLE 1 The Sieve of Eratosthenes.

<i>Integers divisible by 2 other than 2 receive an underline.</i>										<i>Integers divisible by 3 other than 3 receive an underline.</i>									
1	2	3	<u>4</u>	5	<u>6</u>	7	<u>8</u>	9	<u>10</u>	1	2	3	<u>4</u>	5	<u>6</u>	7	8	<u>9</u>	<u>10</u>
11	<u>12</u>	13	<u>14</u>	15	<u>16</u>	17	<u>18</u>	19	<u>20</u>	11	<u>12</u>	13	<u>14</u>	<u>15</u>	<u>16</u>	17	<u>18</u>	19	<u>20</u>
21	<u>22</u>	23	<u>24</u>	25	<u>26</u>	27	<u>28</u>	29	<u>30</u>	21	<u>22</u>	23	<u>24</u>	25	<u>26</u>	<u>27</u>	<u>28</u>	29	<u>30</u>
31	<u>32</u>	33	<u>34</u>	35	<u>36</u>	37	<u>38</u>	39	<u>40</u>	31	<u>32</u>	<u>33</u>	<u>34</u>	35	<u>36</u>	37	<u>38</u>	<u>39</u>	<u>40</u>
41	<u>42</u>	43	<u>44</u>	45	<u>46</u>	47	<u>48</u>	49	<u>50</u>	41	<u>42</u>	43	<u>44</u>	<u>45</u>	<u>46</u>	47	<u>48</u>	49	<u>50</u>
51	<u>52</u>	53	<u>54</u>	55	<u>56</u>	57	<u>58</u>	59	<u>60</u>	51	<u>52</u>	53	<u>54</u>	55	<u>56</u>	<u>57</u>	<u>58</u>	59	<u>60</u>
61	<u>62</u>	63	<u>64</u>	65	<u>66</u>	67	<u>68</u>	69	<u>70</u>	61	<u>62</u>	<u>63</u>	<u>64</u>	65	<u>66</u>	67	<u>68</u>	<u>69</u>	<u>70</u>
71	<u>72</u>	73	<u>74</u>	75	<u>76</u>	77	<u>78</u>	79	<u>80</u>	71	<u>72</u>	73	<u>74</u>	<u>75</u>	<u>76</u>	77	<u>78</u>	79	<u>80</u>
81	<u>82</u>	83	<u>84</u>	85	<u>86</u>	87	<u>88</u>	89	<u>90</u>	81	<u>82</u>	83	<u>84</u>	85	<u>86</u>	<u>87</u>	<u>88</u>	89	<u>90</u>
91	<u>92</u>	93	<u>94</u>	95	<u>96</u>	97	<u>98</u>	99	<u>100</u>	91	<u>92</u>	<u>93</u>	<u>94</u>	95	<u>96</u>	97	<u>98</u>	<u>99</u>	<u>100</u>
<i>Integers divisible by 5 other than 5 receive an underline.</i>										<i>Integers divisible by 7 other than 7 receive an underline; integers in color are prime.</i>									
1	2	3	<u>4</u>	5	<u>6</u>	7	<u>8</u>	<u>9</u>	<u>10</u>	1	2	3	<u>4</u>	5	<u>6</u>	7	8	<u>9</u>	<u>10</u>
11	<u>12</u>	13	<u>14</u>	<u>15</u>	<u>16</u>	17	<u>18</u>	19	<u>20</u>	11	<u>12</u>	<u>13</u>	<u>14</u>	<u>15</u>	<u>16</u>	<u>17</u>	<u>18</u>	<u>19</u>	<u>20</u>
21	<u>22</u>	23	<u>24</u>	<u>25</u>	<u>26</u>	27	<u>28</u>	29	<u>30</u>	21	<u>22</u>	<u>23</u>	<u>24</u>	<u>25</u>	<u>26</u>	<u>27</u>	<u>28</u>	<u>29</u>	<u>30</u>
31	<u>32</u>	<u>33</u>	<u>34</u>	<u>35</u>	<u>36</u>	37	<u>38</u>	<u>39</u>	<u>40</u>	31	<u>32</u>	<u>33</u>	<u>34</u>	<u>35</u>	<u>36</u>	<u>37</u>	<u>38</u>	<u>39</u>	<u>40</u>
41	<u>42</u>	43	<u>44</u>	<u>45</u>	<u>46</u>	47	<u>48</u>	49	<u>50</u>	41	<u>42</u>	43	<u>44</u>	<u>45</u>	<u>46</u>	<u>47</u>	<u>48</u>	49	<u>50</u>
51	<u>52</u>	53	<u>54</u>	<u>55</u>	<u>56</u>	<u>57</u>	<u>58</u>	59	<u>60</u>	51	<u>52</u>	<u>53</u>	<u>54</u>	55	<u>56</u>	<u>57</u>	<u>58</u>	<u>59</u>	<u>60</u>
61	<u>62</u>	<u>63</u>	<u>64</u>	<u>65</u>	<u>66</u>	67	<u>68</u>	69	<u>70</u>	61	<u>62</u>	<u>63</u>	<u>64</u>	<u>65</u>	<u>66</u>	<u>67</u>	68	69	<u>70</u>
71	<u>72</u>	73	<u>74</u>	<u>75</u>	<u>76</u>	77	<u>78</u>	79	<u>80</u>	71	<u>72</u>	<u>73</u>	<u>74</u>	<u>75</u>	<u>76</u>	<u>77</u>	<u>78</u>	<u>79</u>	<u>80</u>
81	<u>82</u>	83	<u>84</u>	<u>85</u>	<u>86</u>	87	<u>88</u>	89	<u>90</u>	81	<u>82</u>	<u>83</u>	<u>84</u>	85	<u>86</u>	<u>87</u>	<u>88</u>	<u>89</u>	<u>90</u>
91	<u>92</u>	<u>93</u>	<u>94</u>	<u>95</u>	<u>96</u>	97	<u>98</u>	<u>99</u>	<u>100</u>	91	<u>92</u>	<u>93</u>	<u>94</u>	95	<u>96</u>	<u>97</u>	<u>98</u>	<u>99</u>	<u>100</u>

Since, if $n = ab$, then $a \leq \sqrt{n}$ or $b \leq \sqrt{n}$.

For determining the prime numbers within 100, we only need to divide with primes that do not exceed $\sqrt{100} = 10$. Meaning simply checking 2, 3, 5 and 7 is enough to find all primes under 100.

Study of Primes

- **Distribution of primes:** The number of primes not exceeding x , can be approximated by $x/\ln x$. (The Prime Number Theorem)
- **Generating primes:** A closed formula that always produces primes has not been found so far. It is proven that there is no **polynomial** with integer coefficients such that $f(n)$ is prime for all positive integers n .
- **Conjectures on primes:**
 - **Goldbach's Conjecture:** Every even integer n , $n > 2$, is the sum of two primes.
 - There are infinitely many primes of the form $n^2 + 1$, where n is a positive integer. But it has been shown that there are infinitely many primes of the form $n^2 + 1$, where n is a positive integer or the product of at most two primes.
 - **The Twin Prime Conjecture:** There are infinitely many pairs of twin primes. Twin primes are pairs of primes that differ by 2. Examples are 3 and 5, 5 and 7, 11 and 13, etc.

Greatest Common Divisor

Definition : Let a and b be integers, not both zero. The largest integer d such that $d \mid a$ and also $d \mid b$ is called the greatest common divisor of a and b . The greatest common divisor of a and b is denoted by $\gcd(a,b)$.

One can find greatest common divisors of small numbers by inspection.

Example : What is the greatest common divisor of 24 and 36?

Solution : $\gcd(24, 36) = 12$

Example : What is the greatest common divisor of 17 and 22?

Solution : $\gcd(17, 22) = 1$

Co-primes / Relatively Primes

Definition : The integers a and b are *relatively prime* if their greatest common divisor is 1.

Example : 17 and 22

Definition : The integers $a_1, a_2, \dots, a_n$ are *pairwise relatively prime* if $\gcd(a_i, a_j) = 1$ whenever $1 \leq i < j \leq n$.

Example : Determine whether the integers 10, 17 and 21 are pairwise relatively prime.

Solution : Because $\gcd(10, 17) = 1$, $\gcd(10, 21) = 1$, and $\gcd(17, 21) = 1$, we can say that 10, 17, and 21 are pairwise relatively prime.

Example : Determine whether the integers 10, 19, and 24 are pairwise relatively prime.

Solution : Because $\gcd(10, 24) = 2$, the given numbers are not pairwise relatively prime.

Finding the Greatest Common Divisor Using Prime Factorizations

- Suppose the prime factorizations of a and b are:

$$a = p_1^{a_1} p_2^{a_2} \cdots p_n^{a_n}, \quad b = p_1^{b_1} p_2^{b_2} \cdots p_n^{b_n},$$

where each exponent is a nonnegative integer, and where all primes occurring in either prime factorization are included in both. Then:

$$\gcd(a, b) = p_1^{\min(a_1, b_1)} p_2^{\min(a_2, b_2)} \cdots p_n^{\min(a_n, b_n)}.$$

- Example :** $120 = 2^3 \cdot 3 \cdot 5$ $500 = 2^2 \cdot 5^3$
 $\gcd(120, 500) = 2^{\min(3, 2)} \cdot 3^{\min(1, 0)} \cdot 5^{\min(1, 3)} = 2^2 \cdot 3^0 \cdot 5^1 = 20$
- Finding the gcd of two positive integers using their prime factorizations is not efficient because there is no efficient algorithm for finding the prime factorization of a positive integer.

Least Common Multiple

Definition : The least common multiple of the positive integers a and b is the smallest positive integer that is divisible by both a and b . It is denoted by $\text{lcm}(a,b)$.

- The least common multiple can also be computed from the prime factorizations.

$$\text{lcm}(a, b) = p_1^{\max(a_1, b_1)} p_2^{\max(a_2, b_2)} \cdots p_n^{\max(a_n, b_n)}$$

No smaller number is divisible by a and b both.

Example: $\text{lcm}(2^3 3^5 7^2, 2^4 3^3) = 2^{\max(3,4)} 3^{\max(5,3)} 7^{\max(2,0)} = 2^4 3^5 7^2$

- The greatest common divisor and the least common multiple of two integers are related by:

Theorem 5: Let a and b be positive integers. Then

$$ab = \text{gcd}(a,b) \cdot \text{lcm}(a,b)$$

(*proof is Exercise 31*)



Euclid

(325 B.C.E. – 265 B.C.E.)

Euclidean Algorithm

- The Euclidean algorithm is an efficient method for computing the greatest common divisor of two integers. It is based on the idea that $\gcd(a, b)$ is equal to $\gcd(b, c)$ when $a > b$ and c is the remainder when a is divided by b .

Example : Find $\gcd(91, 287)$:

- $287 = 91 \cdot 3 + 14$ Divide 287 by 91

- $91 = 14 \cdot 6 + 7$ Divide 91 by 14

- $14 = 7 \cdot 2 + 0$ Divide 14 by 7

Stopping
condition

$$\gcd(287, 91) = \gcd(91, 14) = \gcd(14, 7) = 7$$

continued →

Correctness of Euclidean Algorithm

Lemma 1 : Let $a = bq + r$, where a , b , q , and r are integers. Then $\gcd(a,b) = \gcd(b,r)$.

Proof:

- Suppose that d divides both a and b .
- Then d also divides $a - bq = r$ (by Theorem 1 of Section 4.1).
- Hence, any common divisor of a and b must also be any common divisor of b and r .
- Suppose that d divides both b and r .
- Then d also divides $bq + r = a$.
- Hence, any common divisor of a and b must also be a common divisor of b and r .
- Therefore, $\gcd(a,b) = \gcd(b,r)$. ◀

Correctness of Euclidean Algorithm

- Suppose that a and b are positive integers with $a \geq b$.

Let $r_0 = a$ and $r_1 = b$.

Successive applications of the division algorithm yields:

$$\begin{aligned} r_0 &= r_1 q_1 + r_2 & 0 \leq r_2 < r_1, \\ r_1 &= r_2 q_2 + r_3 & 0 \leq r_3 < r_2, \\ &\vdots \\ &\vdots \\ &\vdots \\ r_{n-2} &= r_{n-1} q_{n-1} + r_n & 0 \leq r_n < r_{n-1}, \\ r_{n-1} &= r_n q_n. \end{aligned}$$

- Eventually, a remainder of zero occurs in the sequence of terms: $a = r_0 > r_1 > r_2 > \cdots \geq 0$. The sequence can't contain more than a terms.
- By Lemma 1, $\gcd(a, b) = \gcd(r_0, r_1) = \cdots = \gcd(r_{n-1}, r_n) = \gcd(r_n, 0) = r_n$.
- Hence the greatest common divisor is the last nonzero remainder in the sequence of divisions.

Dividing Congruences by an Integer

- Dividing both sides of a valid congruence by an integer does not always produce a valid congruence (see Section 4.1).
- But dividing by an integer **relatively prime** to the modulus does produce a valid congruence:

Theorem 7 : Let m be a positive integer and let a , b , and c be integers. If $ac \equiv bc \pmod{m}$ and $\gcd(c, m) = 1$, then $a \equiv b \pmod{m}$.

Proof: Since $ac \equiv bc \pmod{m}$, $m \mid ac - bc = c(a - b)$ by Lemma 2 and the fact that $\gcd(c, m) = 1$, it follows that $m \mid a - b$. Hence, $a \equiv b \pmod{m}$.