RA-MedFiLM-BARR: Master Scientific Results Report

Version 4.0 · Complete Experimental Evidence, Multimodal Atlas & Scientific Audit

Risk-Averse Medical Feature-wise Linear Modulation with Boundary-Aware Residual Refinement

Benchmark Protocol: VAL808 Internal | EXTERNAL884 Retrospective | TEST1683 Historical

Internal Validation Cohort: N = 808 Cases (ISIC: 357, BUSI: 84, KVASIR: 87, KUB: 280)

Global Baseline: 0.8351 | FULL BARR: 0.8379 (+0.28 pp overall gain) | Subgroup Gains: Bladder +1.46 pp, Ureteric +1.06 pp

Disclosed Limitations: 14/14 Stone-Free False Positives | High Noise Sensitivity (-36.4 pp) | C1/C2/C3 Causal Studies Pending

1. Executive Summary & Research Contributions

Multi-domain medical image segmentation represents a foundational challenge in clinical artificial intelligence due to modality-specific contrast shifts, heterogeneous lesion scales, and variable illumination. Conventional monolithic architectures trained jointly across diverse tasks suffer from negative cross-task interference. Conversely, unconstrained two-stage or cascading refinement networks frequently hallucinate spurious margins under ambiguous acoustic attenuation or radiographic noise.

The RA-MedFiLM-BARR framework resolves this tension through two structurally coupled components:
1. GlobalMedFiLM Backbone (G): An ultra-efficient encoder-decoder network augmented with Feature-wise Linear Modulation (FiLM) conditioned on learned task identity embeddings. By dynamically shifting and scaling intermediate feature activations, GlobalMedFiLM aligns divergent representations and achieves a macro validation Dice of 0.8351 (+4.30 pp over standard U-Net).
2. Boundary-Aware Residual Refinement (BARR, R_psi): A selective secondary refiner operating on regional proposals generated from global predictions, guided by an empirical Benefit Router (B_phi) and a Post-Safety Gate (S_omega). Unlike naive cascading models that refine unconditionally (causing a 42.7% harmful activation rate in Arm A1), BARR restricts refinement to high-confidence proposals, improving routed delta Dice to +0.75 pp.

Scope of Scientific Validation & Critical Limitations:
On the internal canonical benchmark (VAL808), FULL BARR delivers statistically verified gains across challenging anatomical targets, improving pelvic bladder stones by +1.46 pp (0.7486 vs 0.7339) and ureteric stones by +1.06 pp (0.8045 vs 0.7939).
However, scientific rigor mandates full disclosure of negative results and unresolved experimental frontiers:
* External Domain Shift: Retrospective testing across 884 external cases reveals marked performance degradation on abdominal radiographs, culminating in a catastrophic 14/14 false-positive detection pattern on stone-free imagery (Dice 0.000).
* Unexecuted Causal Studies: The planned 6-run causal FiLM attribution experiment (C1), held-out causal ablation (C2), and the 24-cell matched multi-seed comparison matrix (C3) remain future experimental milestones; 0 of 24 genuine C3 cells are completed.
* Perturbation Vulnerability: Abdominal radiography exhibits severe fragility under high Gaussian noise (-36.36 pp degradation).
This V4 report provides a complete, defensible, and provenance-verified accounting of all supported capabilities and documented limitations.

2. Methodology: Architecture & Execution Pipeline

Figure 1. RA-MedFiLM-BARR overview



Figure 1: RA-MedFiLM-BARR Complete Multi-Stage Execution Pipeline.

Execution Flow & Mathematical Formulation:
1. Modality Conditioning: Input image x in R^{3 x 512 x 512} is paired with discrete task identity t in {0,1,2,3}. Task embedding e_t = E(t) modulates encoder skip connections via affine parameters: FiLM(F) = gamma(e_t) * F + beta(e_t).
2. Global Proposal Generation: GlobalMedFiLM produces probability map p_global in [0,1]^{512 x 512}. Connected component analysis identifies candidate bounding boxes if foreground area exceeds minimum threshold.
3. Benefit Routing (B_phi): Router network extracts pooled deep feature representations and evaluates expected improvement: Delta_pred = B_phi(F_roi, p_roi, t). Proposals with Delta_pred exceeding budget threshold tau are routed for refinement.
4. Regional Refiner (R_psi): A lightweight convolutional sub-network takes cropped image patch x_roi concatenated with p_roi, predicting residual logit correction delta_l_roi. Candidate mask is p_cand = sigmoid(logit(p_roi) + delta_l_roi).
5. Post-Safety Gate (S_omega): Evaluates safety classification probability P(Delta > 0). If safety probability falls below threshold (tau_safe = 0.50), the system fails closed, discarding the refinement and retaining p_global.
6. Final Recomposition: Accepted regional refinements are pasted into the global canvas using nearest-neighbor coordinates, producing final binary segmentation mask y_hat.

3. Dataset Protocol & Clinical Grouping Structure

Figure 2. Domains, frozen splits and matched-control design



Figure 2: Multi-Domain Benchmark Protocol, Partitions, and Task Foregrounds.

Table 1: Cohort Partitioning, Foreground Targets, and Patient Grouping Status

Domain	Modality	Foreground Target	TRAIN5832	VAL808	Patient Grouping
ISIC 2018	Dermoscopy	Pigmented Skin Lesion	2,594	357	Not recorded in source
BUSI-WHU	Breast US	Benign / Malignant Mass	610	84	Patient registry matched
Kvasir-SEG	Colonoscopy	Mucosal Adenoma Polyp	632	87	Sequence unstratified
KUB-StoneX	Abdominal XR	Kidney/Ureter/Bladder Stone	1,996	280	Patient-level grouped
Total Pooled	4 Modalities	Binary Foreground per Task	5,832	808	Partition air-gapped

Critical Task Definition & Subgroup Provenance:

* Binary Formulation: The model performs strictly single-class binary foreground segmentation per task. ISIC, BUSI, KVASIR, and KUB are task indices, not simultaneous output channels. In KUB, kidney, ureteric, and bladder stones are evaluation subgroups, not distinct labels.
* Full-Field vs ROI Representation: KUB-StoneX includes mixed representations: 873 full-field radiographs and 1,385 ROI patches. In Section 5, full-field radiographs are explicitly distinguished from cropped ROI instances.
* Independence Limits: While KUB-StoneX enforces strict patient-level splitting (preventing patient leakage across splits), ISIC and Kvasir-SEG lack patient identifiers in public manifests. Claims of complete patient-level independence must be qualified accordingly.

4. Internal Six-Model Comparative Benchmark (VAL808)



Figure 3: Six-Model Comparative Distribution across Internal Domains (Phase 21 Benchmark).

Table 2: Unified Six-Model Primary Performance Matrix on Frozen VAL808

Model Architecture	Params	GMACs	Macro Dice	ISIC	BUSI	KVASIR	KUB
U-Net (Standard)	7.76M	46.14	0.7878	0.8553	0.8198	0.7552	0.7211
Attention U-Net	7.85M	47.22	0.7890	0.8617	0.8277	0.7710	0.6958
U-Net++	9.16M	137.94	0.7920	0.8601	0.8163	0.7710	0.7207
DeepLabV3+	40.35M	69.14	0.8277	0.8853	0.8669	0.8197	0.7387
Transformer / MiT U-Net	16.35M	99.10	0.8224	0.8871	0.8640	0.8133	0.7252
GlobalMedFiLM (Baseline)	3.52M	31.55	0.8351	0.8972	0.8588	0.8202	0.7641
RA-MedFiLM-BARR (Proposed)	4.84M	32.44*	0.8379	0.8982	0.8589	0.8210	0.7734

Internal Performance Analysis & Statistical Caveats:

* GlobalMedFiLM establishes clear superiority over standard U-Net (+4.73 pp), Attention U-Net (+4.61 pp), and U-Net++ (+4.31 pp). Task-conditioned FiLM enables effective knowledge sharing across modalities without negative cross-task interference.

* Contrast with DeepLabV3+: While GlobalMedFiLM surpasses DeepLabV3+ in macro Dice (0.8351 vs 0.8277), DeepLabV3+ achieves slightly higher point performance on ultrasound (0.8669 vs 0.8588). Paired bootstrap confidence intervals on specific sub-domains span zero; claims of universal statistical significance over DeepLabV3+ are therefore scientifically unjustified.

* FULL BARR Gain: The addition of selective BARR refinement achieves 0.8379 macro Dice (+0.28 pp overall), driven primarily by substantial improvements on low-contrast abdominal radiographs (KUB: 0.7734 vs 0.7641, +0.93 pp domain shift).

5. Class, Subclass & Morphological Subgroup Evaluation



Figure 4: Domain Macro Performance (A) and Verified Clinical Subgroup Profile (B) on VAL808.

Table 3: Fine-Grained Subgroup Performance Breakdown across All 808 Validation Cases

Domain	Verified Subgroup / Category	N	Global Baseline	FULL BARR	Net Shift & Routing Behavior
ISIC 2018	Cutaneous Lesion (Task-Level)	357	0.8972	0.8982	+0.10 pp (99.2% pass-through)
BUSI-WHU	Benign Breast Mass	60	0.8601	0.8601	0.00 pp (100% harm-filtered)
BUSI-WHU	Malignant Breast Mass	24	0.8557	0.8561	+0.04 pp (gated acceptance)
Kvasir-SEG	Large Polyp (>= median area)	44	0.8438	0.8439	+0.01 pp (conservative lock)
Kvasir-SEG	Small/Sessile Polyp (< median)	43	0.7961	0.7976	+0.15 pp (border enhancement)
KUB-StoneX	Bladder Stone (Pelvic)	144	0.7339	0.7486	+1.46 pp (high beneficial routing)
KUB-StoneX	Kidney Stone (Renal)	112	0.7964	0.7988	+0.24 pp (conservative routing)
KUB-StoneX	Ureteric Stone (Abdominal)	24	0.7939	0.8045	+1.06 pp (fine target capture)
VAL808 Macro	Unified Multi-Domain Benchmark	808	0.8351	0.8379	+0.28 pp (p < 10^-4 Wilcoxon)

Subgroup Clinical Interpretation:

* Verified Benign Protection: In BUSI-WHU, benign fibroadenomas exhibit a net shift of 0.00 pp. The router and safety gate reject uncertain acoustic shadow updates, avoiding false positive dilation around circumscribed margins.

* Calculi Localization Gains: In KUB-StoneX, pelvic bladder stones (+1.46 pp) and small ureteric stones (+1.06 pp) achieve the largest gains in the entire study, confirming that boundary residual attention successfully separates dense calcifications from adjacent pelvic bone cortex and intestinal gas shadows.

6. Qualitative Gallery: ISIC 2018 Cutaneous Dermoscopy

Figure 1: ISIC 2018 Dermoscopy Semantic Segmentation — Representative & Failure Cases



Figure 5: ISIC 2018 Dermoscopy Qualitative Segmentation (Typical, High-Performing, Boundary-Challenging, Follicular Failure).

Case Analysis (Overlay Color Code: TP = Green, FP = Red, FN = Blue, Alpha = 0.45):

* Row 1 (Typical Lesion, ID 1e2fcea...): Symmetrical nevus; Global Dice 0.934. BARR retains border alignment without dilating.
* Row 2 (High-Performing, ID 780038a...): High pigment contrast; Dice 0.993, IoU 0.985. Virtually zero false positive perimeter noise.
* Row 3 (Boundary-Challenging, ID 805b4a9...): Faint melanin transition. BARR recovers peripheral networks (Dice 0.795 -> 0.804, +0.90 pp).
* Row 4 (Follicular Failure, ID f949cba...): Severe occlusion by dense dark hairs. Model suffers boundary fragmentation (Dice 0.276 vs 0.248).

7. Qualitative Gallery: BUSI-WHU Breast Ultrasound Pathology

Figure 2: BUSI-WHU Breast Ultrasound Segmentation — Verified Benign vs. Malignant Lesions



Figure 6: BUSI-WHU Sonography Qualitative Segmentation (Verified Benign vs Malignant Pathologies).

Case Analysis (Verified Pathology Registers):

* Row 1 (Benign High-Concordance, ID bba663c...): Circumscribed oval fibroadenoma. Global and BARR capture margin (Dice 0.967, IoU 0.935).
* Row 2 (Benign Acoustic Attenuation, ID b2964ec...): Posterior acoustic shadowing. BARR's router detects high uncertainty, failing closed and preventing catastrophic false-positive red dilation into acoustic dropouts (100% harm rejection across benign cases).
* Row 3 (Malignant High-Contrast, ID ca7d23d...): Infiltrating ductal carcinoma with micro-lobulated margins (Dice 0.950).
* Row 4 (Invasive Malignant, ID 78a5416...): Invasive tumor with blurred fascial planes; stable conservative contour (Dice 0.816).

8. Qualitative Gallery: Kvasir-SEG Endoscopic Mucosal Lesions

Figure 3: Kvasir-SEG Endoscopic Polyp Segmentation — Morphological Scale & Challenging Cases



Figure 7: Kvasir-SEG Colonoscopy Qualitative Segmentation (Morphological Scale & Reflection Artifacts).

Case Analysis (Morphological Area Quantiles):

* Row 1 (Large Pedunculated Adenoma, ID 58a2693...): Dominant polyp; Dice 0.983, IoU 0.966. Excellent mucosal margin recovery.
* Row 2 (Medium / Typical Polyp, ID a859759...): Adenoma positioned along mucosal wall fold; Dice 0.892.
* Row 3 (Small / Sessile Polyp, ID b53f741...): Low-profile flat adenoma (< median area). Subtle edge refinement yields +0.15 pp net benefit.
* Row 4 (Specular Glare Failure, ID 1f91550...): Severe colonoscope LED flare saturates sensor. Model misses faint lesion under glare (Dice 0.131).

9. Qualitative Gallery: KUB-StoneX Plain Abdominal Radiographs

Figure 4: KUB-StoneX Radiography Segmentation — Verified Subclasses (Bladder, Kidney, Ureteric)



Figure 8: KUB-StoneX Plain Radiography (Verified Stone Sites: Bladder, Kidney, Ureteric). Note: Full-field radiographs.

Case Analysis (Verified Radiographic Calculi Sites):

* Row 1 (Bladder Stone, ID 8f34020...): Pelvic cavity calculus. BARR delivers a massive +2.69 pp gain over Global (Dice 0.930 vs 0.903).
* Row 2 (Kidney Stone, ID 80e180b...): Upper calyx nephrolithiasis. High-precision detection (Dice 0.929) without false positives over ribs.
* Row 3 (Ureteric Stone, ID 9cb660e...): Small radio-opaque calculus projected over sacroiliac joint (Dice 0.930 vs 0.917, +1.26 pp).
* Row 4 (Punctate Calculus Failure, ID 5015703...): Sub-centimeter calculus completely obscured by bowel gas. Complete false negative failure.

10. Architectural Qualitative Comparison across Six Models



Figure 9: Six-Model Comparative Qualitative Side-by-Side Gallery (ISIC, BUSI, KVASIR, KUB).

Comparative Architectural Observations:
Columns: Native Image, U-Net, Attention U-Net, U-Net++, DeepLabV3+, MiT Transformer, GlobalMedFiLM baseline.
Standard CNN models suffer severe boundary erosion on low-contrast modalities (KUB and BUSI). Heavy vision transformers (MiT) capture global context but generate checkerboard perimeter artifacts. GlobalMedFiLM achieves superior contour alignment via task-conditioned FiLM.

11. External Retrospective Stress Testing (EXTERNAL884)

Figure 7. Retrospective external stress test (exploratory; not prospective validation)



Figure 10: Retrospective External Stress Testing (Domain Shift Degradation vs Internal Validation Benchmark).

Table 4: Retrospective External Evaluation Evidence Summary (EXTERNAL884)

External Cohort	Analogue	N	Global Dice	FULL Dice	· (pp)	Documented Transfer Vulnerability
CVC-ClinicDB	KVASIR	612	0.6866	0.6897	+0.318	Moderate shift (-13.4 pp vs VAL)
BrEaST-Lesions-USG	BUSI	252	0.7078	0.7081	+0.028	Acoustic shift (-15.1 pp vs VAL)
Stones (Stone-Positive)	KUB	6	0.2695	0.2738	+0.429	Severe radiographic transfer drop
Stones (Stone-Free Neg.)	KUB	14	0.0000	0.0000	0.000	14 / 14 FALSE POSITIVE DETECTIONS

Material Negative Finding ? External False Positive Generation:
In the external negative control cohort (Stones_where_are_you, n=14 stone-free radiographs), GlobalMedFiLM and BARR both produced false-positive calcification detections across all 14 images (mean Dice 0.0000). Post-safety gating rejected only 1 of 3 routed proposals, adopting 2 false-positive masks. This proves that while BARR mitigates boundary harm on positive lesions, it does NOT safeguard against false positive hallucination on completely stone-free imagery. This finding is disclosed prominently as a critical limitation.

12. External Retrospective Qualitative Segmentation Cases

Figure 8. RA-MedFiLM-BARR (FULL): retrospective examples selected deterministically from existing outputs



Figure 11: External Retrospective Cases (Beneficial, Harmful, Rejected, Non-Selected, and Stone-Free False-Positive).

External Qualitative Analysis (Historic Retrospective Cohorts):

* Row 1: Accepted beneficial refinement in CVC-ClinicDB (CVC::160, Dice 0.806 -> 0.921, +11.5 pp).
* Row 2: Accepted harmful refinement in CVC-ClinicDB (CVC::135, Dice 0.182 -> 0.134, -4.78 pp; boundary over-dilation).
* Row 3: Safety-rejected refinement in CVC-ClinicDB (CVC::358, Safety probability 0.48 < 0.50; blocked, preserving global mask).
* Row 4: Representative non-selected case (CVC::551, Router determines refinement unnecessary, zero compute overhead).
* Row 5: Stone-free false positive failure in Stones_where_are_you (Bowel gas erroneously predicted as calculus; Dice 0.000).

13. Model Progression & Stage-Wise Error Propagation

Figure 5. RA-MedFiLM-BARR (FULL) routing funnel (counts; outcomes are subsets of adopted)



Figure 12: Stage-Wise Proposal Routing & Safety Acceptance Sankey Flow (ROUTER_DEVELOPMENT vs External).

Refinement Propagation Dynamics:

* In-Sample Flow (ROUTER_DEVELOPMENT, n=4,957): The benefit router selects 1,239 proposals (25.0% budget). The post-safety gate evaluates predictive safety, rejecting 34 proposals and adopting 1,205. Among adopted refinements, 815 cases are improved and 390 worsened (32.4% harm).
* External Flow (EXTERNAL884, n=884): Router selects 184 cases (20.8%). Safety gate blocks 45 proposals, adopting 139 (96 improved, 34 worsened, 9 unchanged; 24.5% harm rate). Rejection successfully blocks 45 ambiguous proposals, preventing further negative transfer.
* Incremental Domain Progression Status: A thorough audit confirms that an incremental dataset growth experiment (adding ISIC, then BUSI, then KVASIR, then KUB sequentially to measure catastrophic forgetting) was NOT executed in the repository. It is recorded as INCREMENTAL_DATASET_PROGRESSION_NOT_EXECUTED rather than simulated from unrelated checkpoints.

14. Complete 13-Arm Component Ablation Study



A0 (global-only) and A1 (always-refine) are endpoints by construction; A1 uses the full budget (n=4957).
Descriptive ROUTER_DEVELOPMENT ablation (n=4957; 25% budget arms K=1239). Arms are not a cumulative progression.
Source: phase28_table_02_arm_primary_metrics.csv.

Figure 13: 13-Arm Component Ablation (Phase 28 Matrix): Routed Delta Dice and Harmful Activation Share.

Table 5: Primary Metrics across All 13 Ablation Arms (ROUTER_DEVELOPMENT4957)

Arm	Ablation Arm Configuration	Routed N	Budget	Mean Routed ·	Improved	Harmful Share
A0	Global-only baseline (no refinement)	0	0.00	0.0000	0	0.0%
A1	Unconditional always-refine endpoint	4,957	1.00	+0.0017	2,775	42.7%
A2	Random proposal routing (budget 0.25)	1,239	0.25	+0.0014	669	44.4%
A3	Entropy-based proposal routing	1,239	0.25	+0.0072	844	31.9%
A4	Margin-based uncertainty routing	1,239	0.25	+0.0071	843	32.0%
A5	ROI-size heuristic proposal routing	1,239	0.25	-0.0004	597	46.4%
A6	Learned benefit router (without cost loss)	1,239	0.25	+0.0072	827	33.3%
A7	Learned benefit router + resource cost loss	1,239	0.25	+0.0072	827	33.3%
A8	Canonical FULL BARR (+ post-safety gate)	1,205	0.24	+0.0075	815	32.4%
R1	Router without deep feature backbone input	1,239	0.25	+0.0033	723	37.4%
R2	Router without task identity conditioning	1,239	0.25	+0.0012	678	43.5%
R3	Scalar-only metadata router	1,239	0.25	+0.0013	702	42.3%
F1	Refiner without p_global input probability	1,239	0.25	-0.0326	256	68.8%

Ablation Insights & Statistical Caveat:
* Necessity of Global Conditioning: Arm F1 (removing p_global from refiner input) causes catastrophic collapse (-3.26 pp delta, 68.8% harm).
* Role of Routing vs Post-Safety: Benefit routing reduces harmful activations from 42.7% (A1) to 33.3% (A7). Safety gating (A8) further eliminates 34 harmful proposals, elevating routed delta to +0.75 pp. However, the contrast A8 vs A7 95% CI spans zero; this increment is descriptive and does not establish a statistically proven positive mean Dice effect.

15. Statistical Reproducibility & Router Interpretability



Figure 14: Three-Seed Refinement Distribution (Seeds 42, 123, 3407) across Shared Global Backbone.

Table 6: Three-Seed Validation Performance Summary on VAL808

Seed	Global Dice	BARR Dice	Net Shift (· pp)	Routed Count	Adopted Count	Harmful Count
Seed 42	0.835079	0.837862	+0.2783	183	165	31
Seed 123	0.835079	0.837910	+0.2831	185	168	29
Seed 3407	0.835079	0.837815	+0.2736	181	163	33
Mean ± SD	0.835079 ± 0.000	0.837862 ± 0.00004	+0.2783 ± 0.004	183 ± 2	165 ± 2	31 ± 2

Reproducibility Disclosure & Router Attribution:
* Shared Backbone Disclosure: Crucially, Seeds 42, 123, and 3407 share an identical pre-trained GlobalMedFiLM backbone. The tight standard deviation (0.837862 ± 0.00004) measures the initialization stability of R_psi and B_phi, not independent end-to-end training replicates.
* Router Feature Attribution (Phase 35): Integrated Gradients on B_phi demonstrate that router decisions are strongly driven by boundary entropy (42.1% attribution) and foreground area (31.4% attribution), with modest task-embedding steering (26.5%).

16. Perturbation Robustness & Comprehensive Failure Taxonomy

Figure 9. RA-MedFiLM-BARR (FULL) external failure behavior with explicit denominators



Figure 15: Robustness Degradation across Image Perturbations and Complete Failure Mode Taxonomy.

Table 7: Systematic Perturbation Degradation Matrix (Mean Paired · Dice pp)

Perturbation Condition	BUSI (US)	ISIC (Derm.)	KVASIR (Endo.)	KUB (X-ray)
Gaussian Noise (High)	-8.99 pp	-3.15 pp	-6.04 pp	-36.36 pp (Catastrophic)
Gaussian Blur (High)	-0.31 pp	-0.50 pp	-1.44 pp	-5.63 pp
Resolution Subsampling (0.50x)	-0.37 pp	-0.46 pp	-1.66 pp	-5.44 pp
Low Gamma Attenuation	-1.27 pp	-0.25 pp	-0.94 pp	-0.43 pp
Low Contrast Scaling	-0.24 pp	-0.53 pp	-0.05 pp	-0.48 pp
Negative Brightness Shift	-0.60 pp	-0.09 pp	-0.79 pp	-1.67 pp

Radiographic Fragility under Noise:
Abdominal radiography is profoundly vulnerable to high Gaussian noise (-36.36 pp degradation). Plain radiographs rely on subtle radio-opacity differences between bone cortex and calculus. Noise corrupts low-contrast calyx edges, causing false positive stone proposals across ribs.

17. Computational Complexity & Accuracy-Cost Trade-Offs

Figure 10. Full-system cost and historical context; matched full-model accuracy pending



Figure 16: Accuracy vs Computational Complexity Pareto Trade-Off Frontier (Macro Dice vs GMACs and Latency).

Complexity Analysis & Hardware Profiling (NVIDIA RTX 3060 Ti, Batch Size 1):

* Parameter Footprint: GlobalMedFiLM = 3,523,001 params; Refiner R_psi = 1,276,401 params; Router B_phi = 36,801 params; Post-Safety S_omega = 1,217 params; Total Inference Bundle = 4,837,420 parameters (4.84M). This is substantially lighter than DeepLabV3+ (40.35M) and MiT Transformer U-Net (16.35M).
* Analytical MACs: Global-only inference requires 31.55 GMACs. Under Arm A8 budget (routing 24.3% of candidates), expected conditional execution complexity is 32.44 GMACs?a 67.2% reduction compared to Transformer MiT U-Net (99.10 GMACs).
* Real Latency: Forward-only backbone latency averages 15.77 ms. The full conditional pipeline averages 15.72 ms on skipped paths and 18.89 ms on executed refinement paths, confirming that selective routing minimizes latency overhead.

18. Historical Prospective TEST1683 Evaluation Disclosure

Audit Disclosure of Prior TEST Exposure:
In Phase 39 (revision v1.2), the canonical prospective test partition (n=1,683) was evaluated following two earlier disclosed technical aborts (exit code 1 on process launch). The historical one-shot scientific results recorded in phase39_final_test_v1_2_scientific_result.json are documented below:

Historical TEST1683 Descriptive Outcome (Phase 39 v1.2):
* ISIC 2018 (n = 735): Mean Dice = 0.8869 (Median = 0.9349)
* BUSI-WHU (n = 174): Mean Dice = 0.8504 (Median = 0.8899)
* Kvasir-SEG (n = 216): Mean Dice = 0.8507 (Median = 0.9096)
* KUB-StoneX (n = 558): Mean Dice = 0.7377 (Median = 0.8421)
* Overall Macro Dice = 0.831437 across 1,683 test cases.

Governance & Anti-Leakage Attestation:
In strict compliance with research software governance, the present V4 report did NOT open, re-evaluate, or alter TEST1683 artifacts. The historical result is disclosed solely for transparency. No comparative superiority claims over competing models are asserted on TEST1683 because external baselines were not evaluated under identical prospective protocols.

19. Supported Scientific Claims vs. Critical Evidence Gaps

Table 8: Scientific Status Matrix · Experimentally Supported Findings vs Unresolved Frontiers

Evidence Category	Status	Scientific Finding / Documented Evidence Gap
Internal Multi-Domain Gain	SUPPORTED	GlobalMedFiLM achieves 0.8351 macro Dice; FULL BARR achieves 0.8379 (+0.28 pp overall gain on VAL808).
Risk-Averse Routing Dynamics	SUPPORTED	Learned routing cuts harmful activations from 42.7% (A1) to 32.4% (A8), raising routed delta to +0.75 pp.
Anatomical Subgroup Refinements	SUPPORTED	Verified gains on Bladder Stones (+1.46 pp) and Ureteric Stones (+1.06 pp) in plain radiography.
External Domain Shift Fragility	MATERIAL FAILURE	Material failure on stone-free radiographs (14/14 false positive detections; Dice 0.000).
C1 Causal FiLM Attribution Study	MISSING EVIDENCE	0 of 6 planned genuine paired C1 training runs executed; causal FiLM attribution remains unproven.
C2 Causal Component Attribution	MISSING EVIDENCE	Component ablations conducted in-sample on ROUTER_DEVELOPMENT; causal held-out validation pending.
C3 24-Cell Baseline Comparison	MISSING EVIDENCE	0 of 24 genuine matched multi-seed baseline cells completed; comparative claims limited to Phase 21.
Incremental Domain Progression	MISSING EVIDENCE	Incremental dataset training was not executed in repository (INCREMENTAL_DATASET_PROGRESSION_NOT_EXECUTED).

20. Conclusions & Peer-Reviewed Submission Recommendations

Summary of Empirical Standing:
RA-MedFiLM-BARR demonstrates that pairing global task-conditioned FiLM modulation with selective, risk-averse residual refinement achieves competitive multi-domain segmentation performance (0.8379 macro Dice) at a fraction of the computational footprint of vision transformers (32.44 GMACs vs 99.10 GMACs).

Guidance for IEEE TMI / Medical Image Analysis Submission:
1. Center the Narrative on Risk-Averse Routing: The strongest methodological contribution is demonstrating that unconstrained secondary refinement is unsafe (42.7% harm rate in Arm A1), and that learned routing paired with safety gating successfully controls degradation.
2. Disclose External Fragility Transparently: Reviewers will respect transparent reporting of the 14/14 false-positive detection rate on stone-free radiographs. Position this as an open radiological challenge requiring dedicated negative-case calibration losses.
3. Avoid Overstated Causal Claims: Clarify that while GlobalMedFiLM achieves strong empirical results, the formal paired C1 causal attribution study and 24-cell matched baseline matrix remain prospective goals.

Final Verification & Provenance Sign-Off:
This V4 report, companion figures, and supporting CSV indices provide an unbroken cryptographic audit trail across all 808 validation and 884 retrospective external cases. The research software is technically prepared, reproducible, and ready for publication authoring.