


All Students Take Calculus I-All pos. II-sin III-tan IV-cos	Law of Cosines $a^2 = b^2 + c^2 - 2bc \cos A$	Difference of Cubes $a^3 \pm b^3 = (a \pm b)(a^2 \mp ab + b^2)$	Arc Length $s = r\theta$
Heron's Formula $A = \sqrt{s(s-a)(s-b)(s-c)}$	Change of Base $\log_b m = \frac{\log m}{\log b}$	Choose Formula $C(x, y) = \binom{x}{y} = \frac{x!}{y!(x-y)!}$	Law of Sines $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$
Degrees to Radians $\frac{A \cdot \pi}{180} = \theta$	Sector Area $A = \frac{1}{2}r^2\theta$	Area of Δ $Area = ab \cdot \frac{1}{2} \sin C$	Polar to (x, y) $r^2 = x^2 + y^2$ $\tan \theta = \frac{y}{x}$
			$(\log_a b)(\log_c d) = (\log_a d)(\log_c b)$
$z = a + bi$ $ z = \sqrt{a^2 + b^2}$	$z = r \operatorname{cis} \theta$ $z^n = r^n \operatorname{cis}(n\theta)$	n th roots of $z = r \operatorname{cis} \theta$ $w_k = r^{1/n} \operatorname{cis} \left(\frac{\theta + 2k\pi}{n} \right)$	$z_1 z_2 = r_1 r_2 \operatorname{cis}(\theta_1 + \theta_2)$ $\frac{z_1}{z_2} = \frac{r_1}{r_2} \operatorname{cis}(\theta_1 - \theta_2)$
			$\vec{v} = \langle a, b \rangle = a\hat{i} + b\hat{j}$ $ \vec{v} = \sqrt{a^2 + b^2}$
			$ c\vec{u} = c \vec{u} $
Dot Product $\vec{u} \cdot \vec{v} = a_1 a_2 + b_1 b_2$	Dot Product Theorem $\vec{u} \cdot \vec{v} = \vec{u} \vec{v} \cos \theta$	θ between \vec{u} & \vec{v} $\cos \theta = \frac{\vec{u} \cdot \vec{v}}{ \vec{u} \vec{v} }$	\vec{u} and \vec{v} are perpendicular $\vec{u} \cdot \vec{v} = 0$
			Component of \vec{u} along \vec{v} $(\vec{u} \cdot \vec{v})/ \vec{v} $
			$\operatorname{proj}_{\vec{v}} \vec{u} = \left(\frac{\vec{u} \cdot \vec{v}}{ \vec{v} ^2} \right) \vec{v}$
			Work $W = \vec{F} \cdot \vec{D}$
Trig Identities			
$\sin^2 + \cos^2 = 1$	$\tan^2 + 1 = \sec^2$	$1 + \cot^2 = \csc^2$	$2 \sin u \cos u = \sin(2u)$
			$\cos^2 u - \sin^2 u = \cos(2u)$
			$\frac{2 \tan u}{1 - \tan^2 u} = \tan(2u)$
$\sin u \cos v \pm \cos u \sin v = \sin(u \pm v)$	$\cos u \cos v \mp \sin u \sin v = \cos(u \pm v)$	$\frac{\tan u \pm \tan v}{1 \mp \tan u \tan v} = \tan(u \pm v)$	$\cot = \frac{1}{\tan}$
$\csc = \frac{1}{\sin}$	$\sec = \frac{1}{\cos}$	$\sin\left(\frac{\pi}{2} - u\right) = \cos u$	$\tan\left(\frac{\pi}{2} - u\right) = \cot u$
		$\sec\left(\frac{\pi}{2} - u\right) = \csc u$	$\cos\left(\frac{\pi}{2} - u\right) = \sin u$
$\cot\left(\frac{\pi}{2} - u\right) = \tan u$	$\csc\left(\frac{\pi}{2} - u\right) = \sec u$	$\frac{1 - \cos 2x}{2} = \sin^2 x$	$\frac{1 + \cos 2x}{2} = \cos^2 x$
		$\frac{1 - \cos 2x}{1 + \cos 2x} = \tan^2 x$	$\pm \sqrt{\frac{1 - \cos u}{2}} = \sin \frac{u}{2}$
$\pm \sqrt{\frac{1 + \cos u}{2}} = \cos \frac{u}{2}$	$\frac{1 - \cos u}{\sin u} = \frac{\sin u}{1 + \cos u} = \tan \frac{u}{2}$	$2 \sin \frac{x+y}{2} \cos \frac{x-y}{2} = \sin x \pm \sin y$	$2 \cos \frac{x+y}{2} \cos \frac{x-y}{2} = \cos x + \cos y$
$-2 \sin \frac{x+y}{2} \sin \frac{x-y}{2} = \cos x - \cos y$	$\sin u \cos v = \frac{1}{2}[\sin(u+v) + \sin(u-v)]$	$\cos u \sin v = \frac{1}{2}[\sin(u+v) - \sin(u-v)]$	
	$\cos u \cos v = \frac{1}{2}[\cos(u+v) + \cos(u-v)]$	$\sin u \sin v = \frac{1}{2}[\cos(u+v) - \cos(u-v)]$	

<p style="text-align: center;">Algebra of Functions</p> <p>Let f and g be functions with domains A and B.</p> <p>$(f + g)(x) = f(x) + g(x)$ Domain $A \cap B$</p> <p>$(f - g)(x) = f(x) - g(x)$ Domain $A \cap B$</p> <p>$(fg)(x) = f(x)g(x)$ Domain $A \cap B$</p> <p>$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$ Domain $\{x \in A \cap B \mid g(x) \neq 0\}$</p> <p>$(f \circ g)(x) = f(g(x))$ Domain $\{x \in B \mid g(x) \in A\}$</p>	<p style="text-align: center;">Polynomial Synthetic Division</p> $(x^3 + x^2 - 1) \div (x + 2)$ <table style="margin-left: auto; margin-right: auto;"> <tr><td style="border-right: 1px solid black; padding: 5px 10px;">- 2</td><td style="padding: 5px 10px;">1</td><td style="padding: 5px 10px;">1</td><td style="padding: 5px 10px;">0</td><td style="padding: 5px 10px;">- 1</td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;"></td><td style="padding: 5px 10px;"></td><td style="padding: 5px 10px;">- 2</td><td style="padding: 5px 10px;">2</td><td style="padding: 5px 10px;">- 4</td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;"></td><td style="padding: 5px 10px;">1</td><td style="padding: 5px 10px;">- 1</td><td style="padding: 5px 10px;">2</td><td style="padding: 5px 10px;">- 5</td></tr> </table>	- 2	1	1	0	- 1			- 2	2	- 4		1	- 1	2	- 5	<p style="text-align: center;">Polynomial Long Division</p> $x^2 - x + 2$ <table style="margin-left: auto; margin-right: auto;"> <tr><td style="border-right: 1px solid black; padding: 5px 10px;">$x + 2$</td><td style="padding: 5px 10px;">$x^3 + x^2$</td><td style="padding: 5px 10px;">$- 1$</td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;"></td><td style="padding: 5px 10px;">$- x^3 - 2x^2$</td><td style="padding: 5px 10px;"></td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;"></td><td style="padding: 5px 10px;">$- x^2$</td><td style="padding: 5px 10px;"></td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;"></td><td style="padding: 5px 10px;">$x^2 + 2x$</td><td style="padding: 5px 10px;"></td></tr> <tr><td style="border-right: 1px solid black; padding: 5px 10px;"></td><td style="padding: 5px 10px;">$2x - 1$</td><td style="padding: 5px 10px;"></td></tr> </table>	$x + 2$	$x^3 + x^2$	$- 1$		$- x^3 - 2x^2$			$- x^2$			$x^2 + 2x$			$2x - 1$	
- 2	1	1	0	- 1																												
		- 2	2	- 4																												
	1	- 1	2	- 5																												
$x + 2$	$x^3 + x^2$	$- 1$																														
	$- x^3 - 2x^2$																															
	$- x^2$																															
	$x^2 + 2x$																															
	$2x - 1$																															

<p style="text-align: center;">Rational Roots Theorem</p> <p>$2x^3 + 2x^2 - 3x - 6$</p> <p>$\pm 1, \pm 2$ $\pm 1, \pm 2, \pm 3, \pm 6$</p> <p>Possible rational roots:</p> <p>$\pm 1, \pm \frac{1}{2}, \pm 2, \pm 3, \pm \frac{3}{2}, \pm 6$</p>	<p style="text-align: center;">Decartes' Rule of Signs</p> <p>Count num. of sign changes</p> <p>$P(x) = 3x^6 + 4x^5 + 3x^3 - x - 3$</p> <p>1 positive real root</p> <p>$P(-x) = 3x^6 - 4x^5 - 3x^3 + x - 3$</p> <p>1 or 3 negative real roots</p>	<p style="text-align: center;">Logarithm Formulas</p> <p>$\log(m \cdot n) = \log m + \log n$</p> <p>$\log\left(\frac{m}{n}\right) = \log m - \log n$</p> <p>$\log(m^n) = n \cdot \log m$</p> <p>$\log_b b^x = x = b^{\log_b x}$</p>	<p style="text-align: center;">Other trig stuff</p> <p>$\cot = \frac{1}{\tan}$</p> <p>$\csc = \frac{1}{\sin}$</p> <p>$\sec = \frac{1}{\cos}$</p>
--	--	--	--

<p style="text-align: center;">Horizontal Asymptotes</p> <p>$y = \frac{2x^2 - 4x + 5}{x^2 - 2x + 1}$ Original Equation</p> <p>$= \frac{2x^2}{x^2}$ $x \rightarrow \infty$, other terms \rightarrow tiny</p> <p>$= 2$ Cancel, horizontal asymptote</p>	<p style="text-align: center;">Slant Asymptotes</p> <p>$y = \frac{x^2 - 4x - 5}{x - 3}$ Original Equation</p> <p>$= x - 1 - \frac{8}{x - 3}$ Divide</p> <p>$= x - 1$ $x \rightarrow \infty$, other terms \rightarrow tiny</p>
--	--

<p style="text-align: center;">Vertical Asymptotes</p> <p>$y = \frac{2x^2 - 4x + 5}{x^2 - 2x + 1}$ Original Equation</p> <p>$= \frac{2x^2 - 4x + 5}{(2x - 1)(x + 2)}$ Factor demoniator</p> <p>$x = \frac{1}{2}$ or $x = -2$ Impossible</p>	<p style="text-align: center;">End Behavior</p> <table style="width: 100%;"> <tr> <td style="text-align: center;">$y = x^n$ n is even</td> <td style="text-align: center;">$y = -x^n$ n is even</td> </tr> <tr> <td style="text-align: center;">$y = x^n$ n is odd</td> <td style="text-align: center;">$y = -x^n$ n is odd</td> </tr> </table>	$y = x^n$ n is even	$y = -x^n$ n is even	$y = x^n$ n is odd	$y = -x^n$ n is odd	<p style="text-align: center;">$m \times n$ matrix</p> $\begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$
$y = x^n$ n is even	$y = -x^n$ n is even					
$y = x^n$ n is odd	$y = -x^n$ n is odd					

<p style="text-align: center;">$y = \sin x$ in red; $y = \cos x$ in blue</p> 	<p style="text-align: center;">sin/cos Graph Properties</p> <p>If in form:</p> <p>$y = a \sin k(x - b)$</p> <p>amplitude a, period $2\pi/k$, phase shift b</p>	<p style="text-align: center;">Allowed row operations</p> <ol style="list-style-type: none"> 1. Add a multiple of one row to another 2. Multiply a row by a nonzero constant 3. Interchange two rows
---	---	--

<p style="text-align: center;">Row-Echelon Form</p> $\begin{bmatrix} 1 & 2 & -1 & 1 \\ 0 & 1 & 4 & -7 \\ 0 & 0 & 1 & -2 \end{bmatrix}$	<p style="text-align: center;">Reduced Row-Echelon Form</p> $\begin{bmatrix} 1 & 0 & 0 & -3 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & -2 \end{bmatrix}$	<p style="text-align: center;">Using matrix inverses ($AX = B \Rightarrow X = A^{-1}B$)</p> $\begin{bmatrix} 2 & -5 \\ 3 & -6 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 15 \\ 36 \end{bmatrix} \Rightarrow \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -2 & \frac{5}{3} \\ -1 & \frac{3}{2} \end{bmatrix} \begin{bmatrix} 15 \\ 36 \end{bmatrix} = \begin{bmatrix} 30 \\ 9 \end{bmatrix}$
---	--	---

Matrix Multiplication!

$$\begin{bmatrix} 1 & 3 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} -1 & 5 & 2 \\ 0 & 4 & 7 \end{bmatrix} = \begin{bmatrix} 1 \cdot (-1) + 3 \cdot 0 & 1 \cdot 5 + 3 \cdot 4 & 1 \cdot 2 + 3 \cdot 7 \\ (-1) \cdot (-1) + 0 \cdot 0 & (-1) \cdot 5 + 0 \cdot 4 & (-1) \cdot 2 + 0 \cdot 7 \end{bmatrix} = \begin{bmatrix} -1 & 17 & 23 \\ 1 & -5 & -2 \end{bmatrix}$$

<p style="text-align: center;">2×2 Matrix Inverse</p> <p>If $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ then $A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$</p>	<p style="text-align: center;">$n \times n$ Matrix Inverse</p> $\left[\begin{array}{ccc ccc} 1 & -2 & -4 & 1 & 0 & 0 \\ 2 & -3 & -6 & 0 & 1 & 0 \\ -3 & 6 & 15 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc ccc} 1 & 0 & 0 & -3 & 2 & 0 \\ 0 & 1 & 0 & -4 & 1 & -\frac{2}{3} \\ 0 & 0 & 1 & 1 & 0 & \frac{1}{2} \end{array} \right]$
--	---