

Square Roots $\sqrt{x^6} = x^3 $ $\sqrt{x^8} = x^4$ $\sqrt{x^7} = x^3\sqrt{x}$	Absolute Value Inequalities $ x < c \quad -c < x < c$ $ x > c \quad x < -c \text{ or } c < x$	Distance Formula $A(x_1, y_2) \text{ and } B(x_2, y_2)$ $d(A, B) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$		Midpoint Formula $A(x_1, y_2) \text{ and } B(x_2, y_2)$ $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$
Equation of a Circle $(x-h)^2 + (y-k)^2 = r^2$	Point-Slope Form $y - y_1 = m(x - x_1)$	Standard Form $Ax + By + C = 0$	All Students Take Calculus I-All pos. II-sin III-tan IV-cos	Law of Cosines $a^2 = b^2 + c^2 - 2bc \cdot \cos A$
Joint Variation If z is varies jointly as x and y , $z = kxy$	Perpendicular Lines $m_2 = -\frac{1}{m_1}$	Average Rate of Change $\text{ARoC} = \frac{y \text{ change}}{x \text{ change}} = \frac{f(x_2) - f(x_1)}{x_2 - x_1}$		Difference of Cubes $a^3 + b^3 = (a+b)(a^2 - ab + b^2)$ $a^3 - b^3 = (a-b)(a^2 + ab + b^2)$
Standard Form of a Quadratic Function $f(x) = a(x - h)^2 + k$				
Vertical Shifts of Graphs Suppose $c > 0$. Graph $y = f(x) + c$ by shifting $y = f(x)$ up c . Graph $y = f(x) - c$ by shifting $y = f(x)$ down c .		Horizontal Shifts of Graphs Suppose $c > 0$. Graph $y = f(x - c)$ by shifting $y = f(x)$ right c . Graph $y = f(x + c)$ by shifting $y = f(x)$ left c .		Definition of Log if $a^x = y$, $\log_a y = x$
Arc Length $s = r\theta$				
Reflecting Graphs Graph $y = -f(x)$ by reflecting $y = f(x)$ in the x -axis. Graph $y = f(-x)$ by reflecting $y = f(x)$ in the y -axis.		Vertical Stretching of Graphs To graph $y = cf(x)$, graph $y = f(x)$, then if $c > 1$ stretch vertically a by factor of c if $0 < c < 1$ shrink vertically a by factor of c		Inverse Variation If y is inversly proportional to x . $y = \frac{k}{x}$
Horizontal Stretching of Graphs To graph $y = f(cx)$, graph $y = f(x)$, then if $c > 1$ shrink horizontally by a factor of $\frac{1}{c}$ if $0 < c < 1$ stretch horizontally by a factor of $\frac{1}{c}$		Even and Odd Functions if $f(-x) = f(x)$ $f(x)$ is even if $f(-x) = -f(x)$ $f(x)$ is odd		Heron's Formula $A = \sqrt{s(s-a)(s-b)(s-c)}$
Min or Max of a Quadratic Function $f(x) = x(x - h)^2 + k \quad f(h) = k$ $f(x) = ax^2 + bx + c \quad f(-\frac{b}{2a})$		Change of Base $\log_b m = \frac{\log m}{\log b}$	Completing the Square With a quadratic in form $ax^2 + bx = c$ $(\frac{1}{2} \cdot b)^2 = c$	Hidden quadratic 1 $x^{-3/2} + 2x^{-1/2} + x^{1/2}$ $x^{-3/2}(1 + 2x + x^2)$ $x^{-3/2}(1 + x)^2$
Hidden quadratic 2 $e^{2x} + 2e^x + 1$ $(e^x + 1)^2$				
Permutations $p(x, y) = \frac{x!}{(x - y)!}$	Choose Formula $C(x, y) = \binom{x}{y} = \frac{x!}{y!(x - y)!}$	Law of Sines $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$	Degrees to Radians $\frac{A \cdot \pi}{180} = \theta$	Remainder Theorem If $P(x) \div (x - c)$, the remainder = $P(c)$.
SOH-CAH-TOA $\sin = \frac{\text{opp}}{\text{hyp}} \quad \cos = \frac{\text{adj}}{\text{hyp}} \quad \tan = \frac{\text{opp}}{\text{adj}}$		Sector Area $A = \frac{1}{2}r^2\theta$	Direct Variation If y is directly proportional to x , $y = kx$	Population Growth n is population size, r is relative growth rate, t is time $n = n_0e^{rt}$
Area of Δ $A = ab \cdot \frac{1}{2} \sin C$				
Property of logs $(\log_a b)(\log_c d) = (\log_a d)(\log_c b)$				

Algebra of Functions		Polynomial Synthetic Division		Polynomial Long Division			
Let f and g be functions with domains A and B .		$(x^3 + x^2 - 1) \div (x + 2)$		$x^2 - x + 2$			
$(f + g)(x) = f(x) + g(x)$	Domain $A \cap B$	- 2	1	1	0	- 1	
$(f - g)(x) = f(x) - g(x)$	Domain $A \cap B$			- 2	2	- 4	
$(fg)(x) = f(x)g(x)$	Domain $A \cap B$						
$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$	Domain $\{x \in A \cap B \mid g(x) \neq 0\}$						
$(f \circ g)(x) = f(g(x))$	Domain $\{x \in B \mid g(x) \in A\}$			1	- 1	2	- 5

<p>Rational Roots Theorem</p> <p>$2x^3 + 2x^2 - 3x - 6$</p> <p>$\pm 1, \pm 2$ $\pm 1, \pm 2, \pm 3, \pm 6$</p> <p>Possible rational roots:</p> <p>$\pm 1, \pm \frac{1}{2}, \pm 2, \pm 3, \pm \frac{3}{2}, \pm 6$</p>	<p>Decartes' Rule of Signs</p> <p>Count num. of sign changes</p> <p>$P(x) = 3x^6 + 4x^5 + 3x^3 - x - 3$</p> <p>1 positive real root</p> <p>$P(-x) = 3x^6 - 4x^5 - 3x^3 + x - 3$</p> <p>1 or 3 negative real roots</p>	<p>Logarithm Formulas</p> <p>$\log(m \cdot n) = \log m + \log n$</p> <p>$\log\left(\frac{m}{n}\right) = \log m - \log n$</p> <p>$\log(m^n) = n \cdot \log m$</p> <p>$\log_b b^x = x = b^{\log_b x}$</p>	<p>Other trig stuff</p> <p>$\cot = \frac{1}{\tan}$</p> <p>$\csc = \frac{1}{\sin}$</p> <p>$\sec = \frac{1}{\cos}$</p>
--	---	---	---

<p>Horizontal Asymptotes</p> <p>$y = \frac{2x^2 - 4x + 5}{x^2 - 2x + 1}$ Original Equation</p> <p>$= \frac{2x^2}{x^2}$ $x \rightarrow \infty$, other terms \rightarrow tiny</p> <p>$= 2$ Cancel, horizontal asymptote</p>	<p>Slant Asymptotes</p> <p>$y = \frac{x^2 - 4x - 5}{x - 3}$ Original Equation</p> <p>$= x - 1 - \frac{8}{x - 3}$ Divide</p> <p>$= x - 1$ $x \rightarrow \infty$, other terms \rightarrow tiny</p>
--	--

Vertical Asymptotes		End Behavior		Trig Identities
$y = \frac{2x^2 - 4x + 5}{x^2 - 2x + 1}$	Original Equation	$y = x^n$ n is even	$y = -x^n$ n is even	$\sin^2 + \cos^2 = 1$
$= \frac{2x^2 - 4x + 5}{(2x - 1)(x + 2)}$	Factor demoniator	$y = x^n$ n is odd	$y = -x^n$ n is odd	$\tan^2 + 1 = \sec^2$
$x = \frac{1}{2}$ or $x = -2$	Impossible			$1 + \cot^2 = \csc^2$

