

Cambridge International AS & A Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

MATHEMATICS 9709/52

Paper 5 Probability & Statistics 1

February/March 2020

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has 12 pages. Blank pages are indicated.

JC20 03_9709_52/RP © UCLES 2020

[Turn over

The 40 members of a club include Ranuf and Saed. All 40 members will travel to a concert.

1

air die is thrown repeatedly until a 1 or a 6 is obtained.	An ordinary fair die is thrown repeatedly until a 1 or a 6 is obtained. (a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
air die is thrown repeatedly until a 1 or a 6 is obtained.	An ordinary fair die is thrown repeatedly until a 1 or a 6 is obtained. (a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
air die is thrown repeatedly until a 1 or a 6 is obtained.	An ordinary fair die is thrown repeatedly until a 1 or a 6 is obtained. (a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
air die is thrown repeatedly until a 1 or a 6 is obtained.	An ordinary fair die is thrown repeatedly until a 1 or a 6 is obtained. (a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
air die is thrown repeatedly until a 1 or a 6 is obtained.	An ordinary fair die is thrown repeatedly until a 1 or a 6 is obtained. (a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
air die is thrown repeatedly until a 1 or a 6 is obtained.	An ordinary fair die is thrown repeatedly until a 1 or a 6 is obtained. (a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
air die is thrown repeatedly until a 1 or a 6 is obtained.	An ordinary fair die is thrown repeatedly until a 1 or a 6 is obtained. (a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
air die is thrown repeatedly until a 1 or a 6 is obtained.	An ordinary fair die is thrown repeatedly until a 1 or a 6 is obtained. (a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
air die is thrown repeatedly until a 1 or a 6 is obtained.	An ordinary fair die is thrown repeatedly until a 1 or a 6 is obtained. (a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
air die is thrown repeatedly until a 1 or a 6 is obtained.	An ordinary fair die is thrown repeatedly until a 1 or a 6 is obtained. (a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
air die is thrown repeatedly until a 1 or a 6 is obtained.	An ordinary fair die is thrown repeatedly until a 1 or a 6 is obtained. (a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
	(a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1
	(a) Find the probability that it takes at least 3 throws but no more than 5 throws to obtain a 1

On another occasion, this die is thrown 3 times. The random variable *X* is the number of times that a 1 or a 6 is obtained.

(b)	Draw up the probability distribution table for X .	[3]
(c)	Find $E(X)$.	[2]

a)	Find the standard deviation of the weights of these apples.
h)	Find the probability that the weight of a randomly chosen apple of this variety differs for
b)	Find the probability that the weight of a randomly chosen apple of this variety differs from mean weight by less than 4 grams.
b)	

4

Richard has 3 blue candles, 2 red candles and 6 green candles. The candles are identical apart from

thei	r colours. He arranges the 11 candles in a line.
(a)	Find the number of different arrangements of the 11 candles if there is a red candle at each end. [2]
(b)	Find the number of different arrangements of the 11 candles if all the blue candles are together and the red candles are not together. [4]

a)	Find the probability that the number of adults in this sample who own a car is less than 6.


A random sample of 120 adults from Greenton is now chosen.

)	Use an approximation to find the probability that more than 75 of them own a car. [5]

6 Box *A* contains 7 red balls and 1 blue ball. Box *B* contains 9 red balls and 5 blue balls. A ball is chosen at random from box *A* and placed in box *B*. A ball is then chosen at random from box *B*. The tree diagram below shows the possibilities for the colours of the balls chosen.

(a) Complete the tree diagram to show the probabilities.

[3]




(b)	Find the probability that the two balls chosen are not the same colour.	[2]
(c)	Find the probability that the ball chosen from box A is blue given that the ball chois blue.	osen from box B [4]

7 Helen measures the lengths of 150 fish of a certain species in a large pond. These lengths, correct to the nearest centimetre, are summarised in the following table.

Length (cm)	0-9	10 – 14	15 – 19	20 – 30
Frequency	15	48	66	21

(a) Draw a cumulative frequency graph to illustrate the data.

[4]



(b)	40% of these fish have a length of d cm or more. Use your graph to estimate the value of d . [2]
The	mean length of these 150 fish is 15.295 cm.
(c)	Calculate an estimate for the variance of the lengths of the fish. [3]

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s must be clearly shown.

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.