

Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/21

Paper 2 Pure Mathematics 2

May/June 2020

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages. Blank pages are indicated.

1 Solve the equation

$$\ln(x+1) - \ln x = 2 \ln 2. \quad [3]$$

[illegible]

2 The polynomial $p(x)$ is defined by

$$p(x) = 6x^3 + ax^2 + 9x + b,$$

where a and b are constants. It is given that $(x - 2)$ and $(2x + 1)$ are factors of $p(x)$.

Find the values of a and b .

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

3 A curve has parametric equations

$$x = e^t - 2e^{-t}, \quad y = 3e^{2t} + 1.$$

Find the equation of the tangent to the curve at the point for which $t = 0$. [5]

This image shows a single page of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

- 4 (a) Sketch, on the same diagram, the graphs of $y = |3x + 2a|$ and $y = |3x - 4a|$, where a is a positive constant.

Give the coordinates of the points where each graph meets the axes. [3]

- (b) Find the coordinates of the point of intersection of the two graphs. [3]

.....

.....

.....

.....

.....

.....

.....

.....


- (c) Deduce the solution of the inequality $|3x + 2a| < |3x - 4a|$. [1]

.....

.....

.....

.....



- [illegible]

- (b) Use the equation in part (a) to show by calculation that the x -coordinate of M lies between 0.59 and 0.60. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Use an iterative formula, based on the equation in part (a), to find the x -coordinate of M correct to 3 significant figures. Give the result of each iteration to 5 significant figures. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6 (a) Prove that

$$\sin 2\theta(\operatorname{cosec} \theta - \sec \theta) \equiv \sqrt{8} \cos(\theta + \frac{1}{4}\pi). \quad [5]$$

[illegible]

(b) Solve the equation

$$\sin 2\theta(\operatorname{cosec} \theta - \sec \theta) = 1$$

for $0 < \theta < \frac{1}{2}\pi$. Give the answer correct to 3 significant figures.

[2]

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find $\int \sin x(\operatorname{cosec} \frac{1}{2}x - \sec \frac{1}{2}x) dx$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 7 (a) Find the quotient when $9x^3 - 6x^2 - 20x + 1$ is divided by $(3x + 2)$, and show that the remainder is 9. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Hence find $\int_1^6 \frac{9x^3 - 6x^2 - 20x + 1}{3x + 2} dx$, giving the answer in the form $a + \ln b$ where a and b are integers. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (c) Find the exact root of the equation $9e^{9y} - 6e^{6y} - 20e^{3y} - 8 = 0$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

[illegible]

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.