

Cambridge International AS & A Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

5 8 5 0 0 5 4 7 6 2

FURTHER MATHEMATICS

9231/41

Paper 4 Further Probability & Statistics

May/June 2020

1 hour 30 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has 12 pages. Blank pages are indicated.

DC (LK) 198480 © UCLES 2020

[Turn over

1 Two randomly selected groups of students, with similar ranges of abilities, take the same examination in different rooms. One group of 140 students takes the examination with background music playing. The other group of 210 students takes the examination in silence. Each student is awarded a grade for their performance in the examination and the numbers from each group gaining each grade are shown in the following table.

		Grade awarded	
	A	В	C
Background music	49	51	40
Silence	93	68	49

Test at the 10% significance level whether grades awarded are independent of whether background music is playing during the examination. [6]

2

The times, in milliseconds, taken by a computer to perform a certain task were recorded on 10 randomly chosen occasions. The times were as follows.

1 .									6.34	6.28	
	ned that the								agt this al	aim	г
a) Call	y out a w	riicoxon	signed-i	ank test	at the 3	% Sigiiii	icance i	ever to to	est this cl	aiii.	[
•••••			•••••			••••••	•••••				
•••••	•••••			•••••			•••••				•••••
•••••	•••••	•••••	••••••	••••••	••••••	••••••	•••••				•••••
											•••••
											•••••
							•••••				
•••••	•••••	•••••								••••••	
											•••••
											•••••
•••••	•••••						•••••				
•••••							•••••				
•••••											
b) State	an unde	rlying as	sumptio	n that is	made w	hen usir	ng a Wil	coxon si	gned-ran	k test.	

 $\mathbf{3}$ The continuous random variable X has probability density function f given by

$$f(x) = \begin{cases} \frac{3}{16} (2 - \sqrt{x}) & 0 \le x < 1, \\ \frac{3}{16\sqrt{x}} & 1 \le x \le 9, \\ 0 & \text{otherwise.} \end{cases}$$

Find $E(X)$.	

The random variable *Y* is such that $Y = \sqrt{X}$. **(b)** Find the probability density function of *Y*. [5]

4	A company has two different machines, X and Y, each of which fills empty cups with coffee. The
	manager is investigating the volumes of coffee, x and y, measured in appropriate units, in the cups filled
	by machines X and Y respectively. She chooses a random sample of 50 cups filled by machine X and a
	random sample of 40 cups filled by machine <i>Y</i> . The volumes are summarised as follows.

$$\sum x = 15.2$$
 $\sum x^2 = 5.1$ $\sum y = 13.4$ $\sum y^2 = 4.8$

The manager claims that there is no difference between the mean volume of coffee in cups filled by machine X and the mean volume of coffee in cups filled by machine Y.

Test the manager's claim at the 10% significance level.	[9]

5

A large number of children are competing in a throwing competition. The distances, in metres, thrown by a random sample of 8 children are as follows.

		19.8	22.1	24.4	21.5	20.8	26.3	23.7	25.0			
(a)	Assuming the population is	hat dista mean dis	nces are tance thr	normal own is	ly distril more tha	outed, te in 22.0 m	st, at the	e 5% sig	gnificance	level, w	hether	the [7]
							•••••	•••••		•••••		
		• • • • • • • • • • • • • • • • • • • •	•••••		•••••	• • • • • • • • • • • • • • • • • • • •	•••••					
			••••••		•••••	•••••	•••••			•••••	•••••	••••
			••••••	•••••	•••••	••••••	••••••	••••••		•••••	•••••	••••
		• • • • • • • • • • • • • • • • • • • •										••••
					•••••	• • • • • • • • • • • • • • • • • • • •		••••••			•••••	••••
	•••••	• • • • • • • • • • • • • • • • • • • •		•••••				••••••			•••••	
	•••••			••••••			•••••				•••••	••••
		•	•••••		•••••	• • • • • • • • • • • • • • • • • • • •	•••••					••••
		• • • • • • • • • • • • • • • • • • • •										
		•••••	••••••		•••••	•••••	•••••	•••••			•••••	••••
			•••••		•••••	•••••	•••••			•••••	•••••	••••

Find a 95% co	onfidence interval for	r the population m	nean distance thrown	

(a)	Find the probability generating function, $G_X(t)$, of X .	[2
ias	sa also tosses two coins. One coin is biased so that the probability of a head is $\frac{2}{3}$ sed so that the probability of a head is p . The probability generating function of ds obtained by Rassa, is $G_{\gamma}(t)$. The coefficient of t in $G_{\gamma}(t)$ is $\frac{7}{12}$.	f <i>Y</i> , the number o
l. X		
D)	Find $G_{\gamma}(t)$.	[3
D)	Find $G_{\gamma}(t)$.	[3
))	Find $G_{\gamma}(t)$.	[3
))	Find $G_{\gamma}(t)$.	[3
0)	Find $G_{\gamma}(t)$.	[3
))		[3
0)	Find $G_{\gamma}(t)$.	[3
0)		[3
0)	Find $G_{\gamma}(t)$.	[3
0)		[3
0)		[3
0)	Find G _y (t).	[3
0)		
D)	Find G _y (f).	[3

The random variable Z is the sum of the number of red balls selected and the number of heads obtained by Rassa.

					 	•••••
					 	•••••
					 	•••••
Use the	probability ge	nerating func	tion of Z to fi	and E(Z).		•••••
Use the	probability ge	nerating func	tion of Z to fi	and E(<i>Z</i>).		
Use the	probability gei	nerating func	tion of Z to fi	nd E(<i>Z</i>).		
Use the	probability gei	nerating func	tion of Z to fi	and E(Z).		
Use the	probability gei	nerating func	tion of Z to fi	nd E(<i>Z</i>).		
Use the	probability gei	nerating func	tion of Z to fi	ind E(Z).		
Use the	probability gei	nerating func	tion of Z to fi	and E(Z).		
Use the	probability gei	nerating func	tion of Z to fi	nd E(Z).		
Use the	probability gei	nerating func	tion of Z to fi	ind E(Z).		
Jse the	probability gen	nerating func	tion of Z to fi	ind E(Z).		

Additional Page

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.				
				•••••
				•••••
				•••••
				•••••
				•••••
				•••••
				•••••

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.