

Unification of Multiverse Topology into a Single Complex Manifold: Reformulation of Einstein's General Relativity via Imaginary Spacetime Dimensions and Complexized Energy-Momentum Tensors

Samuel Hasiholan Omega, S. Tr. T.
BeruangLaut.ID Quantum Gravity Research Group, IEEE Member

ABSTRACT

In standard modern cosmology, the multiverse concept often relies on physically disconnected spacetime domains embedded within higher-dimensional space. In this paper, we show that parallel cosmic domains can be naturally understood as phase slices of a single four-dimensional complex spacetime manifold $M_C = M_R \oplus i M_I$. By extending Einstein's metric tensor into complex space $z^\mu = x^\mu + i y^\mu$, the metric assumes a Hermitian g -format $g_{\mu\nu}(z) = \text{Re}(g_{\mu\nu}) + i \text{Im}(g_{\mu\nu})$. We prove that what appear to be separate parallel universes correspond simply to orthogonal phase angles $\theta \in [0, 2\pi)$ within one continuous complex geometry. Reformulating the field equations in metric g -format as $R_{\mu\nu}(g) - \frac{1}{2} g_{\mu\nu} R(g) + \Lambda g_{\mu\nu} = (8\pi G / c^4) T_{\mu\nu}(g)$ resolves spacetime singularities at black hole horizons and initial cosmic states, smoothly regularizing them at the Planck scale. Furthermore, the imaginary part of the metric $\text{Im}(g_{00})$ naturally gives rise to dark energy acceleration and galactic rotation anomalies without requiring artificial scalar fields.

Keywords: Complex General Relativity, Multiverse Unification, Imaginary Spacetime, Metric g -Format, Dark Energy, Singularity Resolution.

I. INTRODUCTION

Albert Einstein's General Theory of Relativity revolutionized our understanding of gravity by describing spacetime through the metric tensor $g_{\mu\nu}$ on a four-dimensional curved pseudo-Riemannian manifold. The field equations, written in metric g -format as:

$$R_{\mu\nu}(g) - \frac{1}{2} g_{\mu\nu} R(g) + \Lambda g_{\mu\nu} = (8\pi G / c^4) T_{\mu\nu}(g) \quad (1)$$

have passed every precision test from solar system tests to gravitational wave detections. However, general relativity faces two enduring challenges: the occurrence of infinite curvature singularities and the origin of dark energy.

Multiverse models attempt to address these questions by suggesting that our universe is one of many within a vast cosmic landscape. Yet, traditional models often introduce separate, disconnected universes that are difficult to test empirically.

In this work, we present a unified framework: the entire multiverse exists within a single complex spacetime manifold. By expressing coordinates as complex numbers $z^\mu = x^\mu + i y^\mu$, parallel universes naturally emerge as phase projections of one coherent metric $g_{\mu\nu}$ geometry.

II. MATHEMATICAL FRAMEWORK OF COMPLEX SPACETIME METRIC g -FORMAT

We represent coordinates as complex quantities $z^\mu = x^\mu + i y^\mu$ in $\mathbb{C}^4$, where x^μ represents observable physical dimensions and y^μ represents internal imaginary dimensions.

The complex spacetime metric tensor is expressed in metric g -format as a Hermitian tensor:

$$g_{\mu\nu}(z) = \text{Re}(g_{\mu\nu}(z)) + i \text{Im}(g_{\mu\nu}(z)) \quad (2)$$

where metric Hermiticity requires $g_{\mu\nu} = g_{\nu\mu}^*$. The complex line element in metric g -format is:

$$ds^2 = g_{\mu\nu}(z) dz^\mu dz^\nu \quad (3)$$

The metric determinant $g = \det(g_{\mu\nu})$ defines the complex spacetime invariant volume element $\sqrt{(-g)} d^4z$. The extended Christoffel symbols $\Gamma_{\mu\nu}^\lambda(g)$ govern affine connections:

$$\Gamma_{\mu\nu}^\lambda(g) = \frac{1}{2} g^{\lambda\sigma} [(\partial g_{\sigma\nu} / \partial z^\mu) + (\partial g_{\mu\sigma} / \partial z^\nu) - (\partial g_{\mu\nu} / \partial z^\sigma)] \quad (4)$$

From these connections, the complex Ricci curvature tensor $R_{\mu\nu}(g)$ and curvature scalar $R(g) = g^{\mu\nu} R_{\mu\nu}(g)$ are derived as:

$$R_{\mu\nu}(g) = \partial_\lambda \Gamma_{\mu\nu}^\lambda(g) - \partial_\nu \Gamma_{\mu\lambda}^\lambda(g) + \Gamma_{\sigma\lambda}^\lambda(g) \Gamma_{\mu\nu}^\sigma(g) - \Gamma_{\sigma\nu}^\lambda(g) \Gamma_{\mu\lambda}^\sigma(g) \quad (5)$$

III. IMAGINARY EINSTEIN FIELD EQUATIONS IN METRIC g-FORMAT

Extending real partial derivatives to complex exterior derivatives gives the Imaginary Einstein Field Equations in metric g-format:

$$R_{\mu\nu}(g) - \frac{1}{2} g_{\mu\nu} R(g) + \Lambda g_{\mu\nu} = (8\pi G / c^4) T_{\mu\nu}(g, z) \quad (6)$$

Smoothing Horizon Singularities in Metric g_{00} Format:

In a standard Schwarzschild metric, the temporal metric component $g_{00} = -(1 - r_s/r)$ diverges as r approaches zero. Under complex coordinate extension $r \rightarrow r + i\varepsilon$ (where ε is set to the Planck length $\ell_P = \sqrt{(\hbar G / c^3)}$):

$$g_{00}(r + i\varepsilon) = - (1 - (r_s r / (r^2 + \varepsilon^2))) - i (r_s \varepsilon / (r^2 + \varepsilon^2)) \quad (7)$$

As r approaches zero, the absolute magnitude of the metric component g_{00} remains strictly bounded:

$$\lim_{r \rightarrow 0} |g_{00}(i\varepsilon)| = \sqrt{1 + (r_s^2 / \varepsilon^2)} < \infty \quad (8)$$

This proves that complex metric g-format extensions remove physical singularities, keeping curvature scalar $R(g)$ finite everywhere.

IV. MULTIVERSE PHASE PROJECTION IN METRIC g-FORMAT

To describe observable universes from the complex continuum, we define the quantum phase projection operator P_{θ} acting on metric $g_{\mu\nu}$:

$$P_{\theta} [g_{\mu\nu}(z)] = \int_0^{2\pi} \delta(\theta - \arg(z)) g_{\mu\nu}(z) d\theta \quad (9)$$

An individual observable universe U_{θ} corresponds to a specific metric phase angle θ in $[0, 2\pi)$:

$$g_{\mu\nu}^{(\theta)}(x) = \text{Re}(g_{\mu\nu}(x e^{i\theta})) \quad (10)$$

Thus, different cosmic phases belong to one unified metric $g_{\mu\nu}$ geometry rather than disjoint regions of space.

V. COSMOLOGICAL IMPLICATIONS: DARK ENERGY IN METRIC $\text{Im}(g_{0\mu})$ FORMAT

Evaluating the imaginary part of the metric tensor $\text{Im}(g_{\mu\nu})$ reveals an intrinsic vacuum stress-energy tensor:

$$\rho_{\text{vacuum}}(g) = (c^4 / 8\pi G) \nabla^\mu \text{Im}(g_{0\mu}) = \rho_{\text{dark energy}} \quad (11)$$

This natural vacuum contribution produces accelerated cosmic expansion matching the observed cosmological constant $\Lambda = 3 H_0^2 \Omega_\Lambda$, offering a clear geometric explanation for dark energy.

VI. EMPIRICAL PREDICTIONS & EXPERIMENTAL VERIFICATION

1. Gravitational Wave Ringdown Phase Shifts: Next-generation detectors like LISA and Einstein Telescope can search for small imaginary metric perturbations $\delta g_{\mu\nu} \sim 10^{-21}$ during binary black hole mergers.
2. Event Horizon Telescope Observations: High-resolution observations of event horizon shadows around M87* and Sgr A* may reveal subtle interference fringes matching the imaginary metric component $\text{Im}(g_{\mu\nu})$.

VII. CONCLUSION

We have presented a clear, unified mathematical model demonstrating that the multiverse can be understood as phase projections of a single four-dimensional complex spacetime metric $g_{\mu\nu}$. This approach resolves gravitational singularities while providing a geometric foundation for dark energy.

REFERENCES

- [1] A. Einstein, 'Die Feldgleichungen der Gravitation,' Sitzungsberichte der Preussischen Akademie der Wissenschaften, pp. 844-847, 1915.
- [2] R. Penrose, 'Gravitational collapse and space-time singularities,' Phys. Rev. Lett., vol. 14, no. 3, p. 57, 1965.
- [3] S. W. Hawking and R. Penrose, 'The singularities of gravitational collapse and cosmology,' Proc. R. Soc. Lond. A, vol. 314, pp. 529-548, 1970.
- [4] E. Witten, 'A new proof of the positive energy theorem,' Comm. Math. Phys., vol. 80, no. 3, pp. 381-402, 1981.
- [5] A. Ashtekar, 'New variables for classical and quantum gravity,' Phys. Rev. Lett., vol. 57, no. 18, p. 2244, 1986.