

Chapter 1: Random Number Generation, Random Variables Simulation (v.17.1)

The objective of this chapter is to learn to generate/simulate sequences of random variables that come from discrete or continuous distributions. The sequences of numbers we will generate are not truly random, but with sufficient care, the sequences can be made to have the most (and major) properties of random numbers. Most algorithms that will be used to generate sequences of “pseudorandom” numbers are of the form $X_{n+1} = g(X_n)$ for some function g . We will consider only linear functions g .

Starting with a seed X_0 , one can build up the succession of “pseudorandom” numbers by the rule

$$X_{n+1} = (aX_n + b) \text{ modulo } m.$$

That is, the next number X_{n+1} in the sequence is the remainder of the division of a linear form $aX_n + b$ by the number m . Two most popular choices for m are:

(1) m is a prime number, and

(2) m is a power of 2.

When $b = 0$, these generators are referred to as *multiplicative congruential generators* and, in general, as *mixed or linear congruential generators*.

One of the concerns about the above algorithm is the length of the sequence of the pseudorandom numbers generated, before they repeat themselves. Thus it is important to be able to generate a long sequence of numbers that also appear to be random. The following result helps to search such algorithms:

Lemma 1 [B.J.T. Morgan]. Define the sequence $\{X_n\}_{n=0}^{\infty}$ as follows:

X_0 is given and define $X_{n+1} = (aX_n + b) \text{ modulo } m$ for $n > 0$, and let $m = 2^k$, $a = 4c + 1$, b be odd and c a positive integer. Then, the sequence of pseudorandom numbers generated by this algorithm has cycle length of $m = 2^k$.

Lemma 2. If m is a prime number, the *multiplicative congruential generator* $X_{n+1} = aX_n \text{ modulo } m$, and $a \neq 0$, has maximal period $(m - 1)$ if and only if, $a^k \neq 1 \text{ modulo } m$, for all $k = 1, 2, \dots, m - 1$.

Lemma 3. If $m = 2^k$ for $k \geq 3$, and if $a \text{ modulo } 8 = \{3 \text{ or } 5\}$, and X_0 is odd, then the *multiplicative congruential generator* has maximal period $= 2^{k-2}$.

Lemma 4. The *mixed congruential generator*, $X_{n+1} = (aX_n + b) \text{ modulo } m$ has full period m if and only if the following three conditions hold:

- (i) m and b are relatively prime.
- (ii) Each prime factor of m is also a factor of $(a - 1)$.
- (iii) If m is a multiple of 4, then so is $(a - 1)$.

When m is prime, then (ii) together with the assumption that $a < m$ implies that m must be a multiple of $(a - 1)$, which implies $a = 1$. Therefore, for prime m the only full-period generators correspond to $a = 1$. Prime numbers m are desirable for long periods in the case of multiplicative generators, but in the case of mixed congruential generators, only the $X_{n+1} = (X_n + b) \text{ modulo } m$ has maximal period m when m is prime. This covers the popular case: $m = 2^{31} - 1$ (a prime number).

For the generators $X_{n+1} = (aX_n + b) \text{ modulo } m$, where $m = 2^k$ for $k \geq 2$, the condition for full period 2^k requires that b be odd, and $a = 4i + 1$ for some integer i .

One of the most popular generators is the RANDU generator which is a particular case of the generator $X_{n+1} = (aX_n + c) \text{ modulo } m$, where $a = 2^{16} + 3, b = 0, c = 2^{31}$.

Another popular one is the generator of Lewis, Goodman, and Miller (LGM algorithm) that has been used by IBM and seems to do a satisfactory job in generation of random sequences.

The algorithm is given by $X_{n+1} = 7^5 X_n \text{ modulo } (2^{31} - 1)$. It has been the basic generator for the IMSL statistical package.

The most important properties we would like these sequences of pseudorandom numbers to have are the following:

- Be uniformly distributed over a certain set,
- Be mutually independent.

The table below shows some popular random generators:

Table 1: Some Random Number Generators.

m	a	b	Name
$2^{31} - 1$	7^5	0	(LGM) Lewis Goodman, Miller - IBM
2^{31}	$2^{16} + 3$	0	RANDU
2^{35}	5^{13}	0	APPLE
2^{32}	134,775,813	1	Turbo-Pascal
$2^{61} - 1$	$2^{19} - 1$	0	Wu-1997

We will assume that the pseudo-random sequences of numbers generated by the above algorithms satisfy the two properties above:

- they are uniformly distributed over the set $X_n \in \{0, 1, \dots, m - 1\}$,
- they are mutually independent.

These algorithms allow one to use a simple transformation and generate a sequence of random variates that come from *Uniform*[0,1] distribution.

Generation of U[0,1]:

Define $U_n = \frac{X_n}{m}$ or $U_n = \frac{X_n + 1/2}{m}$. Then, $U_n \sim \text{Uniform}[0,1]$ and $\{U_n\}_{n=1}^N \sim i.i.d. U[0,1]$.

Comment: Assume a random variable X has *Uniform*[0,1] distribution, then its Cumulative Distribution Function (CDF) is:

$$F(x) = P(U \leq x) = \begin{cases} 0, & \text{if } x \leq 0 \\ x, & \text{if } 0 < x \leq 1, \\ 1, & \text{if } x > 1 \end{cases}$$

and its probability density function (pdf) is:

$$f(x) = \begin{cases} 0, & \text{if } x < 0 \text{ or } x > 1 \\ 1, & \text{if } 0 \leq x \leq 1 \end{cases}$$

Now we assume we are able to generate sequences of independent and identically distributed random variates from the *Uniform*[0,1] distribution. We next will try to generate sequences of random variates that come from various discretely and continuously distributed families of distributions.

1.1 Discrete Distributions

1.1.1 Bernoulli (p).

We will start with the most basic discrete distribution – the Bernoulli distribution. The Bernoulli (p) distribution is defined as follows: X has Bernoulli(p) distribution if

$$X = \begin{cases} 1, & \text{with prob. } p \\ 0, & \text{with prob. } 1 - p \end{cases}$$

The following steps will result in generation of random variates with Bernoulli(p) distribution:

STEP 1: Generate $U \sim U[0,1]$

STEP 2: if $U < p$ then set $X = 1$, else, set $X = 0$.

Then X will have the above distribution because $P(X = 1) = P(U < p) = p$ and $P(X = 0) = 1 - p$.

1.1.2 General Discrete Distributions

Using the same method as in the Bernoulli case, we will generate a general discretely distributed random variates with finitely many values:

$$X = \begin{cases} x_1, & \text{with prob. } p_1 \\ x_2, & \text{with prob. } p_2 \\ x_3, & \text{with prob. } p_3 \\ \vdots & \\ x_m, & \text{with prob. } p_m \end{cases}$$

where $\sum_{i=1}^m p_j = 1$.

The following steps will result in generation of random variates with the above distribution:

STEP 1: Generate $U \sim U[0,1]$,

STEP 2: if $U < p_1$ then set $X = x_1$, else, if $U < p_1 + p_2$ then set $X = x_2$,
else, ... , if $U < p_1 + p_2 + \dots + p_{m-1}$ then set $X = x_{m-1}$, else, set $X = x_m$.

Then X will have the above distribution because

$$P(X = x_j) = P(p_1 + \dots + p_{j-1} \leq U \leq p_1 + \dots + p_{j-1} + p_j) = p_j,$$

for any $j = 1, \dots, m$, and $p_0 = 0$.

1.1.3 Binomial(n, p)

We say that X has Binomial(n, p) distribution if

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k} \text{ for any } k = 0, 1, \dots, n$$

The following steps will result in generation of random variates with Binomial (n, p) distribution:

Method 1: By recursion.

$$\text{Notice that } P(X = k + 1) = \frac{n-k}{k+1} \frac{p}{1-p} P(X = k).$$

$$\text{Define } z = \frac{p}{1-p}, \quad y = (1 - p)^n$$

STEP 1: Generate $U \sim U[0,1]$

STEP 2: $k = 0, x = y$

STEP 3: if $U < x$ then set $X = k$ and exit, else

STEP 4: set $y = \frac{n-k}{k+1}zy, x = x + y, k = k + 1$

STEP 5: Repeat from STEP 3.

The resulting X will have the desired distribution by its construction.

Method 2: We will use the Bernoulli-Decomposition of a Binomial(n, p).

Lemma: Assume $X \sim \text{Binomial}(n, p)$ then it has the same distribution as $Y_1 + Y_2 + \dots + Y_n$ for any sequence $\{Y_i\}_{i=1}^n$ of i.i.d. Bernoulli(p).

The following steps will result in generation of random variates with Binomial (n, p) distribution using the above result:

STEP 1: Generate a sequence $\{U_i\}_{i=1}^n \sim \text{i.i.d. } U[0,1]$,

STEP 2: Using the numbers in STEP 1 and the Bernoulli(p) –generation method, generate a sequence of $\{Y_i\}_{i=1}^n \sim \text{i.i.d. Bernoulli}(p)$,

STEP 3: Define $X = Y_1 + Y_2 + \dots + Y_n$.

Then X will have $\text{Binomial}(n, p)$ distribution by its construction.

1.1.4 Poisson (γ)

X has Poisson(γ) distribution if its distribution is given by $P(X = k) = \frac{e^{-\gamma} \gamma^k}{k!}$ for $k = 0, 1, \dots, n, \dots$

The following steps will result in generation of random variates with Poisson(γ) distribution:

Method: By recursion.

Notice that $P(X = k + 1) = \frac{\gamma}{k+1} P(X = k)$,

Define $z = e^{-\gamma}$.

STEP 1: Generate $U \sim U[0,1]$

STEP 2: $k = 0, x = z$

STEP 3: if $U < x$ then set $X = k$ and exit, else

STEP 4: Set $z = \frac{\gamma}{k+1} z, x = x + z, k = k + 1$

STEP 5: Repeat from STEP 3.

Then X will have $\text{Poisson}(\gamma)$ distribution by its construction.

1.2 Continuous Distributions

To generate non-uniform continuous distributions random number generator from the uniform distribution on the unit interval $U(0,1)$ is used along with the following result:

Theorem. Let X be a continuous random variable with cumulative distribution function $F(\cdot)$ [that is, $F(x) = P(X \leq x)$]. Assume $F(\cdot)$ is an increasing and continuous function. Let U be a $\text{Uniform}[0,1]$ -distributed random variable. Consider the random variable $Y = F^{-1}(U)$. Then Y has the same distribution as X .

Proof:

$$P(Y \leq y) = P(F^{-1}(U) \leq y) = P(F(F^{-1}(U)) \leq F(y)) = P(U \leq F(y)) = F(y) = P(X \leq y).$$

This proves the theorem.

Comment: If $F(\cdot)$ is a non-decreasing and (potentially) discontinuous function, then the inverse of F does not exist in the traditional sense. We will define the pseudo-inverse function of F as follows:

$$F^{-1}(y) = \min\{x: F(x) \geq y\}$$

The above results then holds true for this function F .

1.2.1 Exponential(γ)

Definition: X has Exponential (γ) distribution, $X \sim \text{Exp}(\gamma)$, if the CDF of X is given by

$$F(y) = P(X \leq y) = 1 - e^{-\frac{y}{\gamma}} \text{ for } y \geq 0$$

The exponential distribution is common in many applications of probability theory. Due to its relationship to Poisson distribution, it is used to model the time between two events, the occurrence of which follows a Poisson process. An example of such events could be defaults of securities in a pool.

If $X \sim \text{Exp}(\gamma)$, then the CDF of the random variable X is $F(y) = P(X \leq y) = 1 - e^{-\frac{y}{\gamma}}$ for $y \geq 0$. Using the inverse transformation method, we get the inverse and can generate a variate Y that comes from exponential distribution:

$$Y = -\gamma \ln(1 - U) \sim X \sim \text{Exp}(\gamma)$$

Comment: Notice that $Y = -\gamma \ln(U) \sim \text{Exp}(\gamma)$ since U and $1 - U \sim U[0,1]$.

1.2.2 Gamma(n, γ).

To generate a random variable X that has $\Gamma(n, \gamma)$ distribution, we will use the fact that X can be written as (has the same distribution as) a sum of n independent Exponentially distributed random variables:

$$X = Y_1 + Y_2 + \cdots + Y_n$$

where $Y_1 \sim \text{Exp}(\gamma)$. Since we discussed the generation of exponentially distributed random variables, the rest is a simple application of that method.

1.2.3 Logistic(a, b)

Definition: $X \sim \text{Logistic}(a, b)$ if $P(X \leq t) = \frac{1}{1 + e^{-\frac{t-a}{b}}}$.

To generate Logistic distribution, we would use the inverse-transformation method and define

$$Y = a + b \ln \left(\frac{U}{1-U} \right) \sim X \sim \text{Logistic}(a, b).$$

1.2.4 Student's t

Definition: X has t distribution with $n - 1$ degrees of freedom, $X \sim t_{n-1}$, if its density function is given by

$$f(x) = \frac{\Gamma\left(\frac{n+1}{2}\right)}{\sqrt{n\pi} \Gamma\left(\frac{n}{2}\right)} \left(1 + \frac{x^2}{n}\right)^{-(n+1)/2}$$

Lemma: Assume $\{X_i\}_{i=1}^n$ are i.i.d. $N[0,1]$ random sample and $\bar{X} = \frac{\sum_{i=1}^n X_i}{n}$ is the Sample Mean, and $S = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2}$ is the Sample Standard Deviation. Then $t = \frac{\bar{X}}{S/\sqrt{n}}$ has Student's t distribution with $(n - 1) -$ degrees of freedom.

Problem: Suppose that $\{U_i\}_{i=1}^n \sim \text{i.i.d. } U[0,1]$ and let $S_n = U_1 + U_2 + \dots + U_n$. Define $N = \min\{k: S_k > 1\}$

- Show that $P(S_k \leq x) = \frac{x^k}{k!}$ for $0 \leq x \leq 1$.
- Show that $E(N) = e$
- Use part (b) to estimate e by simulation (using the Law of Large Numbers).
- How can you generate approximate standard normally distributed random variables using S_n ? (Use the Central Limit Theorem)

The following random variables with given distributions can be generated by using the inverse transformation algorithm:

- Rayleigh(b) distribution with CDF $F(x) = 1 - e^{-2x(x-b)}$ for $x \geq b$.

b. *Pereto*(a, b) distribution with CDF $F(x) = 1 - \left(\frac{b}{x}\right)^a$ for $x \geq b > 0$.

c. *Arcsine* distribution with CDF $F(x) = \frac{2}{\pi} \arcsin(\sqrt{x})$ for $0 \leq x \leq 1$.

1.2.5 Normal Distribution

To generate normally distributed random variables one can not use the inverse transformation method as there is no closed-form expression for the inverse CDF function in that case. We will consider two methods for generation of normally distributed random variables. In both cases, the outcome of each step is a pair of independent, standard normally-distributed random variables.

Method 1. Box-Muller Method

Theorem: Assume $\{U_i\}_{i=1}^2 \sim i.i.d. U[0,1]$. Define

$$\begin{cases} Z_1 = \sqrt{-2\ln(U_1)} \cos(2\pi U_2) \\ Z_2 = \sqrt{-2\ln(U_1)} \sin(2\pi U_2) \end{cases}.$$

Then Z_1 and Z_2 are $i.i.d. N(0,1)$.

For the proof of the Theorem we need the following

Lemma:

(a) Let $X, Y \sim i.i.d. N(0,1)$. Define the polar coordinates of the couple (X, Y) : $\begin{cases} R^2 = X^2 + Y^2 \\ \theta = \arctan \frac{Y}{X} \end{cases}$.

Then $R^2 \sim \text{Exp}(2)$, and $\theta \sim U[0, 2\pi]$ and R^2 is independent of θ .

(b) If $R^2 \sim \text{Exp}(2)$, and $\theta \sim U[0, 2\pi]$ and R^2 is independent of θ , then if $X = R \sin \theta$, $Y = R \cos \theta$, then $X, Y \sim i.i.d. N(0,1)$.

Proof of Lemma:

(a) Use the fact that $f_{R^2, \theta}(R^2, \theta) = |J| f_{X, Y}(x(R^2, \theta), y(R^2, \theta))$, where the Jacobian is

$$|J| = \left| \begin{array}{cc} \frac{1}{2\sqrt{R^2}} \cos \theta & -R \sin \theta \\ \frac{1}{2\sqrt{R^2}} \sin \theta & -R \cos \theta \end{array} \right| = \frac{1}{2}. \text{ The joint density of } (R^2, \theta) \text{ can be written as}$$

$$f_{R^2, \theta}(d, \alpha) = \frac{1}{2} \frac{1}{2\pi} e^{-d/2} = \left(\frac{1}{2} e^{-\frac{d}{2}} \right) \left(\frac{1}{2\pi} \right) \text{ for } 0 < d < \infty, \text{ and } 0 < \alpha < 2\pi.$$

This implies that $R^2 \sim \text{Exp}(2)$, and $\theta \sim U[0, 2\pi]$ and that R^2 is independent of θ .

(b) The proof is a replication of (a), in a reversed order.

Proof of Theorem: Let $U_1, U_2 \sim i.i.d. U[0, 1]$. Define: $\begin{cases} R^2 = -2\ln U_1 \\ \theta = 2\pi U_2 \end{cases}$. Then it is easy to see that $R^2 \sim \text{Exp}(2)$, and $\theta \sim U[0, 2\pi]$ and that R^2 is independent of θ .

From the Lemma above it follows that if we define $Z_1 = R \sin \theta, Z_2 = R \cos \theta$, then

$$\begin{cases} Z_1 = \sqrt{-2\ln(U_1)} \cos(2\pi U_2) \\ Z_2 = \sqrt{-2\ln(U_1)} \sin(2\pi U_2) \end{cases} \text{ and } Z_1 \text{ and } Z_2 \text{ are } i.i.d. N(0, 1).$$

Method 2. Polar-Marsaglia Method

Theorem: Assume $\{U_i\}_{i=1}^2 \sim i.i.d. U[0, 1]$. Define $V_1 = 2U_1 - 1, V_2 = 2U_2 - 1, W = V_1^2 + V_2^2$, and if $W \leq 1$, define

$$\begin{cases} Z_1 = V_1 \sqrt{\frac{-2 \ln W}{W}} \\ Z_2 = V_2 \sqrt{\frac{-2 \ln W}{W}} \end{cases}$$

Then Z_1 and Z_2 are $i.i.d. N(0, 1)$.

Proof. $\{U_i\}_{i=1}^2 \sim i.i.d. U[0, 1]$, then $(V_1, V_2) \sim U[-1, 1] \times [-1, 1]$. We will choose those couples (U_1, U_2) that are in the unit circle: $W \leq 1$. Then $(U_1, U_2) \sim U(\text{in unit circle})$. The polar

coordinates of (V_1, V_2) are (R, θ) , and $R \sim U[0,1]$, $\theta \sim U[0,2\pi]$, and R is independent of θ .
Since

$$\begin{cases} \sin\theta = \frac{V_2}{R} = \frac{V_2}{\sqrt{V_1^2 + V_2^2}} \\ \cos\theta = \frac{V_1}{R} = \frac{V_1}{\sqrt{V_1^2 + V_2^2}} \end{cases}$$

Then, it follows from the results of the Box-Muller method that

$$\begin{cases} Z_1 = (-2\ln W)^{1/2} \frac{V_1}{\sqrt{V_1^2 + V_2^2}} \\ Z_2 = (-2\ln W)^{1/2} \frac{V_2}{\sqrt{V_1^2 + V_2^2}} \end{cases}$$

are *i. i. d.* $N(0,1)$ because $(U_1, U_2) \sim U(\text{in the unit circle})$.

Comment: In this method, not every pair (U_1, U_2) of uniforms is utilized in generation of standard normals. We only use those pairs that are inside the unit circle (with its center at the origin and radius 1). That is, the proportion of the pairs utilized will be the probability of a random pair of uniforms $(V_1, V_2) \sim U[-1,1] \times [-1,1]$ falling inside the unit circle: $V_1^2 + V_2^2 \leq 1$. This probability is $\pi/4$.

A few important results:

Law of Large Numbers (LLN)

Suppose $\{X_i\}_{i=1}^n \sim i. i. d.$ sample and $E(X_i) = \mu$, $Var(X_i) = \sigma^2 < \infty$.

Define $S_n = \frac{X_1 + X_2 + \dots + X_n}{n}$. Then, $\lim_{n \rightarrow \infty} E(S_n - \mu)^2 = 0$, that is $S_n \approx \mu$ for large enough n .

Central Limit Theorem (CLT)

Suppose $\{X_i\}_{i=1}^n \sim i.i.d.$ sample and $E(X_i) = \mu$, $Var(X_i) = \sigma^2$. Then

$Z_n = \frac{S_n - \mu}{\sigma\sqrt{n}} \xrightarrow{n \rightarrow \infty} Z \sim N(0,1)$. The convergence is in distribution.

1.3 Acceptance-Rejection Method

Suppose we want to generate random variates that come from a distribution with density function $f(x)$ and that it is difficult or impossible to invert the corresponding CDF. That is, the inverse transform method is not applicable. Assume there exists a function $g(x)$ such that

$$g(x) \geq f(x) \text{ for any } x \text{ for which } f(x) \neq 0.$$

Assume $\int_{-\infty}^{\infty} g(x) < \infty$. Define $h(x) = \frac{g(x)}{\int_{-\infty}^{\infty} g(x)}$. Then $h(\cdot)$ is a density function.

The idea behind the acceptance-rejection method is to select a function g in such a way that it is relatively easy to simulate a random variate that has a density of h . Then, the following steps will result in a random variate X that has a distribution with density function f :

STEP 1. Generate Y from the $h(\cdot)$ distribution,

STEP 2. Generate $U \sim U[0,1]$ that is independent of Y ,

STEP 3. If $U \leq \frac{f(Y)}{g(Y)}$ then ACCEPT Y and set $X = Y$ and exit,

else, if $U > \frac{f(Y)}{g(Y)}$ REJECT Y and go to the next STEP,

STEP 4. Repeat Steps 1-3.

Comment: If the support of the function f is a finite interval, say $[a, b]$, then we can take

$$g(x) = \begin{cases} 0, & \text{if } x < a, \text{ or } x > b \\ \max_{x \in [a, b]} f(x), & \text{if } a \leq x \leq b \end{cases} \quad \text{and thus,}$$

$$h(x) = \begin{cases} 0, & \text{if } x < a, \text{ or } x > b \\ \frac{1}{(b-a)}, & \text{if } a < x \leq b. \end{cases}$$

Proof of the Acceptance-Rejection Algorithm for continuous random variables:

It is necessary to show that the conditional distribution of Y given that $U \leq \frac{f(Y)}{g(Y)}$ is F . Using

Bayes Theorem it follows that

$$P\left(Y \leq y \mid U \leq \frac{f(Y)}{g(Y)}\right) = \frac{P\left(U \leq \frac{f(Y)}{g(Y)} \mid Y \leq y\right) P(Y \leq y)}{P\left(U \leq \frac{f(Y)}{g(Y)}\right)}$$

Notice that

$$P\left(U \leq \frac{f(Y)}{g(Y)}\right) = \int_{-\infty}^{+\infty} P\left(U \leq \frac{f(Y)}{g(Y)} \mid Y = y\right) h(y) dy = \int_{-\infty}^{+\infty} \frac{f(y)}{g(y)} h(y) dy = \frac{1}{\int_{-\infty}^{+\infty} g(x) dx}.$$

For notational convenience define $c = \int_{-\infty}^{+\infty} g(x) dx$. Observe further that

$$\begin{aligned} P\left(U \leq \frac{f(Y)}{g(Y)} \mid Y \leq y\right) &= \frac{P\left(U \leq \frac{f(Y)}{g(Y)}, Y \leq y\right)}{P(Y \leq y)} \\ &= \frac{1}{H(y)} \int_{-\infty}^y P\left(U \leq \frac{f(Y)}{g(Y)} \mid Y = w\right) h(w) dw = \frac{1}{H(y)} \int_{-\infty}^y \frac{f(w)}{g(w)} h(w) dw = \frac{F(y)}{cH(y)}. \end{aligned}$$

Therefore it follows that

$$P\left(Y \leq y \mid U \leq \frac{f(Y)}{g(Y)}\right) = \frac{\frac{F(y)}{cH(y)} H(y)}{\frac{1}{c}} = F(y).$$

This completes the proof.

Remarks: In the Acceptance-Rejection algorithm we accept only a fraction of all generated variates, therefore one needs to evaluate the efficiency of the algorithm. The answers to the following questions will shed light on the efficiency of the algorithm:

- (a) What is the probability of acceptance of this algorithm?
- (b) Define X to be the time of the first “success”, defined as accepting the generated number. What is $E(X)$?
- (c) How to choose $g(x)$ to minimize the computational cost of the algorithm?

Suggested Exercises:

1. Use the Acceptance-Rejection method to generate $N(0, 1)$ -distributed random variables. Use Double-Exponential distribution for g : $g(x) = \frac{1}{2}e^{-|x|}$. Notice that here $\frac{f(x)}{g(x)} < c = 1.32$.
2. The answers to the following questions will shed light on the efficiency of the Acceptance-Rejection algorithm. Define N to be the time of the first “success” – accepting the generated number.
 - (a) What is the probability distribution of N ?
 - (b) Compute EN and show that $EN = c$.
 - (c) How to choose $g(x)$ to minimize the computational cost of the algorithm? Ideally, we would like to choose g ‘close’ to f to minimize the computational load. In such cases, c will be close to 1 and we will need to perform fewer iterations. However, there is a tradeoff: g being ‘close’ to f helps to have fewer iterations, but if it is difficult to generate from f , then it must be difficult to generate from g as well.

Examples:

1. Generate $Beta(a, b)$ random variable using the Acceptance-Rejection Method.

The density function of $Beta(a, b)$ is given by:

$$f(x) = \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} x^{a-1}(1-x)^{b-1} \text{ for } 0 \leq x \leq 1, \text{ where } \Gamma(a) = \int_0^{\infty} x^{a-1} e^{-x} dx$$

Use a specific example for demonstration: $Beta(4,3)$. Then, the density function is

$f(x) = 60x^3(1-x)^2$ for $0 \leq x \leq 1$. It is easy to see that $f(x) \leq 4$ for $0 \leq x \leq 1$ (verify).

We will take

$$g(x) = \begin{cases} 4, & \text{for } 0 \leq x \leq 1 \\ 0, & \text{else} \end{cases}$$

and follow the algorithm.

Comments: 1. Notice that the function $g(x) = \begin{cases} 0, & \text{else} \\ 3, & \text{for } 0 \leq x \leq 1 \end{cases}$ would also work in this case. So, the obvious question is: is there an “optimal” choice of g ?

The answer is: the optimal g (among constants) is the following function:

$$g(x) = \begin{cases} \max \frac{f(x)}{g(x)}, & \text{for } 0 \leq x \leq 1 \\ 0, & \text{else} \end{cases}$$

The reasoning of this is that, the average number of trials in the algorithm until acceptance is the number that we choose in g (3 or 4 above), therefore $\max f(x)$ would be the best choice to minimize that number.

1.4 Evaluating Integrals

Suppose we want to estimate the following integral which cannot be computed explicitly:

$\int_0^1 f(x) dx$. It is easy to see that the integral can be written in the following form $\int_0^1 f(x) dx = Ef(U)$, where $U \sim U[0,1]$. The latter expectation can be estimated by using the Law of Large Numbers (LLN): For a sequence of $\{U_i\}_{i=1}^n \sim i.i.d. U[0,1]$

$$Ef(U) \approx \frac{f(U_1) + f(U_2) + \dots + f(U_n)}{n}$$

for large enough n . This is the main idea behind Monte-Carlo simulation.

Comments:

- (1) If the integral is not on the $[0,1]$ interval, a simple rescaling would suffice to convert it to an integral on the $[0,1]$ interval:

$$\int_a^b f(x)dx = (b-a) \int_0^1 f(a + (b-a)t)dt$$

$$\int_0^\infty f(x)dx = \int_0^1 \frac{f\left(\frac{1}{t}-1\right)}{t^2} dt$$

- (2) Multi-dimensional integrals are computed similarly:

$$\int_0^1 \int_0^1 f(x,y)dx dy = Ef(U_1, U_2)$$

where $U_1, U_2 \sim U[0,1]$ and are independent. LLN implies that

$$Ef(U_1, U_2) \approx \frac{f(U_1^1, U_2^1) + f(U_1^2, U_2^2) + \dots + f(U_1^n, U_2^n)}{n}$$

for large enough n .

- (3) The estimator $\hat{\theta}$ is an unbiased estimator for θ because $E\hat{\theta} = \theta$.

The estimator $\hat{\theta}$ is a consistent estimator for θ because $\hat{\theta} \rightarrow \theta$ almost surely as $n \rightarrow \infty$.

1.5 The Kolmogorov-Smirnov Test

One simple test to determine whether a random number generator is yielding close-to-random realizations from a distribution with CDF $F(x)$ is the Kolmogorov-Smirnov test (KS test).

Kolmogorov and Smirnov developed a goodness of fit test for continuous data to determine if a sample comes from a given distribution. Currently the method continues to be one of the most widely used goodness of fit tests because of its simplicity and because it is based on the

empirical distribution function (EDF), which converges uniformly to the population cumulative distribution function (CDF) with probability one (this is based on the Glivenko-Cantelli Theorem). Even though many of goodness of fit tests have been developed in recent years with higher statistical power than the KS test, the KS test remains popular because it is simple and intuitive, comparing the EDF to the CDF.

Assume we have a sample of size of n . (Simulated by using the random number generators considered in this book). Order the sample the following way: $x_{(1)} \leq x_{(2)} \leq \dots x_{(n-1)} \leq x_{(n)}$.

Define the sample distribution function S_n as follows

$$S_n(x) = \begin{cases} 0 & \text{for } x < x_{(1)} \\ \frac{r}{n} & \text{for } x_{(r)} \leq x < x_{(r+1)} \\ 1 & \text{for } x_{(n)} \leq x \end{cases}$$

Then, for large enough n ($n > 40$), it can be shown that IF x 's are indeed from a distribution with CDF F , then, with probability $1 - \alpha$

$$|S_n(x) - F(x)| < \frac{K_\alpha}{\sqrt{n}}$$

The table below shows the critical values of the KS test.

Table 2: *Approximate Critical Values of the Kolmogorov-Smirnov Test for Large n .*

α	0.20	0.10	0.05	0.02	0.01	0.005	0.001
K_α	1.07	1.22	1.36	1.52	1.63	1.73	1.95

To test for the uniformity and/or randomness of a sequence of generated random variates, one can use one of the following tests:

1. Runs Test – tests if a given sequence has iid property;
2. Swerial Correlation Test – tests for independence;
3. Chi-Squared Test – tests for uniformity of a sequence;
4. Spectral Test – tests for randomness.

1.6 Exercises

1. Use the Random Number generators discussed in the class to do the following:
 - (a) Generate 10,000 Uniformly distributed random numbers on $[0,1]$ and plot the histogram of them using LGM method;
 - (b) Now use built-in functions of whatever software you use to do the same thing as in (a).
 - (c) Compare the histograms of the above random number sequences (in (a) and (b)) – use any method of comparison you like.
2. Use the numbers of part (a) of question 1 to do the following:
 - (a) Generate 10,000 random numbers with the following distribution;

$$X = \begin{cases} -1 & \text{with prob. } 0.3 \\ 0 & \text{with prob. } 0.5 \\ 1 & \text{with prob. } 0.2 \end{cases}$$

- (b) Draw the histogram and the empirical distribution function by using the 10,000 numbers generated above in part (a).
3. Use the idea of part (a) of question 1 to do the following:
 - (a) Generate 1,000 random numbers with Binomial distribution with $n = 44$ and $p = 0.64$.
(*Hint*: A random variable with Binomial distribution (n, p) is a sum of n Bernoulli (p) distributed random variables, so you will need to generate 44,000 Uniformly distributed random numbers, to start with).
 - (b) Draw the histogram. Compute the probability that X , with Binomial $(44, 0.64)$ distribution, is at least 40.
 - (c) Use any statistics textbook for the exact number for the above probability and compare them.
4. Use the numbers of part (a) of question 1 to do the following:
 - (a) Generate 10,000 Exponentially distributed random numbers with parameter $\lambda = 1.5$.
 - (b) Draw the histogram by using the 10,000 numbers of part (a).

5. Use the idea of part (a) of question 1 to do the following:
 - (a) Generate 5,000 Uniformly distributed random numbers on $[0,1]$.
 - (b) Generate 5,000 Normally distributed random numbers with mean 0 and variance 1, by **Box-Muller** Method.
 - (c) Draw the histogram by using the 5,000 numbers of part (b).
 - (d) Now use the **Polar-Marsaglia** method to do the same as in (b). here you will not have the same number of RVs)
 - (e) Draw the histogram by using the numbers you got in (d).
 - (f) Now compare the **times** it takes the computer to generate 5,000 normally distributed random numbers by the two methods. Which one is more efficient?

If you do not see a clear difference, you need to increase the number of generated realizations of random variables.
6. (a) Use the density formula for the standard normal distribution to construct the density curve. Plot it from -4 to 4 by 0.0005 step size.
- (b) Compare it to the ones obtained in question 5. Do you see any differences?
7. (a) Now use the built-in function (of the software you are using : Matlab, C/C++, VBA, etc.) to generate 5,000 Normally distributed random numbers with mean 0 and variance 1.
8. (b) Compare the histogram of these to the other three cases. Are there any biases? (You may compute the means of your sequence of numbers and compare them to 0. Ideally they must be equal to 0. Are they?)
9. Generate 10,000 of each of the following distributions using Excel and draw their histograms: Uniform $[0,1]$, Bernoulli (0.65), Normal(0,1), Exponential(2).
10. How many uniform random variables do you have to generate on average to get 1000 normally distributed random variables via the Polar-Marsaglia method?
11. Estimate the following by using random number generation techniques:
 - a. $P(U^2 + V^2 \leq 0.8)$ for U, V being iid $U[-1/2, 3/2]$
 - b. $E(U^2 V^2 | U \leq 0.8)$ for U, V being iid $U[-1/2, 3/2]$

12. Which algorithm is faster: Box Muller or Polar Marsaglia? Discuss.
13. Let Z be a standard normal random variable. We want to generate random variables which are distributed as the absolute value of Z using the Acceptance-Rejection algorithm.
- Find the probability density function f of $|Z|$.
 - Find a function $g(x)$ such that $g(x) \geq f(x)$ for any x for which $f(x) \neq 0$. (Hint: use exponential density).
 - Formulate the Acceptance-Rejection algorithm to generate random variables distributed as $|Z|$.
 - How can you extend this algorithm to generate standard normal random variables?
14. Suppose you have generated n independent $U[0,1]$ random variables U_1, U_2, \dots . Find an algorithm for generating a random variable X , where:
- X has the following probability density function:

$$f(x) = \frac{1}{\sqrt{2\pi}2x} \exp\left(-\frac{(\ln(x) - 2)^2}{8}\right)$$

- X has the following probability density function:

$$f(x) = \frac{1}{96} x^3 \exp\left(-\frac{x}{2}\right) \text{ for } x \geq 0.$$

15. Evaluate the following integral by simulation:

- $\int_0^1 (1 - x^2)^{\frac{3}{2}} dx$

- $\int_0^1 \int_0^1 (1 - x^2)^{3/2} e^{(x+y)^2/2} dx dy$

16. Which is larger – the Delta of a 3 month or the 6-month Call options on a non-dividend-paying stock?
17. Which is larger – the Gamma of a 3 month or the 6-month At-The-Money Call options on a non-dividend-paying stock?