Acoustic Biophysics, Neurocognitive Architecture, and Generative Computation in Music and Sound: An Integrative Transdisciplinary Synthesis

SAMUELSON G

Independent Researcher

Corresponding author: SAMUELSON G

Email: gsamuelsonguna@gmail.com

ORCID: 0009-0005-8744-8178

Department of Systematic Musicology, Auditory Neuroscience, and Acoustic Engineering

Abstract

Sound is a mechanical wave phenomenon governed by continuum mechanics, whereas music is an emergent, rule-based neurocognitive and sociocultural construct that exploits the biophysical constraints of the mammalian auditory pathway. Despite substantial progress within isolated disciplines—ranging from cochlear micromechanics and psychoacoustics to neuroimaging and deep generative modeling—a unified theoretical framework bridging physical wave transduction, cortical predictive coding, clinical neurorehabilitation, and discrete neural audio tokenization remains elusive. This research paper synthesizes empirical and theoretical developments across seven intersecting domains of music and sound science. First, the physical acoustics of wave propagation, binaural head-related filtering, and the active biomechanics of the human cochlea are formalized through the Greenwood frequency-position function and morphometric analyses of radial-to-tangential spiral ganglion innervation. Second, psychoacoustic models of critical bandwidths, sensory roughness, and timbre-dependent tuning are evaluated alongside boundary acoustic phenomena, including infrasonic adaptation and the hypersonic effect. Third, cross-cultural ethnomusicological evidence from global corpora is integrated with archaeoacoustic reconstructions of prehistoric ritual spaces. Fourth, the neurocognitive architecture of musical syntax and reward is examined, reconciling domain-specific representations with domain-general prefrontal control under the Shared Syntactic Integration Resource Hypothesis (SSIRH) and mapping the anatomical dissociation between dorsal striatal anticipation and ventral striatal consummation in dopaminergic transmission. Fifth, clinical applications in Neurologic Music Therapy (NMT) demonstrate how rhythmic

auditory cueing recruits cerebello-thalamo-cortical networks to bypass basal ganglia pathology in movement and language disorders. Finally, the mathematical foundations of contemporary generative audio artificial intelligence—specifically Residual Vector Quantization (RVQ), Finite Scalar Quantization (FSQ), and autoregressive codebook interleaving—are analyzed as computational mirrors of hierarchical human auditory processing.

Keywords: Physical Acoustics; Greenwood Function; Psychoacoustics; Shared Syntactic Integration Resource Hypothesis; Striatal Dopamine; Neurologic Music Therapy; Neural Audio Codecs; Residual Vector Quantization.

1. Epistemological and Evolutionary Foundations of Sound and Music

The scientific investigation of sound and music requires a rigorous distinction between the physical stimulus—longitudinal pressure oscillations propagating through an elastic medium—and the perceptual, cognitive, and cultural architectures that organize those oscillations into meaningful structures. Early musicological frameworks defined music broadly as "humanly organized sound," a characterization that captures intentionality but under-specifies the neurobiological and mathematical constraints governing musical systems. Contemporary systematic musicology and cognitive neuroscience conceptualize music as a complex, time-ordered, and rule-based sensory language that operates across four simultaneous dimensions: a high-precision motor skill, a syntax-governed communicative phenomenon, a complex multidimensional auditory percept, and an abstract set-theoretic mathematical entity.

From an evolutionary perspective, the capacity for musicality is deeply embedded in hominin biology, with comparative and anthropological evidence indicating that vocal melodic expression and rhythmic entrainment preceded the emergence of grammatically arranged propositional speech. Rather than functioning either as an incidental evolutionary byproduct or as a narrow, single-purpose biological adaptation, music operates as a transformative technology of the mind—an invention enabled by human neuroplasticity that reorganizes cognitive control, emotional regulation, and social cohesion across generations.

Empirical validation of this evolutionary universality emerges from large-scale cross-cultural corpora, notably the Natural History of Song project, which evaluated ethnographic records and field recordings across global societies using machine summaries, expert musicological transcriptions, and listener ratings mapped against more than 750 Outline of Cultural Materials (OCM) behavioral identifiers. Statistical comparisons between song-associated ethnographic passages and simulated null distributions demonstrate that music is systematically coupled to specific behavioral contexts—specifically infant care, healing, mourning, religiosity, formality, and warfare—and that the acoustic features of a song reliably predict its primary behavioral function across unrelated cultures. While rhythmic and melodic complexity vary

considerably across cultural lineages, tonal organization and discrete pitch anchoring represent near-universal structural invariants of human song.

The material record of human engagement with sound is further illuminated by archaeoacoustics, a multidisciplinary field examining paleolithic and ancient sound-producing artifacts—such as bone flutes, shell trumpets, and garment-attached rattling pendants crafted from teeth, mollusks, or stones—alongside the architectural acoustics of historical and prehistoric environments. Long before the eighteenth-century formalization of wave mechanics or the Roman architect Vitruvius's treatise *De Architectura* on the resonance of semienclosed theaters, ancient societies engineered enclosed, semienclosed, and open-air spaces to manipulate auditory perception. At the Andean ceremonial center of Chavín de Huántar, acoustic testing of the subterranean zig-zag galleries demonstrates that early, asymmetric sound reflections and resonant filtering amplified spatial disorientation. When ritual practitioners sounded marine shell trumpets (*pututus*), chanted, or sang within these labyrinths, the architectural geometry projected a numinous auditory illusion in which a booming, *pututu*-like voice appeared to emanate directly from the Lanzón monolithic deity—the "Great Voice" and oracle of Chavín. Conversely, open-air prehistoric aggregation sites and monumental structures, such as the sound-reflecting rock wall at the Cholula pyramid, were acoustically configured to maximize speech clarity and amplify vocal projection across large assemblies. Modern computational acoustics now leverages virtual reconstructions to recover the intangible sonic heritage of historical monuments—including sacred and civic buildings damaged by fire prior to their physical restoration—allowing empirical evaluation of how reverberation, echoes, and spectral filtering shaped ancient ritual cognition.

2. Physical Acoustics, Binaural Spatialization, and Cochlear Biomechanics

2.1 Wave Propagation and Head-Related Directional Filtering

In standard atmospheric conditions, acoustic waves propagate through air as longitudinal travelling pressure disturbances at a velocity of $c \approx 344\text{m/s}$ (1,128ft/s). For a sinusoidal wave of frequency f, the physical wavelength $\lambda = c / f$ determines how the wave interacts with anatomical structures of the listener's body. When λ is large relative to the physical dimensions of the human head and torso (low frequencies), sound waves diffract around the listener with minimal attenuation; however, as frequency increases and λ approaches the dimensions of the shoulders, head, and pinnae, complex scattering, shadowing, and resonance effects emerge.

The external ear consists of the pinna and the external auditory canal (meatus), which terminate at the tympanic membrane (eardrum). Acoustically, the pinna functions both as an energy collector and as a direction-dependent spectral filter, while the auditory canal acts as a quarter-wave acoustic waveguide that guides pressure oscillations to the eardrum. Quantitative ratios of sound pressure at the eardrum relative to the free-field

approaching wave (expressed in decibels, dB) reveal that external ear resonances generate substantial passive amplification between 2,000Hz and 5,000Hz, reaching a primary gain peak at approximately 2,500Hz.

Binaural spatial localization relies on the central nervous system's parallel extraction of temporal and spectral disparities across the two ears. In the azimuthal (horizontal) plane, judging whether a sound source lies to the left or right depends on a duplex mechanism: interaural time differences arising from disparities in the time of arrival of the wavefront at the two ears, which dominate at low frequencies, and interaural level differences generated by the acoustical shadow cast by the head, which become pronounced at high frequencies. Along the median plane—the vertical plane bisecting the head into symmetrical left and right halves—interaural time and level differences vanish entirely. Consequently, resolving whether a sound originates from ahead, behind, or at a specific vertical elevation depends on the convolutions of the pinnae, which impose direction-specific spectral notches and peaks that rise and fall with increasing frequency. When these pinna cues and head-tracking dynamics are bypassed, as occurs during standard stereophonic headphone listening, listeners lose the perceptual externalization of the sound field and instead experience an intracranial lateralization confined to a line connecting the two ears.

2.2 Cochlear Micromechanics and the Greenwood Frequency-Position Function

At the tympanic membrane, airborne pressure fluctuations are converted into mechanical vibrations of the middle ear ossicles, which provide impedance matching to excite travelling fluid waves within the liquid-filled inner ear (cochlea). Rather than behaving as a passive Fourier filter bank, the mammalian cochlea relies on an active biomechanical process localized within the outer and inner hair cells of the Organ of Corti (OC). This active cellular mechanism amplifies weak acoustic signals by several hundred-fold, sharpens frequency selectivity far beyond passive hydrodynamic limits, and compresses a vast range of acoustic intensities into the physiological dynamic range of auditory nerve fibers. Transduction into bioelectric nerve impulses occurs across approximately 3,500 inner hair cells arrayed systematically along the two-and-a-half turns of the basilar membrane, which project via the eighth cranial (vestibulocochlear) nerve toward the auditory brainstem and primary auditory cortex in Heschl's gyrus.




Figure 1: Biophysical Transduction from Free-Field Acoustic Waves to Primary Auditory Cortex

Acoustic pressure waves travelling at 344 meters per second undergo directional spectral filtering by the pinna and resonant amplification peaking near 2500 Hz within the auditory canal. Mechanical vibrations of the tympanic membrane and middle ear ossicles excite travelling waves along the liquid-filled cochlea, where 3500 inner hair cells transduce frequency-place coordinates into phase-locked action potentials along the eighth cranial nerve to Heschl's gyrus.

Along the cochlear duct—measured from the center of the round window at the base to the helicotrema at the apex—the mechanical stiffness and width of the basilar membrane vary continuously, establishing a logarithmic tonotopic map in which high frequencies excite the basal turn and low frequencies propagate to the apical turn. This spatial-spectral projection is formalized across mammalian species by the Greenwood frequency-position function:

$$f(x) = A\left(10^{\{a\,x\}} - K\right)$$

where f(x) represents the characteristic tonotopic frequency in Hertz (Hz), x denotes the fractional distance along the cochlear spiral measured from the apical end ($0 < x < 1$, or alternatively measured in millimeters from the apex), A is a species-specific frequency scaling constant, a governs the logarithmic slope of the tonotopic gradient, and K is a constant of integration reflecting the divergence from pure logarithmic scaling near the lower audible frequency limit. For the human auditory system, empirical histological and psychoacoustic critical-band measurements establish the canonical parameter values A = 165.4Hz, a = 2.1 (when x is expressed as a normalized proportion of total

cochlear length, or $a = 0.06\, mm^{\{-1\}}$ when x is measured in millimeters), and K = 0.88.

2.3 Spiral Ganglion Morphometrics and Clinical Anatomy-Based Fitting

While the Greenwood function accurately defines characteristic frequencies along the Organ of Corti, clinical interventions such as Cochlear Implants (CIs) electrically stimulate the cell somata and axons of the Spiral Ganglion (SG) housed within Rosenthal's canal. High-resolution three-dimensional morphometric reconstructions of nine human temporal bone specimens (Table 1) demonstrate a profound anatomical divergence between the trajectory of the Organ of Corti and that of the Spiral Ganglion. Across these specimens, the human Organ of Corti exhibits an average total length of 33.13 ± 2.11mm (ranging from 30.50mm in specimen 7R to 36.87mm in specimen 14R and extending past 990° of angular rotation from the round window), whereas the Spiral Ganglion averages only 13.69 ± 0.80mm in length (ranging from 12.54mm to 14.62mm, yielding a highly conserved SG-to-OC length ratio of 0.41 ± 0.01 and terminating near 720°). Similarly, the cochlear middle wall (MW) averages 15.49 ± 0.69mm (a mean difference of 1.81pm 0.20mm relative to the SG, and a MW-to-OC length ratio ranging from 0.44 to 0.49 with a mean of 0.47). Furthermore, Rosenthal's canal terminates prior to both extremes of the basilar membrane, leaving the most basal 2.35% ± 0.90% and the most apical 10.72% ± 1.33% of the Organ of Corti without directly radial spiral ganglion cell somata.

Table 1: Three-Dimensional Morphometric Measurements of the Organ of Corti (OC), Spiral Ganglion (SG), and Middle Wall (MW) Across Nine Human Temporal Bone Specimens

Specimen / Metric	4R	5R	6R	7R	8R	13 L	14 R	15 L	16 R	Mean ±\pm± SD
OC Length (mm)	32.31	33.44	31.41	30.50	36.16	33.09	36.87	32.45	31.95	33.13 ± 2.11
SG Length (mm)	14.00	14.20	NA	12.54	14.58	13.41	14.62	13.44	12.70	13.69 ± 0.80

Specimen / Metric	4R	5R	6R	7R	8R	13 L	14 R	15 L	16 R	Mean ±\pm± SD
SG / OC Length Ratio	0.43	0.42	NA	0.41	0.40	0.41	0.40	0.41	0.40	0.41 ±0.01
MW Length (mm)	15.79	15.90	NA	14.35	16.24	15.20	16.33	15.15	14.97	15.49 ± 0.69
MW vs. SG Difference (mm)	1.79	1.70	NA	1.81	1.66	1.79	1.71	1.71	2.27	1.81 ± 0.20
Basal OC without SG (%)	1.98	3.47	NA	1.31	0.40	3.41	1.84	2.13	3.29	2.35 ± 0.90
Apical OC without SG (%)	9.22	11.33	NA	11.64	10.26	9.40	12.99	11.37	9.58	10.72 ± 1.33

Specimen / Metric	4R	5R	6R	7R	8R	13 L	14 R	15 L	16 R	Mean ±\pm± SD
OC Length at Inflection V (%)	59.36	64.74	60.01	59.08	55.56	59.56	NA	61.14	60.31	59.97 ± 2.54
Frequency at Inflection V (Hz)	1,035	764	998	1,051	1,273	1,023	NA	937	982	1,000 ± 140

Section 2.3: Spiral Ganglion Morphometrics and Clinical Anatomy-Based Fitting.

Tracing the radial nerve fibers from the Organ of Corti to the Spiral Ganglion across angular rotation intervals from the round window (Table 2) reveals the precise spatial mechanics of this tonotopic compression. Across the basal 60% of the cochlea, radial nerve fibers follow a nearly perpendicular trajectory, so characteristic frequencies along the Organ of Corti and the Spiral Ganglion remain closely aligned from 0° (17,857Hz OC mean vs. 17,225Hz SG mean) through 90° (6,501Hz OC vs. 6,193Hz SG), 180° (3,239Hz OC vs. 3,174Hz SG), and 270° (1,659Hz OC vs. 1,539Hz SG). However, at a critical anatomical inflection point V—located at an average of 59.97% ± 2.54% of the Organ of Corti length from the base (or 62% ± 3.2% where tangential deviation becomes pronounced, corresponding to approximately 360° of rotation, 81% ± 3.1% of total SG length, and a mean characteristic frequency of 1,000 ± 140Hz)—the nerve fibers shift to an increasingly tangential course. By 360° of rotation, the mean OC frequency reaches 920Hz (range 809 - 1,063Hz) while the corresponding SG frequency drops to 785Hz (range 676 - 916Hz), and by 475.6° ± 23.2° the Greenwood function maps the 500Hz stimulus locus. Consequently, the entire apical 40% of the Organ of Corti (covering all frequencies ≤ 1,000Hz, which requires ~ 20mm of lateral wall insertion depth under the OC to reach but only 12 - 13mm for an ideally positioned perimodiolar array) is compressed into the final ~ 19% of the Spiral Ganglion.

Table 2: Characteristic Tonotopic Frequencies and Inter-Subject Ranges Along the Organ of Corti (OC) and Spiral Ganglion (SG) by Angle of Rotation from the Round Window

Angle of Rotation from Round Window	OC Mean Frequency (Hz\text{Hz}Hz)	OC Frequency Range (Hz\text{Hz}Hz)	SG Mean Frequency (Hz\text{Hz}Hz)	SG Frequency Range (Hz\text{Hz}Hz)
0°	17,857	17,623 - 18,120	17,225	14,734 - 20,677
90°	6,501	5,867 - 7,253	6,193	5,111 - 7,132
180°	3,239	2,812 - 3,567	3,174	2,700 - 3,496
270°	1,659	1,460 - 1,906	1,539	1,381 - 1,832
360°	920	809 - 1,063	785	676 - 916

Section 2.3: Spiral Ganglion Morphometrics and Clinical Anatomy-Based Fitting.

This anatomical compression has profound second-order consequences for musical pitch perception and speech intelligibility in cochlear implant recipients. Because the basal turn alone accounts for the first ~ 20mm of the cochlear duct, conventional short electrode arrays (14 - 19.1mm of active stimulation length, or 20mm insertion depth) stimulate only a single turn. Depending on individual cochlear duct length (CDL, which averages 31.5mm} and ranges up to 35mm), a 20mm} insertion reaches only the 917Hz} tonotopic location in a slightly longer cochlea (32.5mm CDL), 820Hz in an average cochlea (31.5mm CDL), and 640Hz in a smaller-than-average cochlea (29.5mm CDL). However, the fundamental frequencies (F_0) and lower harmonics/formants (F_1 - F_4) of human vocal communication and musical harmony reside below 700Hz}—specifically, male voice exhibits $F_0 \approx 120$, $F_1 \approx 240$Hz, $F_2 \approx 360$Hz, $F_3 \approx 480$Hz, and F_4

≈ 600Hz, while female voice exhibits $F_0 \approx 200$Hz, $F_1 \approx 400$Hz, and $F_2 \approx 600$Hz, and the C_5 key of a piano equates to approximately 513Hz.

When short arrays fail to reach the apical region in the second turn, these low-to-mid pitches cannot be naturally mapped according to the Greenwood function and must be artificially compressed and transposed onto a higher octave, distorting harmonic ratios and degrading musical timbre. Conversely, long, flexible lateral-wall electrode arrays (28mm, 31.5mm, or 34mm) achieve complete cochlear coverage across both the first and second turns down to approximately 100Hz. Clinical trials in twelve single-sided deafness (SSD) cochlear implant patients—utilizing clinical imaging to reconstruct individual electrode contact locations relative to the basilar membrane and testing pure tones with 5Hz triangular frequency modulation (± 5% nominal deviation to prevent standing-wave amplitude fluctuations)—confirm that when long arrays are combined with fine-structure stimulation and Anatomy-Based Fitting (ABF), subjective pitch matches align closely with the Greenwood function without statistically significant basal shift. In these evaluations, pitch-matching consistency across N trials is quantified in semitones via the logarithmic deviation:

$$\Delta f_{PM} = 12 \log_2 \left(\frac{f_{PM}}{f_{PM}}\right), \qquad \Delta f_{RMS} = \sqrt{\frac{1}{N}\sum_{i=1}^{N} (\Delta f_{PM}, i)^2}$$

where smaller values of $\mathbf{\Delta f_{RMS}}$ confirm high tonotopic precision and superior musical sound quality.

3. Psychoacoustics, Sensory Dissonance, and Auditory Scene Analysis

3.1 Perceptual Dimensions and Critical Band Filtering

Psychoacoustics investigates the non-linear mapping between physical parameters of an acoustic waveform and the subjective psychological attributes constructed by the listener. Across the literature, four primary psychoacoustic properties govern music perception: **pitch** (the perceptual correlate of fundamental frequency and periodic temporal fine structure), **loudness** (the non-linear correlate of sound pressure intensity across frequency), **duration** (the subjective scaling of temporal intervals and rhythm), and **timbre** (the multidimensional attribute governed by spectral envelope, attack transients, and harmonicity that enables discrimination between instruments playing at identical pitch and loudness). When multiple sound sources radiate simultaneously, the auditory system relies on **Auditory Scene Analysis**—utilizing cues of common onset, harmonic coherence, spatial proximity, and spectral continuity—to segregate or fuse overlapping acoustic streams into discrete perceptual objects.

At the core of spectral integration is the concept of the **critical band** (quantified psychoacoustically via the Bark scale, Mel scale, or Equivalent Rectangular Bandwidth,

ERB), which corresponds to an approximately constant physical distance and number of hair cells along the basilar membrane. When two sinusoidal components fall within the same critical bandwidth, their mechanical excitation patterns overlap on the basilar membrane, generating rapid amplitude fluctuations perceived as beats and sensory roughness.

3.2 Plomp-Levelt Consonance and Sethares's Spectral-Scale Coupling

Historically, music theory attributed the consonance of musical intervals (such as the octave 2:1 or perfect fifth 3:2) to abstract mathematical ratios, whereas Helmholtz linked consonance to the absence of dissonant interference beats between upper partials. In 1965, Plomp and Levelt advanced this framework empirically by systematically measuring listener consonance ratings for pairs of pure sine waves in a controlled environment across the audible spectrum. Their empirical curves demonstrated that two pure tones are perceived as maximally dissonant when their frequency separation equals approximately one-quarter of the local critical bandwidth, and become fully consonant once their separation exceeds the critical bandwidth.

Building upon Plomp and Levelt's pure-tone interference curves, William Sethares demonstrated that sensory consonance and dissonance are not intrinsic, invariant properties of musical intervals, but rather depend directly on the spectral timbre of the sounding instrument. For two complex tones composed of partials with frequencies $f_1 < f_2$ and amplitudes a_1, a_2, the elementary Plomp-Levelt sensory dissonance $d(f_1, f_2, a_1, a_2)$ is parameterized as a difference of exponentials scaled by the critical bandwidth factor $s(f_1)$:

$$d(f_1, f_2, a_1, a_2) = a_{12} \left[e^{-b_1 s(f_1)(f_2 - f_1)} - e^{-b_2 s(f_1)(f_2 - f_1)} \right],$$

$$s(f_1) = \frac{d^*}{s_1 f_1 + s_2}$$

where a_{12} reflects the combined partial amplitudes (such as $\min(a_1, a_2)$ or $a_1 a_2$) and $b_1 = 3.5$, $b_2 = 5.75$, $d^* = 0.24$, $s_1 = 0.0207$, and $s_2 = 18.96$ fit the empirical Plomp-Levelt roughness curve. Summing $d(f_i, f_j, a_i, a_j)$ over all pairs of partials in a spectrum F and its pitch-shifted copy αF yields the total intrinsic dissonance curve $D_F(\alpha)$ across interval ratios α. When an instrument possesses a harmonic spectrum ($f_n = n f_0$, characteristic of bowed strings and aerophones), local minima of $D_F(\alpha)$ occur precisely at simple integer ratios (2:1, 3:2, 4:3, 5:4), explaining the cultural convergence toward just and twelve-tone equal-tempered scales. Conversely, when an instrument exhibits inharmonic partials (such as metallophones, bells, or synthesized stretched spectra), the minima of $D_F(\alpha)$ shift to non-integer microtonal ratios, proving that musical scales and instrumental timbres co-evolve as coupled psychoacoustic systems.

3.3 Boundary Acoustic Regimes: Infrasound and the Hypersonic Effect

Beyond the canonical human audible spectrum (20Hz to 20kHz), acoustic energy continues to exert measurable physiological and neurobiological effects. At the low-

frequency boundary, **infrasound** (< 20Hz) generated by natural geophysical phenomena or complex industrial environments poses recondite hazards to human health, requiring physiological and vestibular adaptation. Conversely, at the ultra-high-frequency boundary (> 20kHz), neurophysiological studies have uncovered the **hypersonic effect**. Although human listeners cannot consciously detect isolated sine waves above 20kHz, presenting complex acoustic stimuli (such as traditional gamelan music or natural soundscapes) that retain rich, non-stationary high-frequency components above the audible range significantly enhances regional cerebral blood flow in deep brain structures and increases alpha-band electroencephalographic coherence compared to high-cut filtered audio (< 20kHz). This finding has direct implications for digital audio engineering and archival standards, demonstrating that conventional 44.1kHz Nyquist sampling limits omit biologically active ultrasonic modulation.

4. Cognitive Neuroscience of Music: Syntax, Reward, and Genetics

4.1 Hierarchical Processing from Brainstem to Auditory Cortex

The neurocognitive processing of music begins subcortically: electroencephalographic (EEG) recordings of the **Auditory Brainstem Response (ABR)** demonstrate that brainstem nuclei phase-lock to the incoming acoustic signal with sub-millisecond precision, preserving the temporal fine structure and pitch contour of the original sound waveform. As information ascends to the primary auditory cortex—specifically the medial section of **Heschl's gyrus (HG)** and surrounding planum temporale—functional neuroimaging and invasive intracranial recordings reveal a spatially organized, hierarchical representation that transforms raw spectral-temporal features into abstract relative pitch intervals, melodic contours, harmonic progressions, and chromatic tonal contexts.

4.2 Musical and Linguistic Syntax: The Shared Syntactic Integration Resource Hypothesis

A foundational paradox in cognitive neuroscience is the relationship between music and language: neuropsychological case studies demonstrate double dissociations in which individuals suffer selective loss of musical pitch processing (congenital or acquired amusia) without aphasia, or severe linguistic aphasia while retaining musical abilities, yet functional neuroimaging consistently shows overlapping cortical activation during musical and linguistic structural tasks.

To resolve this paradox, Patel (2003, 2008) formulated the **Shared Syntactic Integration Resource Hypothesis (SSIRH)**, which distinguishes between *domain-specific syntactic representations* stored in long-term memory and *domain-general neural resources* that dynamically activate and integrate those representations during real-time processing. Under the SSIRH, linguistic grammar and musical tonal-harmonic

syntax reside in anatomically distinct posterior and temporal storage networks (explaining why focal brain damage can selectively impair one domain while sparing the other), whereas both domains recruit shared frontal cognitive control and structural integration circuits—most notably **Brodmann Area 47 (BA 47)** in the inferior frontal cortex and Broca's territory—to establish temporal coherence and resolve structural expectations.

Empirical confirmation of the SSIRH emerges from converging electrophysiological, behavioral dual-task, and predictive coding investigations. In event-related potential (ERP) studies where listeners process sentences and chord progressions that vary in structural integration difficulty, syntactic violations in both language and music elicit an identical positive-going ERP component—the **P600**—with an indistinguishable centro-parietal scalp distribution, indexing shared neural work during structural reanalysis. Behavioral dual-task experiments recorded at Stanford's Center for Computer Research in Music and Acoustics (CCRMA) provide further causal proof by manipulating linguistic syntactic complexity (comparing difficult object-extracted relative clauses against simpler subject-extracted relative clauses) alongside simultaneous melodic pitch expectations. When the first five notes establish a strong sense of key and the critical sixth note—falling precisely on the final word of the relative clause—is rendered **out-of-key**, a selective super-additive interference interaction occurs during object-extracted relative clause comprehension. Crucially, as predicted by the SSIRH, this cross-domain interference arises specifically for the structural harmonic violation (out-of-key note) and does not occur for a lower-level psychoacoustic deviance (an in-key note presented at an unexpectedly loud intensity), confirming that resource sharing operates at the level of structural integration rather than general auditory attention. Contemporary theoretical refinements comparing the SSIRH with the Syntactic Equivalence Hypothesis situate these shared prefrontal mechanisms within a broader **hierarchical predictive processing** and executive cognitive control framework. As musical and linguistic sequences unfold in time, prefrontal-temporal circuits incrementally generate probabilistic predictions; when an unexpected harmonic modulation or syntactic garden path occurs, domain-general cognitive control resources in the inferior frontal cortex are recruited to override initial expectations and update the internal structural representation.




Figure 2: Cortical Syntactic Integration Networks and Subcortical Striatal Dopamine Dissociation in Music Cognition

Left: Under the Shared Syntactic Integration Resource Hypothesis, domain-specific musical and linguistic representations reside in distinct temporal storage networks but converge on shared structural integration and cognitive control resources in the inferior frontal cortex (Brodmann Area 47), indexed electrophysiologically by the P600 component. Right: Positron emission tomography and functional magnetic resonance imaging reveal anatomically distinct dopaminergic pathways in the striatum: the right caudate nucleus mediates the 15-second anticipatory phase prior to peak emotion, whereas the right nucleus accumbens mediates consummatory euphoria (chills) via mesolimbic projections from the ventral tegmental area.

4.3 Neurochemistry of Musical Pleasure: Striatal Dopamine Dissociation

Why does an abstract sequence of acoustic vibrations—devoid of direct homeostatic survival value like food or shelter—evoke intense euphoria, autonomic arousal, and "chills" (musical frisson) across all human societies? Neuroimaging investigations demonstrate that pleasurable music recruits the core mesolimbic reward circuitry, significantly enhancing functional and effective connectivity between the **Ventral Tegmental Area (VTA)**, the **Nucleus Accumbens (NAcc)**, the **hypothalamus**, and the **insula** (which integrates autonomic visceral states).

In a landmark neurochemical investigation, Salimpoor et al. (2011) combined the molecular specificity of [^{11}C]raclopride Positron Emission Tomography (PET)—which measures endogenous dopamine release via competitive D2/D3 receptor binding potential displacement—with the high temporal resolution of blood-oxygen-level-dependent (BOLD) functional Magnetic Resonance Imaging (fMRI) and continuous autonomic psychophysiology while participants listened to self-selected chill-inducing

instrumental music versus neutral control music. Their data revealed an anatomical and temporal dissociation within the human striatum between the *anticipation* and the *consummation* of musical reward (Table 3):

Table 3: Anatomical and Temporal Dissociation of Striatal Dopamine Release During Anticipatory and Consummatory Phases of Musical Reward

Striatal Subregion	Functional Reward Phase	Temporal Epoch Definition	Stereotaxic Peak Coordinates (x,y,zx, y, zx,y,z)	Statistical Peak & Psychophysiological Coupling
Right Caudate Nucleus (Dorsal Striatum)	**Anticipatory Phase** (Prediction, tension buildup, and incentive salience)	15s immediately preceding peak emotional response	(14, -6, 20)	t = 3.2; significant [^{11}C]raclopride binding potential change; coupled to structural expectation
Right Nucleus Accumbens (Ventral Striatum)	**Consummatory Phase** (Peak emotional pleasure, euphoria, and chills)	Exact duration of reported peak pleasure / autonomic chill	(8, 10, -8)	t = 2.8; PET dopamine release P < 0.001; BOLD signal positively correlated with chill intensity (P < 0.01)

Section 4.3: Neurochemistry of Musical Pleasure: Striatal Dopamine Dissociation.

This dorsal-to-ventral striatal cascade bridges the cognitive neuroscience of syntax with the neurobiology of emotion. During the 15-second anticipatory epoch—when melodic tension, rhythmic syncopation, or harmonic delay suspends resolution—dopaminergic transmission surges in the **right caudate nucleus**, which is heavily interconnected with prefrontal syntactic prediction networks. When the expected resolution arrives (or violates expectations in an aesthetically optimal manner), **musical reward prediction errors** trigger phasic dopamine release in the **right nucleus accumbens**, generating subjective euphoria and reinforcing auditory statistical learning. Pharmacological causal manipulations confirm that modulating dopaminergic transmission directly alters both the subjective pleasure and motivational value elicited by music.

4.4 Genetic Architecture and Clinical Resilience

The biological hardwiring of music is underscored by behavioral genetics and clinical neurology. Pedigree studies of **hereditary congenital amusia** (a selective neurodevelopmental deficit in fine-grained musical pitch discrimination) reveal a high familial aggregation across multiplex families, with a sibling relative risk of $\lambda_s = 10.8$ (95% confidence interval: 8.0 - 13.5), enabling the mapping of specific genetic loci governing auditory-cortical connectivity. Conversely, as documented by neurologist Oliver Sacks (2006), because musical perception and production recruit a distributed, redundant bilateral network spanning emotional, motor, cerebellar, and cognitive regions, musical responsiveness and procedural memory frequently represent the final cognitive faculties preserved in patients suffering from severe brain injury or advanced cortical neurodegeneration.

5. Clinical Neuromusicology and Neurologic Music Therapy (NMT)

The resilience and distributed connectivity of the musical brain provide the neurobiological foundation for **Neurologic Music Therapy (NMT)**, an evidence-based clinical system comprising 20 standardized treatment techniques across sensorimotor, speech-language, and cognitive rehabilitation domains. Unlike traditional interpretive or recreational music therapy, NMT operates on the neuroscience model of rhythmic auditory-motor entrainment and structural neural plasticity.

5.1 Auditory-Motor Entrainment and the Basal Ganglia Bypass in Parkinson's Disease

Human bipedal locomotion relies on the intrinsic generation of alternating left-right rhythmic motor patterns coordinated by the basal ganglia and supplementary motor area. In **Parkinson's disease (PD)**, the progressive degeneration of nigrostriatal dopaminergic neurons disrupts internal neural timing cues, manifesting clinically as bradykinesia, shuffling stride, festination, and debilitating **freezing of gait (FOG)**. Similar temporal motor disruptions occur following **stroke**, **traumatic brain injury (TBI)**, and **spinal cord injury (SCI)**.

Rhythmic Auditory Stimulation (RAS)—the flagship sensorimotor technique of NMT—delivers external isochronous or musically embedded rhythmic pulses matched or incrementally scaled to the patient's cadence. Neurophysiologically, periodic acoustic signals excite the ascending reticular and cochlear pathways, which project via direct structural and functional tracts to spinal motor neurons and the **cerebello-thalamo-cortical network**. By entraining motor neuron excitability to an external auditory clock, RAS **bypasses the defective basal ganglia loop**, synchronizing muscle contraction timing across both spatial and temporal dimensions to normalize stride length, gait velocity, and symmetry. Recent clinical trials demonstrate that coupling NMT and RAS

with **immersive Virtual Reality (VR)** environments creates synergistic multimodal feedback that further enhances functional mobility and gait quality in PD patients.

5.2 Speech, Cognitive, and Non-Musical Acoustic Interventions

Beyond gait rehabilitation, structured musical and acoustic interventions target language, executive function, and autonomic regulation across diverse clinical populations (Table 4):

Table 4: Clinical Neuromusicology and Acoustic Intervention Modalities, Target Pathologies, Neurobiological Mechanisms, and Functional Outcomes

Clinical Modality / Protocol	Target Pathology or Population	Primary Neurobiological / Psychoacoustic Mechanism	Documented Clinical and Functional Outcomes
Rhythmic Auditory Stimulation (RAS) & NMT-VR	Parkinson's disease (PD), stroke, TBI, spinal cord injury	External rhythmic entrainment via cerebello-thalamo-cortical tracts bypassing basal ganglia deficits	Reduction in freezing of gait; improved stride length, cadence symmetry, and functional mobility
Rhythmic Reading Training (RRT)	Acquired dyslexia in post-stroke aphasia	Meter-syntax synchronization and shared prefrontal temporal coherence (BA 47)	Improved phonological decoding, reading fluency, and morpho-syntactic integration
Instrumental & Movement Music Training	Preschool children and older adults	Cross-domain plasticity in prefrontal executive control and auditory working memory	Enhanced inhibitory control in children; long-term memory retention and working memory in older adults

Clinical Modality / Protocol	Target Pathology or Population	Primary Neurobiological / Psychoacoustic Mechanism	Documented Clinical and Functional Outcomes
Brainwave Entrainment (BWE)	Chronic pain, migraines, PMS, stress, and cognitive deficits	Frequency-following cortical synchronization and inter-hemispheric brain coherence	Alleviation of headache/migraine severity, behavioral stabilization, and stress reduction
Therapeutic Prosody & Poetic Acoustics	Anxiety, chronic stress, and psychotherapeutic rapport	Acoustic modulation of vocal warmth, formant spacing, and poetic rhythmic cadence	Reduction in physiological stress markers and strengthening of client-therapist therapeutic alliance

Section 5.2: Speech, Cognitive, and Non-Musical Acoustic Interventions.

Crucially, non-musical sound therapies—including natural environmental soundscapes, rhythmic chanting, and listening to metered poetry—also exert robust anxiolytic and analgesic effects. In clinical psychotherapy, the acoustic prosody of the therapist's voice (conveying warmth, interest, and curiosity through phase-aligned pitch and timbre modulation) serves as a primary determinant of the therapeutic alliance, although disentangling semantic linguistic content from pure speech acoustics remains a central methodological challenge.

6. Computational Phonogram Archiving and Generative Audio Artificial Intelligence

6.1 From Symbolic Representations to Neural Audio Codecs

The computational modeling of music and sound has undergone a paradigm shift from symbolic sequence modeling and feature-engineered Music Information Retrieval (MIR) to end-to-end generative learning directly over raw audio waveforms. In the symbolic domain, representations such as **REMI** (Revamped MIDI) and **REMI+** encode polyphonic multi-track compositions as discrete token sequences representing

Pitch, **Velocity**, **Duration**, **Bar**, **Position**, and multi-instrument **Program** identifiers, enabling Transformer architectures to capture long-range metric and harmonic structure. Similarly, supervised acoustic classifiers have achieved high industrial precision—for example, binary acoustic gunshot detection systems achieving 95.02% sensitivity in gunshot identification and 98.16% specificity in rejecting non-gunshot environmental noise.

However, generating high-fidelity raw audio (24kHz to 48kHz sampling rates) directly in the time domain presents an intractable sequence-length bottleneck for autoregressive Transformers. To bridge continuous waveform physics with discrete language modeling, contemporary generative audio frameworks—such as Meta AI's **AudioCraft** (encompassing **MusicGen**, **AudioGen**, and **EnCodec**), **SoundStream**, and **Descript Audio Codec (DAC)**—employ **Neural Audio Codecs** to compress continuous waveforms into compact streams of **discrete audio tokens**.

6.2 Mathematical Taxonomy of Vector Quantization in Neural Audio Codecs

A neural audio codec comprises three core components: an encoder network E (built from Convolutional Neural Networks [CNN], Recurrent Neural Networks [RNN], Transformers, or hybrid CNN+Transformer blocks) that maps a raw waveform or time-frequency representation into continuous latent frames $z_t \in R^D$; a discrete quantizer Q that maps z_t to codebook indices; and a decoder G that reconstructs the waveform $\hat{x}$. Across contemporary architectures, discrete audio tokenization is governed by several distinct quantization paradigms (Table 5).

In semantic self-supervised learning (SSL) models such as Wav2Vec-BERT and USM, **Post-Training K-Means Quantization** extracts discrete tokens from frozen latent representations by assigning each continuous latent frame z_t to the nearest centroid c_k in a codebook of size K via Euclidean distance minimization, $q_t = \arg\min_{k\in\{1,\ldots,K\}} ||z_t - c_k||_2^2$, whereas **Product Quantization (PQ)** partitions embeddings into sub-vectors and quantizes each partition independently. For acoustic waveform reconstruction in **EnCodec**, **SoundStream**, and **DAC**, **Residual Vector Quantization (RVQ)** avoids exponential codebook growth by cascading M hierarchical vector quantizers, where each stage $m \in \{1, \ldots, M\}$ recursively quantizes the residual error left by the preceding stage:

$$r_t^{(1)} \leftarrow z_t, \quad q_t^{(m)} \leftarrow \arg\min_{k\in\{1,\ldots,K\}} |r_t^{(m)} - c_k^{(m)}|_2^2, \quad \hat{z}_t^{(m)} \leftarrow c_{q_t^{(m)}}^{(m)}, \quad r_t^{(m+1)} \leftarrow r_t^{(m)} - \hat{z}_t^{(m)}$$

The final quantized latent vector is reconstructed via the linear superposition $\hat{z}_t = \sum_{m=1}^{M} \hat{z}_t^{(m)}$. Because $r_t^{(1)}$ captures the dominant spectral-temporal envelope while higher stages $m > 1$ encode progressively finer acoustic residuals, RVQ naturally mirrors the hierarchical coarse-to-fine processing of the human auditory pathway and supports scalable bitrate streaming.

To eliminate multi-stream interleaving complexity during downstream autoregressive language modeling, **Single Vector Quantization (SVQ)** restricts discretization to a single codebook (M = 1, utilized in **BigCodec**, **TS3-Codec**, and **WavTokenizer**), whereas **Group Vector Quantization (GVQ)** divides the latent vector z_t into independent feature groups and applies RVQ to each subspace separately (as implemented in **HiFi-Codec** and **Prompt Codec**). Alternatively, **Finite Scalar Quantization (FSQ)** projects z_t into a low-dimensional feature space and rounds each scalar coordinate independently to a fixed set of discrete scalar values (deployed in **SQ-Codec**, **Spectral Codecs**, and **HARP-Net**). Crucially, channel transmission simulations demonstrate that **FSQ achieves vastly superior bit-level perturbation robustness compared to RVQ** when code sequences are transmitted over noisy communication channels, because isolated bit errors in FSQ do not trigger catastrophic hierarchical residual misalignment. Furthermore, **Multi-Scale RVQ (MSRVQ)** and **Cross-Scale RVQ (CSRVQ)** integrate multi-scale features and encode residuals across multiple hierarchical temporal resolutions (such as in **S-TFNet** and **ESC**), achieving high audio fidelity at substantially reduced quantizer complexity.

Table 5: Mathematical Taxonomy of Vector Quantization Paradigms in Neural Audio Codecs and Self-Supervised Audio Architectures

Quantization Paradigm	Codebook Structure & Mathematical Mechanism	Representative Codec & SSL Architectures	Primary Computational and Perceptual Trade-offs
K-Means & Product Quantization (PQ)	Post-training centroid assignment $q_t = argmin_k \|(z_t - c_k)_2^2$ or sub-vector partitioning	Wav2Vec-BERT, USM, EncodecMAE	Rich semantic representation from frozen SSL features; limited raw acoustic waveform fidelity
Residual Vector Quantization (RVQ)	Recursive cascade of M codebooks quantizing residual $r_t^{((m+1))} = r_t^{(m)} - \hat{z}_t^{(m)}$	SoundStream, EnCodec, DAC	High waveform fidelity and scalable bitrate; requires multi-codebook delay patterns in language models

Quantization Paradigm	Codebook Structure & Mathematical Mechanism	Representative Codec & SSL Architectures	Primary Computational and Perceptual Trade-offs
Single Vector Quantization (SVQ)	Single unified codebook (M = 1) without residual cascading	BigCodec, TS3-Codec, WavTokenizer	Simplifies downstream autoregressive acoustic language modeling to a single token stream
Group Vector Quantization (GVQ)	Partitions latent frame z_t into feature groups and applies RVQ independently per group	HiFi-Codec, Prompt Codec	Expands representational subspace capacity for complex polyphonic and speech textures
Finite Scalar Quantization (FSQ)	Projects latent vector and quantizes each scalar dimension to fixed discrete values	SQ-Codec, Spectral Codecs, HARP-Net	Superior bit-level perturbation robustness over noisy transmission channels; eliminates codebook collapse
Multi-Scale & Cross-Scale RVQ (MSRVQ / CSRVQ)	Applies RVQ across multiple temporal resolutions and cross-scale residual hierarchies	S-TFNet, ESC	High perceptual audio quality with substantially lower computational and quantizer complexity

Section 6.2: Mathematical Taxonomy of Vector Quantization in Neural Audio Codecs.

Recent codec innovations have pushed rate-distortion boundaries to extreme limits: **MuCodec** reconstructs high-fidelity music at ultra-low bitrates of 0.35kbps to 1.35kbps; **FlowMAC** leverages conditional flow matching at 3kbps to match the perceptual quality of GAN- and DDPM-based codecs operating at double the bitrate; **MUFFIN** achieves a state-of-the-art 12.5Hz token frame rate; and **SemantiCodec** embeds rich semantic structure at ultra-low bitrates.




Figure 3: Neural Audio Codec Quantization and Multimodal Transformer Architecture for Controllable Music Generation

End-to-end architecture for controllable neural music generation. Continuous audio waveforms are encoded into latent vectors and discretized across hierarchical codebooks via Residual Vector Quantization (RVQ). To model parallel codebooks within a single autoregressive Transformer, the Codebook Interleaving and Delay Pattern offsets secondary codebooks temporally while inserting explicit start and end boundary tokens for seamless loop synthesis. Multi-Stage Cross Attention integrates textual prompts and symbolic REMI+ MIDI tokens, guided by Instrument-Aware Reinforcement Learning prior to neural waveform decoding.

6.3 Autoregressive Codebook Interleaving and Controllable Music Generation

Because RVQ produces M parallel discrete token streams per time step t, modeling all M × T tokens flattened sequentially would multiply the context length by M, whereas predicting all M codebooks independently in parallel at step t ignores conditional dependencies between coarse (m=1) and fine (m > 1) acoustic residuals. **MusicGen** (within the **AudioCraft** framework) resolves this trade-off via the **Codebook Delay /**

Interleaving Pattern, which shifts each codebook m by m - 1 temporal steps so that at diagonal step τ, the Transformer predicts $\left(q_{\tau}^{(1)}, q_{(\tau-1)}^{(2)}, \ldots, q_{(\tau-M+1)}^{(M)}\right)$ conditioned on all preceding steps.

To satisfy the structural demands of contemporary music production, advanced architectures such as the **Loops Transformer** (32kHz) extend the EnCodec RVQ delay pattern through four integrated algorithmic mechanisms. First, explicit loop boundary tokenization injects dedicated start (s) and end (e) tokens directly into the interleaved RVQ codebook stream, enabling the autoregressive Transformer to learn the exact metric and phase boundaries required for seamless cyclic playback. Second, a **Loop Shift** data augmentation mechanism randomly samples two integer offsets s_1 and s_2 during pre-training to vary the number of acoustic time steps padded before the start token s and after the end token e, forcing the self-attention layers to establish phase continuity between the post-s onset and the pre-e cadence. Third, a **Multi-Stage Cross Attention** architecture integrates global semantic text descriptions (capturing style, mood, and genre) with fine-grained symbolic **REMI+** MIDI sequences, bridging abstract textual intent with precise multi-track harmonic and rhythmic control. Finally, **Instrument-Aware Reinforcement Learning** fine-tunes the generative policy to ensure timbral fidelity and multi-instrument separation, while emerging omnimodal frameworks—such as **Qwen3.5-Omni** and the **VERSA** toolkit—unify audio coding, speech and singing synthesis, signal enhancement, and audio-visual scene grounding within a single generative paradigm.

7. Synthesis and Conclusions

A rigorous transdisciplinary examination of music and sound reveals a profound structural isomorphism across physical, neurobiological, and computational scales. At the peripheral transduction stage, just as the mammalian cochlea logarithmically projects a 1,000-fold frequency spectrum onto a 33.13mm basilar membrane via the Greenwood function and compresses the apical 40% of low-frequency nerve fibers (≤ 1,000Hz) into the final 19% of a 13.69mm spiral ganglion, neural audio codecs compress high-rate continuous waveforms into hierarchical residual vector spaces where coarse fundamental structures precede fine spectral residuals.

At the psychoacoustic and clinical interface, Plomp-Levelt and Sethares sensory dissonance models prove that musical harmony and tuning systems are dynamically constrained by cochlear critical bandwidths and instrumental spectral timbre rather than static numerological ratios. Consequently, restoring natural music perception and vocal formant discrimination in patients with sensorineural hearing loss requires long electrode arrays (28 - 34mm) and Anatomy-Based Fitting protocols that respect the 1,000Hz (360°) radial-to-tangential spiral ganglion inflection point V.

At the central neurocognitive level, music operates neither as an evolutionary accident nor as a purely hedonic stimulus, but as a predictive neurocognitive technology. The inferior frontal cortex (Brodmann Area 47) continuously computes structural expectations over incoming acoustic streams (indexed electrophysiologically by the P600 component under the Shared Syntactic Integration Resource Hypothesis), driving anticipatory dopamine release in the dorsal striatum (right caudate nucleus) and consummatory reward prediction signals in the ventral striatum (right nucleus accumbens), while simultaneously entraining cerebello-thalamo-cortical motor loops to bypass basal ganglia pathology in clinical neurorehabilitation.

8. Declarations

Author Identification and Contributions: SAMUELSON G conceived the transdisciplinary theoretical framework, performed the formal mathematical and morphometric synthesis across physical acoustics, cochlear neurobiology, psychoacoustics, clinical neuromusicology, and generative audio artificial intelligence, designed the data visualizations and structural tables, and wrote the final manuscript.

Competing Interests: The author (**SAMUELSON G**) declares no competing financial, commercial, or non-financial interests related to the content of this publication.

Ethical Approval and Consent to Participate: Not applicable. This study synthesizes published peer-reviewed histological, neuroimaging, clinical, and computational datasets without direct recruitment of new human or animal subjects.

Data and Code Availability: All morphometric parameters, stereotaxic neuroimaging coordinates, psychoacoustic constants, and open-source generative audio frameworks (including AudioCraft, EnCodec, MusicGen, VERSA, and MuCodec) analyzed in this manuscript are publicly accessible through their respective institutional repositories and peer-reviewed archives.

Funding Declaration: This research received no specific grant from any commercial, public, or not-for-profit funding agency.

9. References

1. Ackerman, S. J., & Hilsenroth, M. J. (2003). A review of therapist characteristics and techniques positively impacting the therapeutic alliance. *Clinical Psychology Review*, 23(1), 1–33.
2. Aletta, F., & Kang, J. (Eds.). (2020). *Historical acoustics: Relationships between people and sound over time*. MDPI Acoustics.
3. Alvarez-Morales, L., & Díaz-Andreu, M. (2024). Acoustics, soundscapes and sounds as intangible heritage. *Acoustics*.

4. Blood, A. J., & Zatorre, R. J. (2001). Intensely pleasurable responses to music correlate with activity in brain regions implicated in reward and emotion. *Proceedings of the National Academy of Sciences*, 98(20), 11818–11823.

5. Chanda, M. L., & Levitin, D. J. (2013). The neurochemistry of music. *Trends in Cognitive Sciences*, 17(4), 179–193.

6. Cook, P. R. (Ed.). (1999). *Music, cognition, and computerized sound: An introduction to psychoacoustics*. MIT Press.

7. Copet, J., Kreuk, F., Gat, I., Remez, T., Kant, D., Synnaeve, G., Adi, Y., & Défossez, A. (2023). Simple and controllable music generation (AudioCraft / MusicGen). *Advances in Neural Information Processing Systems (NeurIPS)*.

8. Díaz-Andreu, M. (2024). Archaeoacoustics and the sounds of the past. *Annual Review of Anthropology*, 53, 113–130.

9. Fedorenko, E., Patel, A., Casasanto, D., Winawer, J., & Gibson, E. (2009). Structural integration in language and music: Evidence for a shared system. *Memory & Cognition*, 37(1), 1–9.

10. Greenwood, D. D. (1961). Critical bandwidth and the frequency coordinates of the basilar membrane. *Journal of the Acoustical Society of America*, 33(10), 1344–1356.

11. Greenwood, D. D. (1990). A cochlear frequency-position function for several species—29 years later. *Journal of the Acoustical Society of America*, 87(6), 2592–2605.

12. Hudspeth, A. J. (2014). Integrating the active process of hair cells with cochlear function. *Nature Reviews Neuroscience*, 15(9), 600–614.

13. Koelsch, S. (2014). Brain correlates of music-evoked emotions. *Nature Reviews Neuroscience*, 15(3), 170–180.

14. Kolar, M. A., Rick, J. W., Cook, P. R., & Abel, J. S. (2012). AncientInca and Andean ritual acoustics at Chavín de Huántar. *Archaeological Acoustics*.

15. Koshimori, Y., & Thaut, M. H. (2023). Music therapy for gait and speech deficits in Parkinson's disease: A mini-review. *Brain Sciences*, 13(7), 993.

16. Levitin, D. J., & Menon, V. (2003). Musical structure is processed in "language" areas of the brain: A possible role for Brodmann Area 47 in temporal coherence. *NeuroImage*, 20(4), 2142–2152. 17.Mehr, S. A., Singh, M., Knox, D., Ketter, D. M., Pickens-Jones, D., Atwood, S., Lucas, C., Jacoby, N., Egner, A. A., Hopkins, E. J., Howard, R. M., Hartshorne, J. K., Jennings, M. V., Simson, J., Bainbridge, C. M., Pinkert, S., O'Donnell, T. J., Krasnow, M. M., & Glowacki, L. (2019). Universality and diversity in human song. *Science*, 366(6468), eaax0868.

17. Menon, V., & Levitin, D. J. (2005). The rewards of music listening: Response and physiological connectivity of the mesolimbic system. *NeuroImage*, 28(1), 175–184.

18. Patel, A. D., Gibson, E., Ratner, J., Besson, M., & Holcomb, P. J. (1998). Processing syntactic relations in language and music: An event-related potential study. *Journal of Cognitive Neuroscience*, 10(6), 717–733.

19. Patel, A. D. (2003). Language, music, syntax and the brain. *Nature Neuroscience*, 6(7), 674–681.

20. Patel, A. D. (2008). *Music, language, and the brain*. Oxford University Press.

21. Plomp, R., & Levelt, W. J. M. (1965). Tonal consonance and critical bandwidth. *Journal of the Acoustical Society of America*, 38(4), 548–560.

22. Sacks, O. (2006). The power of music. *Brain*, 129(10), 2528–2532.

23. Salimpoor, V. N., Benovoy, M., Larcher, K., Dagher, A., & Zatorre, R. J. (2011). Anatomically distinct dopamine release during anticipation and experience of peak emotion to music. *Nature Neuroscience*, 14(2), 257–262.

24. Salimpoor, V. N., van den Bosch, I., Kovacevic, N., McIntosh, A. R., Dagher, A., & Zatorre, R. J. (2013). Interactions between the nucleus accumbens and auditory cortices predict music reward value. *Science*, 340(6129), 216–219.

25. Sethares, W. A. (2005). *Tuning, timbre, spectrum, scale* (2nd ed.). Springer.

26. Slevc, L. R., & Okada, B. M. (2015). Processing structure in language and music: A case for shared reliance on cognitive control. *Psychonomic Bulletin & Review*, 22(3), 637–652.

27. Stakhovskaya, O., Sridhar, D., Bonham, B. H., & Leake, P. A. (2007). Frequency map for the human cochlear spiral ganglion: Implications for cochlear implants. *Journal of the Association for Research in Otolaryngology (JARO)*, 8(2), 220–233.

28. Thaut, M. H., & Braun Janzen, T. (2019). Neurologic music therapy. In *Foundations of Music Therapy*.

29. Wilson, B. S. (2004). Engineering design of cochlear implants. In F. G. Zeng, A. N. Popper, & R. R. Fay (Eds.), *Cochlear Implants: Auditory Prostheses and Electric Hearing* (pp. 14–52). Springer.

Works cited

1. Psychoacoustics in Music and Cognition | PDF | Neuron - Scribd, https://www.scribd.com/document/623862607/Music-Cognition-and-Computerized-Sound

2. The Psychoacoustic Properties Of Sound: An Introduction - SciSpace, https://scispace.com/pdf/the-psychoacoustic-properties-of-sound-an-introduction-3prhnnw1vk.pdf
3. Effects of Sound Interventions on the Mental Stress Response ... - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC11976171/
4. Greenwood function - Wikipedia, https://en.wikipedia.org/wiki/Greenwood_function
5. Current Research in Systematic Musicology | springerprofessional.de, https://www.springerprofessional.de/en/current-research-in-systematic-musicology/1990002
6. Processing structure in language and music: a case for shared, https://www.researchgate.net/publication/264539027_Processing_structure_in_language_and_music_a_case_for_shared_reliance_on_cognitive_control
7. [Literature Review] Discrete Audio Tokens: More Than a Survey!, https://www.themoonlight.io/en/review/discrete-audio-tokens-more-than-a-survey
8. Greenwood function | 1035 Citations | Top Authors | Related Topics, https://scispace.com/topics/greenwood-function-206am1fy
9. Archaeoacoustics: Research on Past Musics and Sounds, https://www.annualreviews.org/content/journals/10.1146/annurev-anthro-071323-113540?crawler=true&mimetype=application/pdf
10. (PDF) Anatomically distinct dopamine release during anticipation, https://www.researchgate.net/publication/49739517_Anatomically_distinct_dopamine_release_during_anticipation_and_experience_of_peak_emotion_to_music
11. Frequency Map for the Human Cochlear Spiral Ganglion - PMC - NIH, https://pmc.ncbi.nlm.nih.gov/articles/PMC2394499/
12. The Psychoacoustic Properties Of Sound: An Introduction, https://www.researchgate.net/publication/335943960_The_Psychoacoustic_Properties_Of_Sound_An_Introduction
13. Tuning, Timbre, Spectrum, Scale, http://www.r-5.org/files/books/rx-music/tuning/William_A_Sethares-Tuning_Timbre_Spectrum_Scale-EN.pdf
14. Universality and diversity in human song - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC7001657/
15. Structural integration in language and music: Evidence for a shared, https://web.mit.edu/evlab/assets/papers/Fedorenko_et_al_2009_MemCogn.pdf
16. Processing structure in language and music: a case for shared, https://pubmed.ncbi.nlm.nih.gov/25092390/
17. Neurobiological mechanism of music improving gait disorder in, https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2024.1502561/full
18. thaut-braunjanzen2019_chapter_neurologicmusictherapy.pdf, https://nmtacademy.co/wp-content/uploads/2019/05/thaut-braunjanzen2019_chapter_neurologicmusictherapy.pdf
19. Daily Papers - Hugging Face, https://huggingface.co/papers?q=audio%20coding
20. Controllable Music Loops Generation with MIDI and Text via Multi,

https://openreview.net/pdf?id=Whe0cYAui4
21. Review of Aniruddh D. Patel. Music, language, and the brain., https://casasanto.com/papers/Casasanto_PatelReview.pdf
22. Universality and diversity in human song - Penn State, https://pure.psu.edu/en/publications/universality-and-diversity-in-human-song/
23. The Greenwood function shows close alignment with pitch, https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2025.1624499/full
24. Cognitive Neuroscience of Music | PDF | Auditory System - Scribd, https://www.scribd.com/document/468598016/Cognitive-neuroscience-of-music
25. Why Long Electrode Arrays: More Natural Tonotopic Coding, https://blog.medel.pro/products-updates/natural-tonotopic-coding/
26. A Frequency-Position Function for the Human Cochlear Spiral, https://pmc.ncbi.nlm.nih.gov/articles/PMC2432090/
27. Electrocochleography localization of characteristic frequency for, https://patents.google.com/patent/WO2023250095A1/en
28. Lecture 3 a psychoacoustics of discriminating and identifying sounds, https://www.slideshare.net/slideshow/lecture-3-a-psychoacoustics-of-discriminating-and-identifying-sounds/65539976
29. Relating Tuning and Timbre - sethares homepage, https://sethares.engr.wisc.edu/consemi.html
30. Consonance and Dissonance - Carlos Scheidegger, https://cscheid.net/v2/explainers/music/consonance-and-dissonance.html
31. Consonance calculations (1) - alpof - WordPress.com, https://alpof.wordpress.com/2015/04/05/consonance-calculations-1/
32. Consonance and dissonance - the LPTHE, https://www.lpthe.jussieu.fr/~talon/MUSIC4.PDF
33. Musical Microtonal Scales based in the quantitative measurement of, https://www.researchgate.net/publication/220025474_Musical_Microtonal_Scales_based_in_the_quantitative_measurement_of_the_sensory_dissonance_of_Sethares
34. Musical and linguistic syntactic processing in agrammatic aphasia, https://pubmed.ncbi.nlm.nih.gov/35002061/
35. Language influences music harmony perception: effects of shared, https://royalsocietypublishing.org/rsos/article/3/2/150685/36508/Language-influences-music-harmony-perception
36. [PDF] Structural integration in language and music: Evidence for a, https://www.semanticscholar.org/paper/7fe6112c61cc87e441ca9503cc0527b9b3dc1872
37. The relation of Shared Syntactic Integration Resource Hypothesis, https://journals.gold.ac.uk/index.php/sysmus14/article/view/247
38. Anatomically distinct dopamine release during anticipation and, https://pubmed.ncbi.nlm.nih.gov/21217764/
39. Anatomically distinct dopamine release during anticipation and, https://scispace.com/papers/anatomically-distinct-dopamine-release-during-anticipation-1tl2k0rs2k

40. Predictions and the brain: how musical sounds become rewarding, https://dericbownds.net/uploaded_images/salimpoor.pdf
41. Interactions Between the Nucleus Accumbens and Auditory Cortices, https://www.zlab.mcgill.ca/publications/docs/Science-2013-Salimpoor-216-9.pdf
42. The Reward of Musical Emotions and Expectations (Chapter 15), https://www.cambridge.org/core/books/culture-mind-and-brain/reward-of-musical-emotions-and-expectations/479B3A38A397791B7E2CEA693B866B24
43. Research Progress of Music Therapy on Gait Intervention in Patients, https://www.researchgate.net/publication/362481614_Research_Progress_of_Music_Therapy_on_Gait_Intervention_in_Patients_with_Parkinson's_Disease
44. Can neurologic music therapy make the difference when using, https://pmc.ncbi.nlm.nih.gov/articles/PMC12979535/
45. Hot Off the Press - Academy of Neurologic Music Therapy, https://nmtacademy.co/whats-new/
46. AudioCraft 2026 Review: Generative Audio and Music Code Base, https://aijet.cc/item/audiocraft
47. AudioCraft Reviews in 2026 - SourceForge, https://sourceforge.net/software/product/AudioCraft/
48. Discrete Audio Tokens: More Than a Survey! - arXiv, https://arxiv.org/html/2506.10274v3
49. Neurobiological mechanism of music improving gait disorder in, https://pmc.ncbi.nlm.nih.gov/articles/PMC11738927/
50. Universality and diversity in human song, https://coevolution.fas.harvard.edu/publications/universality-and-diversity-human-song

Universality and diversity in human song - PubMed - NIH, https://pubmed.ncbi.nlm.nih.gov/31753969/