

Comparative Biophysics, Thermodynamic Scaling, and Evolutionary Dynamics of Neural Fluid, Epithelial Appendages, and Botanical Meristematic Growth

Author: SAMUELSON G

Independent Researcher

Abstract

The fundamental mechanisms that govern biological growth, fluid dynamics, and cellular metabolism across the eukaryotic domain exhibit profound evolutionary divergence. This comprehensive review explores the distinct biophysical and thermodynamic strategies employed by three radically different biological systems: the human central nervous system, human epithelial hair follicles, and terrestrial botanical meristems. In the human brain, water functions as a precisely calibrated homeostatic and structural medium governed by the Monro-Kellie doctrine, facilitating neuro-metabolic waste clearance via the glymphatic system without driving expansive growth. Conversely, human hair growth relies on deterministic, ATP-intensive cellular proliferation, utilizing specialized metabolic adaptations such as localized lactate shuttling to fuel rapid division. In stark contrast to mammalian systems, the indeterminate growth of terrestrial grasses and roots relies on massive hydrostatic turgor pressure to drive anisotropic cell elongation. By synthesizing hydrodynamics, metabolic opportunity costs, and evolutionary biology—including the transition from determinate to indeterminate growth paradigms—this paper delineates how disparate organisms engineer growth and maintain physiological homeostasis in response to distinct environmental and structural constraints.

Keywords: Biophysics, Glymphatic System, Turgor Pressure, Meristematic Growth, Cellular Proliferation, Evolutionary Scaling, Hair Follicle Metabolism.

1. Introduction

Despite sharing conserved underlying cellular machinery—such as DNA replication, basic organelle function, and lipid bilayer boundaries—the utilization of water, thermodynamic energy, and structural proteins varies dramatically across the eukaryotic domain. The human brain, human epithelial hair follicles, and terrestrial grasses (along with their root systems) represent three radically distinct paradigms of biological organization.

In the human central nervous system, water is not a driver of expansive growth but rather a precisely calibrated homeostatic, structural, and physiological medium essential for neuro-metabolic waste clearance and intracranial pressure regulation.¹ Conversely, human hair growth is characterized by cyclical, division-driven cellular proliferation that relies heavily on localized, highly specialized metabolic adaptations—such as a localized Warburg effect—and is sustained by systemic hydration for nutrient delivery rather than cellular turgidity.³ In stark contrast to both mammalian systems, the growth of terrestrial grasses and their root systems relies fundamentally on turgor pressure—a massive hydrostatic force generated by vacuolar water accumulation—to drive anisotropic cell elongation, coordinated by localized meristematic activity and sophisticated environmental tropisms.⁵

The comprehensive analysis presented herein explores the biophysical, metabolic, and evolutionary differences defining these three distinct biological phenomena.

2. Hydrodynamic Architecture and Fluid Constraints of the Human Brain

The human central nervous system operates within a highly regulated, volume-restricted fluid environment, where macroscopic fluid flow dictates both physiological function and pathological vulnerability. Unlike plant tissues or epithelial appendages, the adult human brain does not undergo macroscopic expansive growth; its fluid dynamics are optimized for metabolic homeostasis.

2.1 The Monro-Kellie Doctrine and Intracranial Pressure Regulation

The biophysics of the human brain is absolutely constrained by the rigid, unyielding structure of the cranium. According to the Monro-Kellie doctrine, the total volume of the intracranial compartment is fixed, consisting primarily of three essentially incompressible components: brain tissue (averaging approximately 1400 mL), cerebral blood volume

(150 mL), and cerebrospinal fluid (CSF) (150 mL).¹ An increase in the volume of one compartment will invariably lead to a compensatory decrease in another, or else result in a rapid rise in intracranial pressure (ICP).¹

Under normal physiological conditions, ICP is maintained at approximately 10 mmHg in a supine individual, and drops to between 0 and 2 mmHg when the individual is in an upright position.¹ This delicate neural balance is highly sensitive to external biochemical interference; for instance, steroidal alkaloids from the *Veratrum* plant species can act as neurotoxic substances, causing DNA damage, decreasing blood pressure, and disrupting the brain's delicate biophysical state.⁸

2.2 The Glymphatic System and State-Dependent Water Dynamics

The brain consists of four primary fluid compartments: the intracellular fluid (ICF), the interstitial fluid (ISF), the cerebrospinal fluid (CSF), and the cerebral vasculature.⁹ The central nervous system completely lacks the conventional lymphatic vessels that perform waste clearance in peripheral tissues.⁹ This paradox is resolved by the glymphatic system, a highly specialized macroscopic waste clearance pathway that facilitates the exchange of CSF and ISF to flush neurotoxins, including β -amyloid, from the brain parenchyma.⁹

CSF flows through the subarachnoid space surrounding the cerebral arteries and is driven into the perivascular spaces.¹⁰ Regulated by Aquaporin-4 (AQP4) water channels, CSF enters the brain parenchyma, mixes with the ISF, and mobilizes interstitial solutes toward the perivenous spaces for systemic clearance.¹⁰ This mechanism exhibits profound, state-dependent temporal variations, becoming highly active during sleep.⁹ During transitions into non-rapid eye movement (NREM) sleep, the volume of the intracellular fluid (ICF) compartment physically contracts, reducing hydraulic resistance and allowing for a massive influx of CSF.¹¹

2.3 Water as a Structural Hydrogel Matrix

Water represents the primary structural component of the brain's soft tissue matrix. The structural organization of water in biological tissues includes bulk (or "free") water and interfacial (or "bound") water, the latter forming a macromolecular network that dictates the structural properties of surrounding proteins and lipids.¹² The brain represents a hydrostatic material whose resistance to mechanical stress is highly dependent on its water content; however, unlike plant tissues, the brain relies entirely on hydration to maintain its viscoelastic properties without the active, cell-wall-constrained turgidity

characteristic of botanical organisms.¹²

3. Cellular Dynamics and Metabolic Cost in Human Hair Proliferation

Human hair growth represents one of the most metabolically demanding and highly proliferative continuous processes in mammalian biology. Hair growth is not driven by hydrostatic inflation but by the deterministic, highly regulated division and differentiation of stem cells residing within the complex microenvironment of the hair follicle.⁴

3.1 Anatomy and Cyclical Dynamics of the Hair Follicle

Hair is formed through the rapid division and differentiation of stem cells, producing keratinocytes that migrate, flatten, and die to become the non-living, keratinized epithelial cells of the visible hair shaft.⁴ The growth originates in the hair bulb, located at the proximal end of the follicle.¹³ Below the Auber line lies the matrix, a germination center composed of highly mitotically active cells.⁴

Hair growth is cyclical, alternating between periods of rapid elongation and quiescence. The cycle is divided into four primary phases:

1. **Anagen (Growth Phase):** The highly mitotic active phase where the follicle works continuously to produce the hair fiber.⁴ It is subdivided into proanagen (proliferation and differentiation) and metanagen (emergence of the shaft).⁴ Scalp hair grows approximately 0.35 mm/day or 1 cm/month.⁴
2. **Catagen (Transition Phase):** A short transition stage marking the cessation of cellular proliferation.¹⁴ The hair follicle undergoes apoptosis-driven regression, losing about one-sixth of its standard diameter to form a "club hair".⁴
3. **Telogen (Resting Phase):** The follicle remains dormant to conserve energy.¹⁴ Approximately 10 to 15% of all body hairs reside in this resting stage at any given time.⁴
4. **Exogen (Shedding Phase):** The mechanical release and shedding of the fully formed club hair.¹⁵

3.2 Metabolic Compartmentalization and the Lactate Shuttle

The proliferation of matrix cells cannot rely solely on conventional oxidative phosphorylation due to the sheer speed of required biomass accumulation. The human hair follicle operates under a spatiotemporally controlled metabolic paradigm akin to the Warburg effect.³ Epithelial stem cell quiescence in the bulge region is maintained through

distinct metabolic signatures, whereas the rapid proliferation of keratinocytes demands alternative energy routing.³

As hair follicle stem cells consume glucose, they process it into pyruvate. Preventing this pyruvate from entering the mitochondria and forcing its conversion into lactate rapidly activates quiescent stem cells.¹⁶ Glycolytic GLUT1+ keratinocytes located in the lower outer root sheath (ORS) predominantly produce this lactate, which is shuttled to neighboring, highly proliferative matrix keratinocytes via MCT1-mediated uptake to fuel oxidative metabolism.³

3.3 Microenvironmental Stress and Integrated Stress Responses

Energy metabolism in the hair follicle must be highly adaptive to systemic stress. The Mitochondrial Pyruvate Carrier (MPC) imports glycolysis-derived pyruvate into the mitochondria.¹⁷ Pharmacologically targeting the MPC in human hair follicles with inhibitors like UK-5099 induces a metabolic stress-responsive proliferative arrest.¹⁷ This inhibition activates the Integrated Stress Response (ISR), mediated by the transcription factor ATF4, which responds to stressors by halting cell cycle progression.¹⁷ Activation of the ISR results in the upregulation of ATF4-target genes like ADM2, which imposes a proliferative block. Treatment with ISR inhibitors (e.g., ISRIB) can attenuate this upregulation, highlighting the delicate energetic balance required to maintain anagen.¹⁷

3.4 Hydration versus Turgor in Mammalian Epithelia

Unlike botanical systems, human hair expansion relies on ATP-intensive cell division. Water does not act as a turgor-generating force within the hair shaft.⁵ However, systemic water remains critical. Proper hydration ensures adequate blood plasma volume to deliver oxygen, vitamins, and minerals to the follicle.¹⁸ Clinical assessments of dehydration often rely on testing skin turgor; mild dehydration causes the skin to return slowly to its normal flat state, reflecting a loss of internal moisture retention and elasticity, presenting as "doughy skin".¹⁹ Furthermore, adequate internal hydration helps regulate sebum production, maintaining the hydrolipidic film that protects the hair shaft from desiccation.²¹

4. Botanical Meristematic Growth and Hydrostatic Expansion

In extreme evolutionary contrast to the ATP-intensive growth of human tissue, terrestrial grasses (family Poaceae) rely on continuous cell generation at localized meristems,

followed by massive, turgor-driven volumetric expansion.⁵

4.1 Shoot Architecture, Intercalary Meristems, and Photosynthesis

Grasses employ a unique dual-meristem system: the apical meristem and the intercalary meristem.²³ Apical meristems push new leaves upward, while intercalary meristems (located at the bases of leaf blades and nodes) allow continuous basal elongation.²³ This adaptation allows grass leaves to elongate even after repeated defoliation by herbivores or mowing.²³

The metabolic engines driving grass growth differ based on ecological adaptation, categorized into C3 and C4 photosynthetic pathways.²⁴ C3 grasses (cool-season) perform optimally in shaded environments.²⁵ C4 grasses (warm-season) evolved an additional pathway that initially fixes CO₂ into a 4-carbon molecule, drastically increasing Nitrogen Use Efficiency (NUE) and Water Use Efficiency (WUE) during high-heat and high-light conditions.²⁴

4.2 Root Dynamics, Tropisms, and the ROOTFLOW Paradigm

Roots anchor the plant, extract water, and synthesize phytohormones.²⁶ The root tip is divided into the zone of cell division (meristem), the zone of elongation, and the zone of maturation.⁷ The ROOTFLOW project revealed that the boundary where cells transition from meristematic division to rapid elongation oscillates in a negative feedback loop with a periodicity of about 90 minutes and over a distance of 50 μm .²⁷

Root growth is guided by directional tropisms. Gravitropism (growth toward gravity) relies on an auxin gradient; starch-filled amyloplasts settle at the bottom of root cap cells, triggering an "auxin fountain" that redistributes auxin to the elongation zone, inhibiting growth on the lower flank to bend the root downward.²⁸ Conversely, hydrotropism steers roots toward moisture gradients via abscisic acid (ABA) signaling.³⁰ While reactive oxygen species (ROS) are required for gravitropic bending, they actively inhibit hydrotropism.³⁰

4.3 Turgor Pressure Biophysics and Dimensional Analysis

Plant cells are encased in a rigid cellulosic cell wall; therefore, cellular division alone only partitions existing space.³² True volumetric growth requires expansion driven by turgor pressure (hydrostatic pressure), generated by the osmotic flow of water into the central vacuole.⁵

Elongation requires anisotropic growth (expansion directed preferentially along a single

axis), governed by localized cell wall yielding.³² In root hairs, expansion is restricted entirely to the extreme apex via *tip growth*, utilizing specialized cell wall-loosening expansins such as EXPA7 and EXPA18. This tubular morphology represents a striking evolutionary convergence with mammalian intestinal villi, as both independently evolved to vastly enlarge surface area for nutrient absorption.

Biophysical equations utilizing a dimensionless number, Π_{pw} , reveal that both turgor pressure (P) and expansive growth rate (v) decrease curvilinearly as environmental water deficit (the magnitude of Π_{pw}) becomes more severe.³³

4.4 Systemic Water Transport Mechanics

The ascent of sap through the xylem is predominantly governed by the Cohesion-Tension Theory.³⁴ Evaporation from stomata creates massive negative pressure (tension) within the leaf apoplast, transmitted continuously down the rigid xylem column to pull water upward against gravity.³⁴ At night, active ion transport into the stele generates "root pressure" (a positive hydrostatic force of approx. 5-6 bars or 75-90 psi), which, while notable, is mathematically inadequate for lifting water to the canopies of tall trees without transpirational pull.³⁴

5. Comparative Thermodynamic and Metabolic Scaling

Metabolism encompasses the interconnected pathways of anabolism and catabolism required to supply the cell with ATP.³⁵ The energetic strategies underlying mammalian cellular proliferation versus botanical cellular expansion scale radically differently.

5.1 ATP Allocation Profiles and Thermodynamic Cost

Biological Process	Primary Growth Mechanism	Metabolic Driver / ATP Sink	Structural Substrate
Human Brain Homeostasis	Zero macroscopic growth; strict volume	Continuous high metabolic rate (TCA cycle) for	Viscoelastic hydrogel matrix; bound water

	maintenance. ¹	neural firing; glymphatic fluid shifts. ⁹	networks. ¹²
Human Hair Proliferation	Rapid, continuous mitotic cellular division (Anagen). ⁴	Warburg effect; lactate shuttling (GLUT1 to MCT1); high anabolism for keratin synthesis. ³	Proteinaceous (Keratin) synthesized entirely from raw precursor metabolites. ⁴
Terrestrial Grass Elongation	Vacuolar inflation; targeted cell wall yielding. ³²	Plasma membrane H ⁺ -ATPases; structural carbohydrate synthesis (cellulose, pectin). ²⁷	Environmental water driven by osmotic potential. ⁵

In the mammalian hair matrix, elongation requires the continuous biochemical synthesis of new cellular entities.⁴ The direct metabolic cost of synthesizing a molecular building block involves significant ATP hydrolysis.³⁹ Measurements utilizing suspended microchannel resonators demonstrate that translation rates and biomass accumulation in prophase are commensurate with, or even higher than, interphase growth rates.⁴⁰ The construction of structural proteins like keratin diverts metabolic precursors away from ATP generation, establishing a massive opportunity cost.³⁹

In contrast, plant cell elongation circumvents this prohibitive opportunity cost by substituting free environmental water for dense protoplasm.⁴² During elongation, the plant vastly expands its central vacuole.⁵ The primary metabolic cost is restricted to the synthesis of cell wall components and the operation of plasma membrane H⁺-ATPases. Thus, botanical expansive growth represents a thermodynamically "cheap" inflation mechanism compared to the heavy anabolism required to build mammalian tissue structure.⁴²

6. Evolutionary Divergence: Deterministic vs. Indeterminate Architectures

The distinct growth behaviors across these systems reflect adaptations categorized into deterministic and indeterminate biological architectures.⁶

Neural Determinism: The human brain represents deterministic growth. Once adult volume is reached, the tissue shifts completely to maintenance, preventing pathological metabolic scaling issues within the fixed cranial vault.¹

Botanical Indeterminacy: Land grasses exhibit indeterminate growth; they continue to add new organs indefinitely via meristems, provided resources are available.⁶ This architectural plasticity is an evolutionary necessity for sessile organisms, allowing them to continually forage through their environment and regenerate foliage after defoliation.⁶

The Evolutionary Anomaly of Human Scalp Hair: While individual follicles undergo a deterministic cycle, the anagen phase on the human scalp has become wildly protracted. In most mammals, hair stops growing at a genetically predetermined length.⁴³ Charles Darwin postulated that the indeterminate growth of human scalp hair was sexually selected, serving as an adornment and an indicator of genetic fitness once early humans developed knife-blade tools to trim it.⁴⁴

7. Conclusions

A comparative examination of human brain hydrodynamics, human hair proliferation, and terrestrial grass growth reveals the extraordinary versatility of biological mechanisms. Within the human central nervous system, water operates as a strictly constrained, incompressible fluid governed by the Monro-Kellie doctrine, crucial for waste clearance but divorced from expansive growth. In human hair, elongation is a product of raw biosynthesis and intense metabolic activity driven by unique bioenergetic pathways like the localized lactate shuttle. Conversely, terrestrial grasses merge environmental water directly with the physical mechanics of growth, weaponizing hydrostatic turgor pressure to execute rapid, thermodynamically efficient, indeterminate tissue elongation. These divergent mechanisms showcase the fundamental distinction between mammalian metabolic engineering—which prioritizes complex homeostasis and high-cost biosynthesis—and botanical hydraulic engineering, optimized for rapid environmental colonization.

Works cited

1. Regulation of intracranial pressure - Deranged Physiology, <https://derangedphysiology.com/main/cicm-primary-exam/nervous-system/Chapter-112/regulation-intracranial-pressure>
2. The Glymphatic System in Cerebrospinal Fluid Dynamics: Clinical Implications, Its Evaluation, and Application to Therapeutics - PubMed, <https://pubmed.ncbi.nlm.nih.gov/40367923/>
3. Compartmentalised metabolic programmes in human anagen hair follicles: New targets to modulate epithelial stem cell behaviour, keratinocyte proliferation and hair follicle immune status? - PubMed, <https://pubmed.ncbi.nlm.nih.gov/33548088/>
4. Physiology, Hair - StatPearls - NCBI Bookshelf, <https://www.ncbi.nlm.nih.gov/books/NBK499948/>
5. Turgor pressure - Wikipedia, https://en.wikipedia.org/wiki/Turgor_pressure
6. Plant Development II: Primary and Secondary Growth | Organismal Biology, <https://organismalbio.biosci.gatech.edu/growth-and-reproduction/plant-development-ii-primary-and-secondary-growth/>
7. 35.3.1: Types of Root Systems and Zones of Growth - Biology LibreTexts, https://bio.libretexts.org/Bookshelves/Introductory_and_General_Biology/Map%3A_A_Raven_Biology_12th_Edition/35%3A_Plant_Form/35.03%3A_Roots-_Anchoring_and_Absorption_Structures/35.3.1%3A__Types_of_Root_Systems_and_Zones_of_Growth
8. Plants' Impact on the Human Brain—Exploring the Neuroprotective and Neurotoxic Potential of Plants - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11510325/>
9. The Glymphatic System – A Beginner's Guide - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4636982/>
10. Glymphatic system: a self-purification circulation in brain - Frontiers, <https://www.frontiersin.org/journals/cellular-neuroscience/articles/10.3389/fncel.2025.1528995/full>
11. Brain water dynamics across sleep stages measured by near-infrared spectroscopy: Implications for glymphatic function - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12202386/>
12. Water Content of Plant Tissues: So Simple That Almost Forgotten? - MDPI, <https://www.mdpi.com/2223-7747/12/6/1238>
13. Hair follicle - Wikipedia, https://en.wikipedia.org/wiki/Hair_follicle
14. Morphogenesis, Growth Cycle and Molecular Regulation of Hair Follicles - Frontiers, <https://www.frontiersin.org/journals/cell-and-developmental-biology/articles/10.3389/fcell.2022.899095/full>
15. The Hair Growth Cycle: 4 Stages Explained | Good Health by Hims, <https://www.hims.com/blog/everything-you-need-to-know-about-the-hair-growth-pr>

- [rocess-regrowing-hair](#)
16. UCLA scientists identify a new way to activate stem cells to make hair grow,
<https://newsroom.ucla.edu/releases/ucla-scientists-identify-a-new-way-to-activate-stem-cells-to-make-hair-grow>
 17. Activation of the integrated stress response in human hair follicles - Research journals - PLOS,
<https://journals.plos.org/plosone/article/file?type=printable&id=10.1371/journal.pone.0303742>
 18. The Importance of Hydration for Hair - How Drinking Water Affects Hair - RKT Hair Care,
<https://www.rkthaircare.com/blogs/wellness/the-importance-of-hydration-for-hair>
 19. Definition of Skin Turgor | BHSkin Dermatology,
<https://ai.bhskin.com/glossary/skin-turgor-dermatology-explained>
 20. Skin turgor: MedlinePlus Medical Encyclopedia,
<https://medlineplus.gov/ency/article/003281.htm>
 21. Does Drinking Water Help Hair Growth?: Role of Hydration - Theradome,
<https://theradome.com/blogs/blog/does-drinking-water-help-hair-growth>
 22. What is hair hydration: myths and reality - Seppic,
<https://www.seppic.com/en-US/article/hair-hydration-myths-reality>
 23. Mechanisms for Growth | Forage Information System | Oregon State ...,
<https://forages.oregonstate.edu/regrowth/how-does-grass-grow/grass-structures/mechanisms-growth>
 24. What are C3 and C4 Native Grass? - NSW Department of Primary Industries,
<https://www.dpi.nsw.gov.au/agriculture/pastures-and-rangelands/native-pastures/what-are-c3-and-c4-native-grass>
 25. Guide to C3 & C4 Grasses: Growth, Photosynthesis, and Turf Management - AgriGro, <https://agrigro.com/blogs/blog/understanding-c3-c4-grasses>
 26. Root health drives grass growth | Hay and Forage Magazine,
<https://hayandforage.com/article-4416-Root-health-drives-grass-growth.html>
 27. How do plants coordinate root growth? | ROOTFLOW Project | Results in Brief - CORDIS,
<https://cordis.europa.eu/article/id/181139-how-do-plants-coordinate-root-growth>
 28. BOTANY (PLANTS) / TROPISMS - Pathwayz,
<https://www.pathwayz.org/Tree/Plain/TROPISMS>
 29. Root Tropisms: Investigations on Earth and in Space to Unravel Plant Growth Direction,
<https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2019.01807/full>
 30. Growth-mediated plant movements: hidden in plain sight Short title ...,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC5826749/>
 31. How Plants Move - Penn State Extension,

- <https://extension.psu.edu/how-plants-move>
32. The dynamics and biophysics of shape formation: common themes in plant and animal morphogenesis - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10752614/>
 33. Theoretical Analyses of Turgor Pressure and Expansive Growth Rate of Plant Cells During Water Deficit - MDPI, <https://www.mdpi.com/2223-7747/14/22/3538>
 34. Lecture 9 Water Relations of Plants,
<https://www.appstate.edu/~neufeldhs/bio1102/lectures/lecture9.htm>
 35. Metabolism - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7545035/>
 36. Energy metabolism and ATP balance in animal cell cultivation using a stoichiometrically based reaction network - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/18629932/>
 37. Integrative and Mechanistic Approach to the Hair Growth Cycle and Hair Loss - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC9917549/>
 38. Root hair growth from the pH point of view - Frontiers,
<https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2022.949672/full>
 39. Defining the Energetic Costs of Cellular Structures | bioRxiv,
<https://www.biorxiv.org/content/10.1101/666040.full>
 40. Mammalian cell growth dynamics in mitosis - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC6534395/>
 41. Mammalian cell growth dynamics in mitosis - eLife,
<https://elifesciences.org/articles/44700>
 42. How Metabolic Rate Relates to Cell Size - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9332559/>
 43. Indeterminate Growth: Could It Represent the Ancestral Condition? - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4743077/>
 44. Why does animal hair has a maximum length whereas human hair can grow indefinitely?,
https://www.reddit.com/r/askscience/comments/3bmr0v/why_does_animal_hair_has_a_maximum_length_whereas/