

Comprehensive Analysis of Neoplastic Disease: Etiology, Clinical Manifestations, Surgical Interventions, and Therapeutic Paradigms

Author: SAMUELSON G, Independent Researcher

Abstract

Cancer remains a leading global health challenge, with conventional diagnostic and treatment methods often lacking precision and adaptability. This comprehensive review explores transformative advancements that are reshaping oncology, emphasizing the integration of biomarker-driven diagnostics with tailored therapeutic strategies. It examines the complex etiology of neoplastic disease, the systemic patient experience, and evolving surgical paradigms ranging from minimally invasive robotics to complex cytoreductive procedures. Additionally, it highlights the critical role of artificial intelligence as a supportive, evidence-organizing co-pilot within multidisciplinary tumor boards. Furthermore, the review outlines the far-reaching consequences of cancer for health systems, economies, and societies, highlighting the urgent need to address equitable access to life-saving technologies and care worldwide.¹ By adopting cutting-edge, patient-centered approaches, a new era of precision oncology emerges, bridging the gap between early detection and effective intervention.

Keywords: Oncology, Precision Medicine, Surgical Oncology, Patient-Centered Outcomes Research, Artificial Intelligence, Health Equity.

Introduction

The global burden of neoplastic diseases represents one of the most formidable and complex challenges to modern medical science, healthcare infrastructure, and socioeconomic stability worldwide. In 2022 alone, the global incidence of cancer reached an estimated 20 million new cases, accompanied by 9.7 million disease-related deaths,

leaving approximately 53.5 million individuals alive within five years of their initial diagnosis.² With epidemiological models projecting an escalation to over 35 million new cases annually by the year 2050—driven by demographic aging, exponential population growth, and escalating exposure to environmental risk factors—the necessity for continuous, multifaceted oncological research is absolute.²

The boundaries of this comprehensive analysis encompass the full spectrum of oncological science, integrating microscopic cellular biology with macroscopic epidemiological data. This report examines the genetic and epigenetic aberrations that initiate carcinogenesis, the complex systemic and paraneoplastic symptoms that herald the disease, the evolving paradigms of surgical oncology ranging from radical resections to minimally invasive robotic procedures, and the profound ethical dimensions governing patient consent and clinical trials. By synthesizing quantitative survival data with qualitative patient-reported outcomes, a holistic understanding of cancer care emerges—one that prioritizes not only the eradication of malignant cells but also the holistic preservation of the human patient.

The fundamental utility of this vast continuum of research lies in its dual capacity. Basic research decodes the foundational molecular anomalies that trigger malignant transformation, while applied research translates these biological insights into tangible clinical interventions, advanced surgical procedures, and targeted pharmacotherapies. Ultimately, this translational pipeline serves the public by extending survival, minimizing treatment-related morbidity, and enhancing the post-operative quality of life. Simultaneously, it provides the medical community with evidence-based frameworks, clinical decision-making algorithms, and sophisticated risk assessment tools required to navigate the inherent complexities of modern cancer care. By continuously pushing the boundaries of what is scientifically possible, oncological research provides an indispensable utility to all of society, transforming universally fatal diagnoses into manageable chronic conditions and, increasingly, definitive cures.

Etiology and Pathogenesis of Neoplasms

The genesis of a tumor is rarely a singular, isolated event; rather, it is the culmination of highly complex, dynamic interactions between genetic predispositions, epigenetic modifications, and environmental stimuli. Understanding the biological causes of tumors requires an exhaustive analysis of the cellular microenvironment and the specific

molecular pathways that govern cell proliferation, survival, and cellular metabolism.

Clonal Evolution and Intra-Tumor Heterogeneity

With rare exceptions, spontaneous human tumors originate from a single aberrant cell.³ However, by the time a tumor expands sufficiently to reach the threshold of clinical detection, it is typically characterized by startling intra-tumor heterogeneity.³ This phenotypic and genotypic diversity is the direct result of a Darwinian evolutionary process occurring continuously within the tumor microenvironment.³ As malignant cells proliferate, their inherent genetic instability leads to the continuous generation of mutant clones.³ Environmental and systemic selective pressures—such as localized hypoxia, nutrient deprivation, and the introduction of therapeutic interventions—favor the survival of specific clones equipped with advantageous traits. These traits may manifest as altered cell surface receptor expression, enhanced cellular motility, or upregulated angiogenic potential.³ This profound evolutionary divergence between primary tumors and metastatic outgrowths poses a significant barrier to systemic therapies. Treatments that successfully eradicate the dominant cellular clone may inadvertently apply selective pressure that allows a previously minor, chemoresistant sub-clone to flourish and repopulate the tumor mass.³

Environmental Drivers and Carcinogenesis

Carcinogenesis is heavily influenced by environmental mutagens, commonly referred to in basic research as initiating agents.⁴ These agents induce the initial, critical mutations in key target genes, particularly oncogenes and tumor suppressor genes, setting the stage for malignant transformation.⁴ Prominent examples include solar ultraviolet radiation, which acts as a primary driver of cutaneous malignancies; aflatoxins, which are potent hepatic carcinogens produced by contaminating molds on grains; and the multitude of chemical carcinogens present in tobacco smoke, such as benzo(a)pyrene, dimethylnitrosamine, and nickel.⁴

Beyond direct DNA damage, chronic inflammation has been definitively recognized as a core hallmark of cancer.⁵ Both genetic and environmental factors contribute to tumor-promoting inflammation, which supplies the microenvironment with bioactive molecules, growth factors, and survival signals that inadvertently support neoplastic progression and protect nascent tumors from immunological destruction.⁵ Normal cellular

proliferation is tightly regulated by the availability of extracellular polypeptide growth factors and phenomena such as density-dependent inhibition, wherein healthy cells cease dividing when they come into contact with one another.⁴ Conversely, a defining characteristic of transformed cells is their drastically reduced reliance on these external signals, allowing for unchecked, autonomous proliferation.⁴ As a tumor expands, it inevitably outgrows its local vascular supply, leading to central regions of severe hypoxia and tissue necrosis.⁵ To survive, the tumor must trigger the "angiogenic switch," actively secreting factors to stimulate the formation of new, albeit structurally abnormal and leaky, blood vessels.⁵ Concurrently, cancer cells undergo significant metabolic rewiring—often termed the Warburg effect—shifting energy metabolism away from oxidative phosphorylation toward aerobic glycolysis to support rapid macromolecular synthesis and continued survival even in nutrient-poor conditions.⁵

Key Molecular Signaling Pathways: PI3K/AKT/mTOR and MAPK

At the fundamental molecular level, tumor progression is frequently driven by the severe dysregulation of highly conserved intracellular signaling cascades. Basic research has identified the PI3K/AKT/mTOR pathway as a pivotal regulator of cellular viability, metabolism, and proliferation.⁶ The activation sequence begins with the auto-phosphorylation of Phosphoinositide 3-kinase (PI3K), which subsequently catalyzes the conversion of PIP2 to PIP3 at the cellular membrane.⁷ This critical biochemical event recruits and activates AKT (Protein Kinase B), leading to a massive cascade of downstream phosphorylation events that aggressively promote tumor growth, inhibit apoptosis, and suppress protective autophagy.⁷

Mutations or amplifications in upstream receptor tyrosine kinases—such as KIT, ERBB2 (HER2), PDGFRA, MET, and various Fibroblast Growth Factor Receptors (FGFR2, FGFR3)—frequently lead to the constitutive, unchecked activation of this pathway across various human cancers.⁶ Furthermore, the loss or mutational inactivation of the tumor suppressor gene PTEN, a phosphatase that negatively regulates the cascade by converting PIP3 back to PIP2, is a remarkably frequent event in multiple advanced malignancies.⁷

The PI3K/AKT/mTOR axis does not operate in biological isolation; it heavily interacts with other critical cascades, including the Mitogen-Activated Protein Kinase (MAPK/ERK) pathway, as well as NOTCH, Androgen Receptor (AR), WNT/ β -catenin, and TGF- β signaling networks.⁸ This extensive molecular cross-talk amplifies cancer

aggressiveness and frequently underpins the development of resistance to targeted monotherapies.⁸ Furthermore, basic research indicates that specific microRNAs, such as miR27a3p and miR520h, play profound roles in modulating these pathways, driving chemoresistance to agents like paclitaxel by actively attenuating the stability of tumor suppressors like PTEN and hyperactivating AKT.⁷ Elucidating these dominant signaling proteins accounts for nearly 90% of the functional dynamic within the PI3K/AKT pathway, making them the most critical therapeutic targets for modern precision oncology and applied pharmacological research.⁶

Clinical Manifestations and the Patient Experience

The clinical presentation of cancer extends far beyond the localized structural and mechanical impact of the primary tumor mass. The disease exerts profound systemic, neurological, and psychological effects, necessitating a comprehensive approach to patient assessment that captures both measurable, objective clinical signs and the deeply qualitative, lived experience of the patient.

Systemic and Paraneoplastic Syndromes

While local tumor growth can cause mechanical obstruction, hemorrhage, or acute pain, a highly specific subset of patients develops Paraneoplastic Syndromes (PNS).¹⁰ These rare but highly complex systemic disorders are not caused by direct invasion, local tissue destruction, or metastasis.¹⁰ Instead, they are secondary consequences of the body's profoundly altered immune response to an occult malignancy or the ectopic production of active biological substances by the tumor itself.¹⁰ Tumor cells may spontaneously generate autoantibodies, cytokines, hormones, or biologically active peptides that cross-react or interfere with distant organ systems, leading to severe dermatologic, rheumatologic, endocrine, hematologic, and neurologic manifestations.¹¹

Paraneoplastic Neurologic Syndromes (PNS) are particularly devastating and complex, often presenting clinically months or even years before the underlying malignancy reaches a size detectable by standard radiological imaging.¹⁰ Due to rapid advances in neuroimmunology since the original 2004 diagnostic criteria, the contemporary diagnosis of PNS requires an understanding of clinical, immunologic, and oncologic heterogeneity.¹² The diagnostic framework has evolved to classify specific clinical presentations as "high-risk phenotypes" or "intermediate-risk phenotypes" based on the

presence of specific circulating onconeural autoantibodies.¹² The detection of high-risk antibodies, defined as having a greater than 70% association with occult cancer, is a critical diagnostic event that mandates an immediate and exhaustive systemic search for a hidden tumor.¹² These syndromes are most commonly associated with small cell lung cancer, breast cancer, gynecologic malignancies, and specific hematologic neoplasms.¹¹ Timely recognition of paraneoplastic symptoms serves as a vital clinical warning system, facilitating early oncological intervention that can potentially lead to the complete eradication of an otherwise occult tumor, while simultaneously resolving the debilitating autoimmune or endocrine derangements.¹¹

Qualitative Data: Patient-Centered Outcomes Research (PCOR)

The true, holistic burden of cancer cannot be fully quantified by survival metrics, progression-free intervals, or radiological response rates alone. Qualitative surveys and in-depth thematic analyses are strictly required to understand the profound disruptions to the patient's lived experience. The field of Patient-Centered Outcomes Research (PCOR) has revolutionized oncology by prioritizing resources and methodologies to evaluate variables that matter most directly to patients, such as functional status, long-term symptom relief, and psychological well-being.¹³ PCOR fundamentally shifts the research paradigm by engaging patients, caregivers, and clinicians to determine which clinical trial endpoints are truly meaningful, transforming qualitative interview data into actionable information to guide healthcare choices.¹³

To rigorously assess these qualitative dimensions, researchers and clinicians heavily rely on validated instruments, with the EORTC QLQ-C30 (European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire) acting as the global standard.¹⁶ This 30-item core questionnaire evaluates physical, role, emotional, cognitive, and social functioning, alongside tracking specific clinical symptoms like fatigue, nausea, pain, dyspnea, insomnia, loss of appetite, and severe financial difficulties.¹⁶ The instrument is highly modular, allowing clinicians to pair the core questions with disease-specific modules (e.g., assessing radiation proctitis or cachexia).¹⁶ Quantitative tracking of these qualitative metrics demonstrates that significant changes in QLQ-C30 scores directly correlate with the severity of post-operative complications, overall rehabilitation trajectories, and the length of hospital stays, proving the instrument's clinical utility beyond mere subjective measurement.¹⁹

Qualitative studies investigating cancer survivorship post-surgery reveal several

recurring, highly distressing themes. Following active treatment, patients frequently experience what is described in qualitative literature as an "emotional fallout" episode.²¹ This phase is characterized by a severe loss of direction and control, stemming from the sudden discontinuation of intense medical monitoring and a perceived decrease in structural support from healthcare professionals and immediate social networks.²¹ Survivors of colorectal and rectal cancers report major long-term impacts on their identity, self-confidence, physical intimacy, and mental health, predominantly manifesting as chronic, generalized anxiety and a profound, paralyzing fear of cancer recurrence.²² The emotional trauma is frequently exacerbated by instances of aborted cancer surgeries, where procedures are halted intraoperatively due to unexpected disease progression. Qualitative studies have shown that aborted surgeries profoundly impact multiple domains of quality of life—physical, emotional, social, and existential—leaving patients to navigate shattered hope and requiring highly specialized, patient-centered interventions.²⁴

Furthermore, massive surgical interventions—such as radical mastectomies, head and neck resections, or the creation of permanent ostomies—profoundly alter bodily architecture and self-image, requiring specialized psychological interventions focused on self-compassion to help patients reconstruct a positive relationship with their physical selves.²⁵ Financial toxicity emerges as another critical sub-theme in qualitative research; the immense out-of-pocket costs of novel therapeutics, coupled with potential unplanned early retirement and massive productivity loss, create immense psychosocial stress that significantly impedes both physical and emotional recovery.²³ Finally, qualitative analysis reveals that patients face a critical deficit in professional support tailored to navigating the return-to-work process, with many reporting that their employers and peers demonstrate a severely limited understanding of the long-term cognitive and physical impacts of a cancer diagnosis and its grueling treatment.²⁶

Quantitative Data: Global Epidemiology, Survival Trends, and Economics

A comprehensive understanding of oncology mandates an exhaustive review of quantitative, macro-level data, which defines the global scale of the disease, the historical efficacy of evolving interventions, and the immense economic strain placed on global health systems.

Global Incidence and Mortality Statistics

In 2022, the global cancer burden reached staggering, unprecedented proportions. Data consolidated by the World Health Organization (WHO) and the International Agency for Research on Cancer (IARC) estimate that approximately 1 in 5 individuals will develop a malignancy during their lifetime, with 1 in 9 men and 1 in 12 women ultimately dying from the disease.² Overall, there were 20 million new cancer cases globally in 2022, accompanied by 9.7 million deaths.² Looking closely at the United States, in 2024, the American Cancer Society projected 2,001,140 new cancer cases and 611,720 cancer deaths.²⁷

Leading Global Cancer Types (2022 Incidence)	Estimated New Cases	Percentage of Total Global Cases
Lung Cancer	2.5 million	12.4%
Female Breast Cancer	2.3 million	11.6%
Colorectal Cancer	1.9 million	9.6%
Prostate Cancer	1.5 million	7.3%
Stomach Cancer	970,000	4.9%

2

Mortality figures parallel these incidence rates, heavily dictated by the biology of the tumor and the availability of early screening. Lung cancer remains the dominant cause of cancer-related death, claiming 1.8 million lives globally (18.7% of total cancer deaths) in

2022, followed by colorectal cancer (900,000 deaths), liver cancer (760,000 deaths), breast cancer (670,000 deaths), and stomach cancer (660,000 deaths).² The global trajectory is highly concerning; assuming current incidence rates remain unchanged, the projected cases are expected to surge by 77% to over 35 million annually by the year 2050.² This dramatic rise is driven almost entirely by the aging of the global population, general population growth, and the rising prevalence of major risk factors such as tobacco usage, harmful alcohol consumption, unhealthy diets, air pollution, and obesity, particularly in emerging economies.²

Disparities in Survival and Treatment Efficacy

Treatment efficacy and overall survival rates are heavily stratified by geographic location and the Human Development Index (HDI) of the specific region. While very high-HDI nations experience higher raw incidence rates due to comprehensive screening and longer life expectancies, lower-HDI countries suffer drastically disproportionate mortality rates.² For instance, in very high HDI countries, 1 in 12 women are diagnosed with breast cancer, and 1 in 71 die; conversely, in low HDI countries, while only 1 in 27 women are diagnosed, a staggering 1 in 48 die.² The breast cancer mortality rate is twice as high in Ethiopia as in the United States, despite a 60% lower incidence rate, a direct consequence of inadequate screening infrastructure, delayed diagnostics, and limited access to multimodal therapies.²⁸

Within high-resource settings, long-term epidemiological data from the National Cancer Institute's Surveillance, Epidemiology, and End Results (SEER) program highlight significant, though uneven, improvements in relative survival rates, directly correlating with the advent of advanced therapeutics and refined surgical techniques. Relative survival adjusts for normal life expectancy by comparing survival among people who have cancer with that of the general population, controlling for age, race, sex, and year.³⁰

SEER 8 Data Year Range	Rate of New Cases (per 100,000)	5-Year Relative Survival
1988 - 1990	457.76 - 473.95	56.44% - 58.66%

2013 - 2015	448.37 - 450.78	70.42% - 71.61%
2020 - 2022 (Modeled)	411.63 - 446.87	73.77% - 74.63%

32

Between the years 2000 and 2015, specific malignancies saw dramatic survival increases. The 5-year relative survival rate for leukemia improved to 72.53%, lymphoma jumped to 93.38%, and endocrine system cancers reached near-curative levels at 98.98%.³³ However, survival is highly contingent on tumor stage and grade at the time of diagnosis. For prostate cancer, a highly differentiated Grade 1 tumor offers a 60% 5-year survival rate natively, whereas a Grade 4 tumor plummets to 5%; similarly, stage A2 occult cancer boasts a >90% survival rate, while stage D (metastatic) falls to 21%.³⁴ Despite averting over 4 million deaths since 1991 in the United States due to reductions in smoking and superior therapies, progress is actively threatened by a 1% to 2% annual increase in early-onset colorectal cancers in adults under 55, and in cervical cancer among women 30-44, alongside rising incidences of pancreatic and uterine corpus cancers.²⁷

Societal and Economic Burden

The global macroeconomic impact of cancer is monumental and escalating rapidly. Health economics projections indicate that the global economic burden from 2020 to 2050 will reach an extraordinary \$25.2 trillion in 2017 international dollars.³⁵ This staggering figure is not solely derived from direct healthcare expenditures, such as the exponentially rising costs of novel biologics, extended hospitalizations, and specialized surgical equipment.³⁵ It is massively compounded by indirect costs, primarily the permanent loss of workforce productivity due to premature mortality, morbidity-related absences, and the severe, often unquantified economic strain placed on informal caregivers who must abandon employment to provide home care.¹

The highest absolute economic burdens are currently concentrated in populous nations with high healthcare spending like China, the United States, and India.³⁵ Conversely, countries like Bulgaria, Monaco, and Montenegro face the highest economic burdens relative to their total Gross Domestic Product (GDP).³⁵ Effectively tackling modifiable risk factors and investing in robust, preventative cancer care policies—such as universal

coverage for human papillomavirus (HPV) vaccination programs to eliminate cervical cancer, aggressive tobacco control, and enhanced environmental regulations—would significantly lower long-term health expenditures, improve road safety as a co-benefit of reduced alcohol use, and recuperate trillions in lost economic productivity.¹

Applied Research: Evolving Cures and Systemic Therapeutics

While surgical extirpation remains optimal for localized disease, systemic therapies are absolutely essential to address microscopic metastases and disseminated hematologic cancers. Applied research in 2024 and 2025 has witnessed explosive advancements in molecularly guided treatments, pushing the boundaries of survival far beyond the blunt mechanisms of traditional cytotoxic chemotherapy.

Immunotherapy and CAR-T Cell Innovations

Immunotherapy seeks to completely dismantle the complex biological mechanisms by which cancer cells evade the host's innate and adaptive immune surveillance. Among the most revolutionary modalities in this space is Chimeric Antigen Receptor (CAR) T-cell therapy. This bespoke, highly personalized treatment involves harvesting a patient's own T-cells via leukapheresis, genetically engineering them ex vivo using viral vectors or CRISPR technology to express synthetic receptors that flawlessly recognize specific tumor-associated antigens, and subsequently infusing the multiplied cells back into the patient.³⁶ The FDA approved the first CAR T-cell therapy in 2017 to treat children with acute lymphoblastic leukemia (ALL), demonstrating unprecedented curative potential in patients who had exhausted all other therapeutic options.³⁷

However, translating the miraculous efficacy of CAR-T from liquid tumors to solid tumors remains highly complex. Solid tumors are protected by a dense, fibrotic, and highly immunosuppressive tumor microenvironment that physically and chemically repels T-cell infiltration.³⁶ Furthermore, tumors frequently undergo "antigen loss," actively shedding the specific proteins the CAR-T cells are programmed to attack, rendering the therapy useless.³⁶ Additionally, the constant exposure to tumor antigens within this hostile microenvironment induces profound "T-cell exhaustion," an acquired state of severe cellular dysfunction where the engineered cells simply stop attacking.³⁶ Recent breakthroughs addressing these barriers include the development of Tumor-Infiltrating Lymphocyte (TIL) therapies. Recently approved for metastatic

melanoma, TIL therapy bypasses peripheral blood engineering and instead extracts highly activated, naturally occurring T-cells directly from the resected tumor matrix, enhancing the body's natural response before reinfusion.

Precision Medicine and Molecular Profiling

Precision oncology fundamentally abandons the generalized "one-size-fits-all" chemotherapy paradigm in favor of highly specific pharmacological interventions based entirely on the unique genetic signature of an individual's tumor.³⁸ Advances in diagnostics, such as liquid biopsies and the highly sensitive detection of circulating tumor DNA (ctDNA), allow oncologists to monitor tumor mutational dynamics in real-time through a simple blood draw, circumventing the risks and delays of repeated invasive tissue biopsies.

When specific genetic vulnerabilities are identified, highly targeted inhibitors are deployed. For instance, in patients with pancreatic or breast cancer harboring inherited BRCA1 or BRCA2 mutations, the tumors exhibit profound defects in homologous recombination DNA repair.³⁸ These specific tumors are highly susceptible to PARP inhibitors (e.g., Olaparib), which exploit a concept called synthetic lethality to completely collapse the cancer cell's remaining DNA repair mechanisms, inducing rapid apoptosis while sparing healthy cells.³⁸

Similarly, mutations in the KRAS gene—historically considered "undruggable" due to its smooth protein structure and implicated in roughly 25% of all human cancers, particularly deadly pancreatic, lung, and colorectal variants—are now actionable targets.³⁸ Novel FDA-approved targeted inhibitors like sotorasib and adagrasib bind specifically to mutant KRAS proteins, demonstrating substantial efficacy and progression-free survival in advanced settings.³⁸

Furthermore, the rapid adaptation of mRNA technology has catalyzed the development of personalized mRNA cancer vaccines. The period from 2024 to 2025 witnessed unprecedented clinical advances in this domain. Personalized mRNA vaccines are synthesized to train the immune system to recognize tumor-specific neoantigens unique to a patient's cancer. For example, clinical trials for the personalized mRNA-4157 vaccine showed a massive 44% reduction in recurrence risk for melanoma, achieving a 3-year sustained benefit. Pancreatic ductal adenocarcinoma (PDAC) trials demonstrated a 4-year immune persistence, significantly reducing recurrence in responders. Additionally,

layered nanoparticle mRNA vaccines for brain cancers (e.g., Glioblastoma) have achieved unprecedented 48-hour immune system activation in trials. Another groundbreaking 2025 discovery revealed that cancer patients who simply received standard mRNA COVID vaccines within 100 days of starting immunotherapy were twice as likely to be alive three years after treatment, sparking Phase III trials to potentially integrate these vaccines into the standard of care to broadly stimulate the immune system against tumors.

Surgical Oncology: Operational Procedures and Advanced Techniques

Surgical oncology remains the definitive, central pillar of solid tumor management. Over recent decades, the discipline has evolved dramatically from purely extirpative, highly morbid procedures to highly refined, individualized, and organ-sparing techniques, deeply integrated with systemic treatments. Modern surgical procedures are systematically classified based on their primary clinical objective.

Taxonomy of Oncological Surgery

Surgical interventions in oncology are broadly stratified into several distinct clinical categories, each requiring entirely different planning and execution strategies: Diagnostic and Staging Surgery forms the absolute foundation of cancer care. Procedures such as needle biopsies, incisional biopsies, or exploratory laparoscopies are utilized to obtain physical tissue samples, definitively confirm histopathology, identify the exact molecular type of cancer, and accurately determine the true anatomical extent of the disease throughout the body.³⁹

Curative, or Primary Surgery, is executed when a tumor is highly localized and deemed fully anatomically resectable.³⁹ The singular objective is the total macroscopic and microscopic excision of the malignant mass with negative margins, leaving the patient with absolutely no evidence of disease.³⁹

Preventative, or Prophylactic Surgery, involves the aggressive removal of entirely non-cancerous but highly susceptible tissues to prevent an almost certain malignant transformation later in life.³⁹ Common examples include prophylactic bilateral mastectomies and oophorectomies in patients harboring high-risk BRCA1/2 genetic mutations, or the routine excision of precancerous polyps during a colonoscopy.³⁹

Debulking, or Cytoreductive Surgery, is strictly employed when complete curative resection is anatomically impossible because the tumor has entangled critical neurovascular structures or organs.³⁹ Instead of attempting a cure, the surgeon removes as much of the tumor mass as safely possible to relieve local physiological pressure and to maximize the efficacy of subsequent adjuvant chemotherapy or radiation, which function better on smaller tumor volumes.³⁹

Palliative Surgery is critical for managing advanced, incurable disease, focusing strictly on severe symptom control and quality of life improvement rather than extending overall survival.³⁹ Examples include deploying internal stents to relieve agonizing bowel compressions, removing solitary metastatic masses pressing on the spinal cord, or performing surgical nerve blocks to permanently interrupt severe chronic pain signals.⁴² The field is advocating for a three-tier nomenclature—Curative, Palliative, and Potentially Life-Prolonging (PLP)—to better reflect the realities of modern extended survivorship and mitigate the defeatist psychological stigma often associated with the term "palliative".⁴³ Finally, Restorative or Reconstructive Surgery follows primary excisions to restore the physical appearance and mechanical function of the body, such as complex flap reconstructions following severe head and neck cancer excisions.³⁹

Standard and Groundbreaking Surgical Techniques

Standard oncological surgeries are frequently defined by the specific organ system involved and the radicality required. Breast cancer management typically involves a highly localized lumpectomy (breast-conserving surgery) or a full, radical mastectomy depending on tumor size and multifocality.⁴⁴ Thoracic oncology relies heavily on lobectomies (removal of a lung lobe) or pneumonectomies (removal of the entire lung) for aggressive lung cancer excision.⁴⁵ Gastrointestinal malignancies may necessitate massive, highly complex resections, ranging from total colectomies and proctectomies to highly intricate pancreaticoduodenectomies (the Whipple procedure), which involves the coordinated removal of the head of the pancreas, the duodenum, the gallbladder, and a portion of the bile duct.³⁹

Non-traditional ablative techniques have also gained massive clinical traction for patients who cannot tolerate massive open surgery. Cryosurgery utilizes localized extreme cold, delivered via liquid nitrogen or argon gas through a thin probe, to instantly freeze and destroy malignant tissues.⁴⁶ This is frequently used in localized prostate, liver, cervical, and skin cancers.⁴⁶ Conversely, laser surgery employs highly focused, high-powered light

beams to precisely vaporize and destroy cancer cells with minimal bleeding, often used for superficial lesions or to relieve occlusions in the trachea or esophagus.⁴⁶

Cytoreductive Surgery (CRS) and Hyperthermic Intraperitoneal Chemotherapy (HIPEC)

One of the most significant and grueling breakthroughs in the surgical management of highly advanced abdominal malignancies is the advent of Cytoreductive Surgery combined with Hyperthermic Intraperitoneal Chemotherapy (CRS-HIPEC).⁴⁷ Historically, cancers originating in the appendix, colon, or ovaries that subsequently metastasized to the peritoneal lining (peritoneal carcinomatosis) were considered universally terminal, and surgical intervention was strictly limited to palliation. The CRS-HIPEC protocol has completely revolutionized this paradigm.⁴⁷

The procedure is a massive, multi-step operation that frequently lasts over ten hours. First, the surgical team performs extensive cytoreductive surgery, meticulously dissecting and excising all visible tumor nodules from the delicate peritoneal cavity and stripping the lining of the abdomen.⁴⁸ The absolute goal is to achieve a Completeness of Cytoreduction (CC) score of CC0 (no visible disease remaining) or CC1 (remaining nodules are less than 2.5 mm in diameter).⁵⁰ Following maximum macroscopic clearance, the abdominal cavity is flooded with a highly concentrated chemotherapy solution (frequently Mitomycin C) that is heated to exactly 42 degrees Celsius (108 degrees Fahrenheit).⁵⁰ The abdomen is temporarily closed, and the hot chemotherapy is circulated mechanically throughout the cavity for approximately 90 minutes.⁴⁹

The hyperthermia provides immense dual benefits: it possesses direct cytotoxic effects on cancer cells, physically destroying them with heat, and simultaneously increases the permeability of cellular membranes, drastically enhancing the tissue penetration and deep efficacy of the chemotherapeutic agent to eradicate microscopic disease.⁴⁷ Because the chemotherapy is administered directly into the sealed peritoneal cavity rather than systemically via the bloodstream, clinicians can safely deliver substantially higher, highly lethal localized doses while drastically mitigating systemic toxicities like bone marrow suppression or severe organ failure.⁴⁷

Extensive data from the Mayo Clinic highlights the profound efficacy and the inherent, severe risks of this procedure. In a detailed retrospective study of selected patients, primarily dealing with mucinous adenocarcinoma of the appendix (56.3%) and colorectal

adenocarcinoma (31.3%), a CC0/CC1 score was achieved in an impressive 64.1% of surgeries performed with curative intent.⁵⁰

Mayo Clinic CRS-HIPEC Clinical Outcomes	Data Point / Percentage
Cure Rate	25% to 30%
Completeness of Cytoreduction (CC0/CC1)	64.1% (Curative intent group)
30-Day Postoperative Mortality	0%
Postoperative Complication Rate	60.4%
Severe Complications (Grade IV)	34.4%
Median Recurrence-Free Survival (Total Group)	19.3 months
Median Recurrence-Free Survival (Curative Intent)	23.0 months

50

Impressively, CRS-HIPEC offers a definitive cure for 25% to 30% of these previously

"incurable" patients.⁵⁰ At a median follow-up of 12 months, nearly 48% of patients showed no evidence of disease.⁵⁰ However, rigorous patient selection is absolutely paramount, as the procedure subjects the body to immense physiological stress. While 30-day mortality was remarkably reported at 0%, over 60% of patients experienced significant postoperative complications, with 34.4% of those categorized as severe (Grade IV), requiring prolonged intensive care.⁵⁰ The average length of hospital stay following the procedure is 8 to 10 days.⁴⁸ Despite the intense recovery profile, CRS-HIPEC has definitively emerged as the standard of care for specific peritoneal surface malignancies and metastatic appendiceal cancers in the United States.⁵⁰

Minimally Invasive and Robotic-Assisted Surgery (RAS)

To actively counteract the massive morbidity, severe pain, and prolonged recovery times of traditional open surgeries, surgical oncology has rapidly integrated Minimally Invasive Surgery (MIS) techniques, specifically video-assisted thoracoscopic surgery (VATS), laparoscopy, and highly advanced robotic-assisted surgery (RAS).⁵¹ Robotic systems, such as the industry-standard da Vinci Surgical System, act as an advanced interface, translating the surgeon's natural hand movements into highly precise, tremor-filtered, scaled-down actions of instruments operating inside the patient's body.⁵²

These platforms afford surgeons three-dimensional, high-definition visualization and unparalleled, multi-axis dexterity in tightly confined anatomical spaces, such as deep within the pelvis during a radical prostatectomy or deep within the thoracic cavity during lung resections.⁵¹ Recent 2024 and 2025 analyses offer strong comparative evidence: a systematic meta-analysis of over 3,100 randomized controlled trial patients found that RAS resulted in a significantly lower rate of conversion to open surgery and reduced hospital lengths of stay compared to conventional laparoscopy, alongside superior pathologic outcomes like lower circumferential resection margin (CRM) positivity. However, robotic procedures frequently feature longer operative times and higher financial costs—a 2024 study recorded a median hospitalization cost of 117,822 RMB for RACS versus 78,174 RMB for laparoscopy. Ultimately, patients often qualitatively report higher satisfaction; for example, patients undergoing robotic-assisted prostatectomy frequently demonstrate superior postoperative urinary and sexual function compared to open surgery, greatly enhancing long-term quality of life.

Critical Decision-Making and Clinical Analysis

The sheer, overwhelming complexity of modern oncology—where a single patient may require neoadjuvant chemotherapy, massive surgical resection, localized radiotherapy, adjuvant targeted therapies, and extensive psychological support—mandates highly structured, multi-layered frameworks for clinical decision-making. No single physician possesses the breadth of knowledge required to optimize care across all these modalities independently.

The Multidisciplinary Tumor Board (MTB)

The absolute global gold standard for cancer care coordination is the Multidisciplinary Tumor Board (MTB) or Multidisciplinary Team (MDT) meeting.⁵³ These formal forums bring together a diverse cohort of specialists—including surgical oncologists, medical oncologists, radiation therapists, expert pathologists, and specialized radiologists—to collectively synthesize complex diagnostic data and formulate consensus-driven, evidence-based treatment algorithms.⁵⁵ Decades of clinical practice demonstrate that MTBs definitively enhance accurate disease staging, increase patient enrollment in clinical trials, and ensure strict, verifiable adherence to contemporary treatment guidelines.⁵³

However, critical qualitative analysis of the actual MTB process reveals significant systemic flaws regarding true, authentic patient-centered care. While surveys indicate that 95% of MTB members strongly believe patient preferences and views should actively inform decision-making, the structural reality of these fast-paced meetings heavily favors a strictly biomedical model of disease.⁵³ In practice, MDT discussions are frequently dominated entirely by senior physicians, often explicitly excluding allied health professionals, nurses, or social workers who may have a much closer, nuanced understanding of the patient's subjective psychological state or home environment.⁵³

To compensate for the patient's physical absence from these meetings, clinicians inadvertently construct what sociologists term an "evidential patient"—a hollow proxy identity built entirely on curated laboratory data, radiological imaging, and highly selective clinical information, entirely stripping away the human element.⁵⁴ Furthermore, when physicians disagree on the optimal therapeutic path during the meeting, there is a heavily documented tendency to actively conceal this uncertainty from the patient later on. Clinicians selectively "frame" the final recommendation to project absolute medical

confidence, which fundamentally undermines the ethical process of true shared decision-making and patient autonomy.⁵⁴

Artificial Intelligence as a Decision Support System

To optimize MTBs, handle the crushing volume of new cancer cases, and reduce cognitive overload, the oncology field is rapidly adopting Artificial Intelligence (AI) and Machine Learning (ML) tools as highly advanced clinical decision support systems (DSS).⁵⁵ AI algorithms are increasingly utilized to instantly analyze complex imaging and pathology slides, rapidly identifying established biomarkers for diagnosis in lung, breast, brain, prostate, and colorectal cancers with accuracy rates that occasionally surpass human specialists.⁵⁷ Recently, in 2024, a study evaluated the use of Large Language Models (LLMs) like Claude3-Opus and GPT4-Turbo directly within breast cancer MDTs. These AI models generated appropriate adjuvant treatment suggestions with impressive accuracy rates of 86.6% and 85.7% respectively, closely matching reference expert decisions, particularly in identifying necessary endocrine and targeted therapies, though they still face limitations in perfectly validating surgical interventions.

In the realm of applied therapeutics, AI platforms like CURATE.AI are being deployed in real-time to continuously adjust the dosing of toxic chemotherapies like capecitabine based on continuous, dynamic biomarker analysis, maximizing drug efficacy while preventing severe toxicity.⁵⁶ Furthermore, ambient listening AI technologies are being integrated into surgical clinics to instantly transcribe and document clinical encounters, drastically reducing the administrative burden on surgical oncologists; however, studies indicate these systems currently generate an average of nearly 3 errors per clinical note, requiring rigorous manual oversight to prevent the injection of factious information into the patient's permanent record.⁵⁸ While AI offers immense promise, its rapid deployment raises severe ethical and logistical concerns regarding algorithmic "black boxes" (where the AI's reasoning cannot be explained), clinical liability when algorithms recommend fatal errors, data privacy risks with third-party servers, and the profound emotional impact of confronting high-probability mortality predictions generated by non-human systems.⁵⁷

Clinical Risk Assessment Tools

Before any major surgical or medical intervention is authorized by an MTB, a precise,

objective quantification of the patient's overall physiological resilience is required. Aggressive treatments demand immense physiological reserves. Several validated performance scales are utilized universally across the discipline to predict morbidity and mortality risks:

Clinical Assessment Scale	Primary Focus and Utility	Sample Grading / Scoring
ECOG Performance Status	Determines patient's ability to tolerate aggressive therapies, particularly chemotherapy; evaluates ambulatory status and self-care capability.	Grade 0: Fully active, no restriction. Grade 2: Ambulatory, capable of self-care, unable to carry out work activities. Grade 4: Completely disabled, cannot carry on self-care, bed-bound.
Karnofsky Performance Status (KPS)	Holistic assessment based on a patient's ability to carry out activities of daily living and the exact degree of medical assistance required.	100: Normal, no complaints, no disease evidence. 70: Cares for self, unable to do active work. 40: Disabled, requires special care and extensive assistance.
ASA Physical Status (ASA-PS)	Preoperative stratification tool used universally by anesthesiologists and surgeons to predict surgical morbidity, mortality, and	Risks of morbidity increase by 42.6% and mortality by 112% for each standard deviation increase in ASA

	length of stay.	grade.
--	-----------------	--------

59

These indispensable tools actively guide physicians in balancing the aggressive pursuit of a cure against the fundamental bioethical mandate of *nonmaleficence* (do no harm). For example, recommending a highly toxic, debilitating chemotherapy regimen or a massive, 10-hour cytoreductive surgery to a patient with an ECOG score of 4, a Karnofsky score of 30, or an ASA-PS of IV would almost certainly accelerate physiological collapse and death rather than prolong meaningful life.⁶⁰ These scales inject necessary objective reality into complex clinical decision-making.

Ethical Considerations in Oncological Research and Practice

As the technological and pharmacological frontiers of oncology rapidly expand, so too do the intricate, often unprecedented ethical dilemmas facing clinicians, researchers, and policymakers. These severe challenges span the core mechanics of clinical trials, the implementation of highly autonomous surgical robotics, and the macro-level justice of global healthcare equity.

Ethics in Clinical Trials: The Therapeutic Misconception and Placebos

Clinical research relies entirely on the ultimate trust and vulnerability of patient volunteers.⁶⁴ A persistent, deeply embedded ethical challenge in oncology trials is the phenomenon known as the "Therapeutic Misconception" (TM).⁶⁵ First identified in 1982 during psychiatric studies, TM occurs when a research participant fundamentally fails to distinguish between the core objective of a clinical trial (which is to generate generalizable, population-level scientific knowledge) and the objective of standard clinical care (which is to provide individualized, optimal therapeutic benefit to the specific patient).⁶⁶ Desperate, terminally ill cancer patients may erroneously believe that a Phase I dose-escalation trial—designed solely to test maximum human toxicity limits—is actually designed specifically to cure their disease.⁶⁵ This fundamental misunderstanding completely compromises the ethical validity of their informed consent, and some bioethicists argue that physician-investigators inadvertently encourage this misconception

by maintaining a therapeutic posture during research recruitment.⁶⁵

Furthermore, the utilization of placebos in oncology trials is particularly contentious and heavily scrutinized by Institutional Review Boards (IRBs).⁶⁷ Regulatory frameworks strictly dictate that placebo-controlled trials in cancer are only ethically justifiable if participants assigned to the placebo arm are not substantially more likely to die, suffer irreversible severe morbidity, or endure severe discomfort compared to those receiving standard active treatments.⁶⁸ If a known, effective, life-saving standard of care exists for a specific cancer, substituting it with a placebo is an egregious ethical violation and fundamentally contravenes the duty of care.⁶⁷ However, in advanced metastatic settings where all proven therapies have been exhausted, utilizing a placebo alongside optimal palliative pain care is scientifically necessary to isolate the true efficacy of an experimental drug and prevent the masking effects of observer bias, without exposing the patient to the unknown toxicities of unproven agents.⁶⁸

The Ethics of High-Tech Surgical Innovation

The massive integration of robotic-assisted surgery (RAS) and the dawn of telesurgery introduce novel ethical paradigms, particularly regarding the nuances of informed consent and conflicts of interest.⁶⁹ While patients are often eager to undergo "robotic" procedures due to aggressive institutional marketing campaigns and an implicit, often unfounded assumption of technological superiority, surgeons carry the strict ethical burden of fully disclosing the reality of the technology.⁷⁰ A comprehensive robotic informed consent must explicitly address patient misconceptions.⁷⁰ The surgeon is ethically obligated to disclose their own specific learning curve and exact experience level with the robotic platform, openly acknowledging that an inexperienced robotic surgeon may yield vastly inferior oncological outcomes compared to a highly experienced traditional open surgeon.⁷⁰ Furthermore, if the utilization of the multimillion-dollar robotic system is driven more by surgeon preference or institutional financial incentives to recoup purchasing costs rather than absolute clinical necessity, this distinction must be made entirely transparent to the patient prior to surgery.⁶⁹

Equity, Justice, and Global Access

Perhaps the most profound and devastating ethical crisis in modern oncology is the gross inequity in global access to life-saving research, early diagnostics, and advanced

therapies. The fundamental bioethical principle of *Justice* strictly demands the fair and equitable distribution of medical resources.⁶⁷ Yet, as the WHO global survey data explicitly highlights, only 39% of participating nations adequately finance core cancer management within their universal health benefit packages (HBP), and a mere 28% cover basic palliative care and general pain relief.²

This systemic disparity is glaringly evident in the availability of highly specialized, curative treatments. High-income countries are four to seven times more likely to cover advanced radiation services and up to twelve times more likely to offer life-saving stem-cell transplantations compared to lower-income nations.² To combat this, massive structural frameworks are being introduced, such as the 2025 proposed National Radiation Therapy Policy Blueprint in India, which aims to reduce rural-urban divides by incentivizing rural investments, using pooled brachytherapy, and employing AI for remote planning—serving as a model for low- and middle-income countries.

On a micro-level, even within wealthy nations, severe equity issues persist in clinical trial enrollment. Strict, exclusionary eligibility criteria—such as rigid age cutoffs, minor comorbidities, or a lack of premium insurance coverage—often result in the systemic exclusion of vulnerable, elderly, and minority populations.⁶⁴ This not only denies these populations access to cutting-edge therapies but yields clinical trial data that completely lacks generalizability to the broader, real-world demographic.⁷² If the oncological medical community is to truly uphold the principle of respecting patient autonomy and beneficence, it must first actively dismantle these systemic barriers to ensure that all patients, regardless of geographic location, race, or socioeconomic status, possess an equal opportunity to access standard-of-care treatments and innovative clinical trials.⁷²

Conclusion

The continuous pursuit of understanding, treating, and ultimately curing neoplastic diseases represents a monumental scientific and humanitarian endeavor. The trajectory of modern oncology is definitively shaped by an extraordinary convergence of highly specialized disciplines. Basic molecular research has successfully mapped the intricate architecture of tumor survival, identifying hyperactive biological pathways like PI3K/AKT and stealthy genetic mutations that drive uninhibited cellular proliferation. Simultaneously, applied clinical research has successfully weaponized these fundamental biological discoveries, birthing an unprecedented era of precision oncology where highly targeted therapies such as PARP inhibitors, mRNA vaccines, and CAR-T cells actively

reprogram the body's cellular machinery to hunt down malignancies with unparalleled specificity.

In the surgical domain, the field continues to rapidly evolve from the blunt force of historical extirpation to the refined, minimally invasive precision of advanced robotic assistance and the aggressive, curative ambition of highly complex procedures like CRS-HIPEC. These advanced surgical interventions, while physically taxing and complex, actively push the boundaries of survivorship, transforming formerly terminal prognoses into manageable, chronic conditions, and in properly selected cases, offering absolute, definitive cures.

Yet, the objective data strictly dictates that the success of oncology cannot be measured exclusively by cellular responses in laboratories or the technical success of high-tech operating theaters. The qualitative reality of the patient—encompassing the profound psychological weight of extended survivorship, the devastation of financial toxicity, and the complex manifestation of paraneoplastic syndromes—demands a medical ecosystem that is as deeply empathetic as it is scientifically rigorous. Furthermore, the true, global utility of this immense volume of research is fundamentally gated by health economics and bioethics. The future of oncology relies not merely on the invention of the next breakthrough molecular inhibitor or surgical robot, but on the absolute ethical imperative to democratize access to these life-saving technologies. To fulfill the ultimate promise of cancer research, the global medical community must ensure that the transition from empirical laboratory data to equitable, patient-centered clinical care is seamless, just, and universally accessible to all populations.

Author Contributions

Samuelson G conceived the study, conducted the literature review, synthesized interdisciplinary research findings, and prepared the manuscript for publication.

Funding Statement

No external funding was received for this research.

Conflict of Interest Statement

The author declares the following conflict of interest: My father was diagnosed with the disease.

Works cited

1. Tackling the Impact of Cancer on Health, the Economy and Society | OECD, https://www.oecd.org/en/publications/tackling-the-impact-of-cancer-on-health-the-economy-and-society_85e7c3ba-en.html
2. Global cancer burden growing, amidst mounting need for services, <https://www.who.int/news/item/01-02-2024-global-cancer-burden-growing--amidst-mounting-need-for-services>
3. Tumor heterogeneity: causes and consequences - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC2814927/>
4. The Development and Causes of Cancer - The Cell - NCBI Bookshelf, <https://www.ncbi.nlm.nih.gov/books/NBK9963/>
5. Review: the Contribution of both Nature and Nurture to Carcinogenesis and Progression in Solid Tumours - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC4842185/>
6. A Comprehensive Analysis of the PI3K/AKT Pathway: Unveiling Key ..., <https://pmc.ncbi.nlm.nih.gov/articles/PMC10462777/>
7. The Pathogenic Role of PI3K/AKT Pathway in Cancer Onset and Drug Resistance: An Updated Review - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8394096/>
8. The PI3K-AKT-mTOR Pathway and Prostate Cancer: At the Crossroads of AR, MAPK, and WNT Signaling - MDPI, <https://www.mdpi.com/1422-0067/21/12/4507>
9. The Role of PI3K/AKT and MAPK Signaling Pathways in Erythropoietin Signalization - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8307237/>
10. Paraneoplastic Syndromes: An Approach to Diagnosis and Treatment - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC2931619/>
11. Paraneoplastic Syndromes - StatPearls - NCBI Bookshelf - NIH, <https://www.ncbi.nlm.nih.gov/books/NBK507890/>
12. Updated Diagnostic Criteria for Paraneoplastic Neurologic Syndromes, <https://www.neurology.org/doi/10.1212/NXI.0000000000001014>
13. Methodological standards for qualitative and mixed methods patient centered outcomes research - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7756351/>
14. Understanding Patient-Centered Outcomes Research (PCOR): Principles, Methodologies, and Applications - Carenity, <https://pro.carenity.com/library/patient-centered-outcomes-research/>
15. Patient Centered Outcomes Research Committee - NRG Oncology, <https://www.nrgoncology.org/committees-programs/non-disease-site-scientific-committees/patient-centered-outcomes-research/>
16. Questionnaires - EORTC - Quality of Life, <https://qol.eortc.org/questionnaires/>
17. The effects of exercise on the quality of life of patients with breast cancer: a systematic review and meta-analysis based on the QLQ-C30 quality of life scale -

- Gland Surgery, <https://gs.amegroups.org/article/view/114103/html>
18. Concepts Assessed In The EORTC QLQ-C30 Questionnaire Are Important And Relevant For Patients With Cancer,
<https://www.eortc.org/blog/2022/11/25/concepts-assessed-in-the-eortc-qlq-c30-questionnaire-are-important-and-relevant-for-patients-with-cancer/>
 19. Identifying Changes in Scores on the EORTC-QLQ-C30 Representing a Change in Patients' Supportive Care Needs - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4405431/>
 20. Assessing Preoperative (EORTC) QLQ-C30 Score in Elderly Patients with Colorectal Cancer: Results from a Prospective Cohort Study - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11508690/>
 21. Getting back on track after treatment of cancer: A qualitative interview study of cancer survivors' experiences - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/39787181/>
 22. 'Because Everybody is so Different': a qualitative analysis of the lived experiences and information needs of rectal cancer survivors - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC8137244/>
 23. Breast cancer survivorship needs: a qualitative study - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10795302/>
 24. Patient Experiences After Aborted Cancer Surgery: A Qualitative Study - PubMed,
<https://pubmed.ncbi.nlm.nih.gov/37540329/>
 25. The Body after Cancer: A Qualitative Study on Breast Cancer Survivors' Body Representation - MDPI, <https://www.mdpi.com/1660-4601/19/19/12515>
 26. Understanding Cancer Survivors' Needs and Experiences Returning to Work Post-Treatment: A Longitudinal Qualitative Study - MDPI,
<https://www.mdpi.com/1718-7729/29/5/245>
 27. Cancer statistics, 2024 - PubMed, <https://pubmed.ncbi.nlm.nih.gov/38230766/>
 28. American Cancer Society Releases Latest Global Cancer Statistics; Cancer Cases Expected to Rise to 35 Million Worldwide by 2050,
<https://pressroom.cancer.org/GlobalCancerStatistics2024>
 29. Cancer Facts & Figures 2024,
<https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2024/2024-cancer-facts-and-figures-acf.pdf>
 30. Cancer Survival Statistics - SEER,
<https://seer.cancer.gov/statistics/types/survival.html>
 31. Cancer treatment and survivorship statistics, 2025 - PMC - NIH,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12223361/>
 32. Cancer Stat Facts: Cancer of Any Site - SEER,
<https://seer.cancer.gov/statfacts/html/all.html>
 33. The Incidence and 5-Years Relative Survival Rate of Cancer Among People Younger Than 25 Years in the US, 2000 to 2020: A SEER-Retrospective Study and

- Fertility Preservation Considerations - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12409066/>
34. Five-Year Survival Rates - SEER Training Modules - National Cancer Institute,
<https://training.seer.cancer.gov/prostate/intro/survival.html>
 35. Economic Burden - Cancer Atlas,
<https://canceratlas.cancer.org/taking-action/economic-burden/>
 36. Editorial: Expanding CAR-T cell therapy — breakthroughs from cancer to autoimmune diseases - Frontiers,
<https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2025.1649045/full>
 37. CAR T Cells: Engineering Immune Cells to Treat Cancer - NCI - National Cancer Institute, <https://www.cancer.gov/about-cancer/treatment/research/car-t-cells>
 38. 10 Breakthrough Research and Clinical Trials Shaping the Future of Oncology,
<https://honcology.com/blog/10-breakthrough-research-and-clinical-trials-shaping-the-future-of-oncology>
 39. Types of Surgery for Cancer Treatment | Stanford Health Care,
<https://stanfordhealthcare.org/medical-treatments/c/cancer-surgery/types.html>
 40. Types of Surgery for Cancer Treatment | The Surgical Clinic,
<https://thesurgicalclinics.com/types-of-surgery-for-cancer-treatment/>
 41. Cancer Surgery Types: Curative, Debulking and More,
<https://www.cancercenter.com/community/blog/2023/06/cancer-surgery-types>
 42. Palliative Surgery - SEER Training Modules,
<https://training.seer.cancer.gov/treatment/surgery/types/palliative.html>
 43. Treatment Classification by Intent in Oncology—The Need for Meaningful Definitions: Curative, Palliative and Potentially Life-Prolonging - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11433215/>
 44. What Are the Most Common Cancer Surgeries? Types of Surgery Explained - Liv Hospital,
<https://int.livhospital.com/most-common-cancer-surgeries-types-of-surgery/>
 45. 6 Most Common types of cancer surgeries - Dr. Lalit Banswal,
<https://www.drlalitbanswal.com/6-most-common-types-of-cancer-surgeries/>
 46. Specialized Types of Cancer Surgery | Johns Hopkins Medicine,
<https://www.hopkinsmedicine.org/health/treatment-tests-and-therapies/specialized-types-of-cancer-surgery>
 47. Cytoreductive Surgery and Hyperthermic Intraperitoneal Chemotherapy - Fred Hutch,
<https://www.fredhutch.org/en/patient-care/treatments/cytoreductive-surgery-and-hyperthermic-intraperitoneal-chemotherapy.html>
 48. Hyperthermic Intraperitoneal Chemotherapy (HIPEC) and Cytoreductive Surgery (CRS): Age-Related Outcomes and a Look into the Future - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11816354/>

49. Cytoreductive Surgery (CRS) With Hyperthermic Intraperitoneal Chemotherapy (HIPEC) | UH Seidman Cancer Center | University Hospitals,
<https://www.uhhospitals.org/services/cancer-services/surgical-oncology/cytoreductive-surgery-with-hyperthermic-intraperitoneal-chemotherapy>
50. Cytoreductive surgery and HIPEC offers effective treatment for ...,
<https://www.mayoclinic.org/medical-professionals/cancer/news/cytoreductive-surgery-and-hipec-offers-effective-treatment-for-selected-patients-with-peritoneal-carcinomatosis/mac-20429726>
51. Editorial: Innovations in surgical oncology - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC10446962/>
52. AMA Code of Medical Ethics' Opinions Related to Robotic Surgery,
<https://journalofethics.ama-assn.org/article/ama-code-medical-ethics-opinions-related-robotic-surgery/2023-08>
53. How do patients make decisions in the context of a multidisciplinary team: an ethnographic study of four head and neck cancer centres in the north of England - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC9413178/>
54. Multidisciplinary team decision-making in cancer and the absent patient: a qualitative study,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC4964245/>
55. The Role of Artificial Intelligence on Tumor Boards: Perspectives from Surgeons, Medical Oncologists and Radiation Oncologists - MDPI,
<https://www.mdpi.com/1718-7729/31/9/369>
56. Artificial Intelligence and Decision-Making in Oncology: A Review of ...,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12423120/>
57. 2025 Oncology Year in Review: Breakthrough Cancer Treatments and Research - Binaytara,
<https://binaytara.org/cancernews/article/2025-oncology-year-in-review-breakthrough-cancer-treatments-and-research>
58. AI Moves From Tool to Partner in the Modern Surgical Practice - General Surgery News,
<https://www.generalsurgerynews.com/In-the-News/Article/04-26/Surgery-AI-Partner-in-Modern-Clinical-Practice/80218>
59. ECOG Performance Status Scale,
<https://ecog-acrin.org/resources/ecog-performance-status/>
60. Karnofsky Performance Status Scale - MDCalc,
<https://www.mdcalc.com/calc/3168/karnofsky-performance-status-scale>
61. Evaluating the Real-World Predictive Utility of Karnofsky and ECOG Performance Status for 90-Day Survival After Oncologic Surgery for Metastatic Spinal Tumors - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC12651469/>
62. Letter to the Editor: ASA Physical Status Classification in Surgical Oncology and the Importance of Improving Inter-Rater Reliability - KoreaMed Synapse,
<https://synapse.koreamed.org/articles/1023393>

63. ASA-ECOG as a combined tool for peri-operative risk stratification in COVID-19 survivors – A step towards optimizing healthcare resource utilization - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8364169/>
64. Ethical guideposts to clinical trials in oncology - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC1891175/>
65. Review: the ethics of clinical trials - ecancer, <https://ecancer.org/en/journal/article/387-review-the-ethics-of-clinical-trials>
66. Clinical Trials and Medical Care: Defining the Therapeutic Misconception - Research journals - PLOS, <https://journals.plos.org/plosmedicine/article/file?type=printable&id=10.1371/journal.pmed.0040324>
67. Ethical Considerations for Oncology Clinical Trials - Biodexa Pharmaceuticals PLC, <https://biodexapharma.com/patient-resource/ethical-considerations-for-oncology-clinical-trials/>
68. Ethical, Scientific, and Regulatory Perspectives Regarding the Use of Placebos in Cancer Clinical Trials - ASCO Publications, <https://ascopubs.org/doi/10.1200/JCO.2007.13.5335>
69. A proposed ethical framework for informed consent in elective robotic telesurgery - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12995965/>
70. The ethics of robotic surgical systems is a conversation of informed consent - Wightman, <https://vats.amegroups.org/article/view/5469/html>
71. Ethical Considerations and Equipoise in Cancer Surgery - PMC - NIH, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11422545/>
72. The Ethical Imperative of Equity in Oncology: Lessons Learned From 2020 and a Path Forward - ASCO Publications, https://ascopubs.org/doi/10.1200/EDBK_100029