Neural Epilepsy: A Comprehensive Multi-Scalar Analysis of Pathogenesis, Diagnostics, and Advanced Therapeutics

Author: Samuelson G.

Independent Researcher

Abstract

Epilepsy remains one of the most neurobiologically complex and globally debilitating neurological diseases, characterized by an enduring, pathological predisposition to generate spontaneous, recurrent seizures. This comprehensive review synthesizes the multi-scalar pathophysiology of epileptogenesis—ranging from subcellular ion channelopathies, epigenetic reprogramming, and autophagic dysregulation, to the macroscopic topological reorganization of thalamocortical neural networks. We detail the latest global epidemiological trends and the modernized 2025 International League Against Epilepsy (ILAE) classification framework. Furthermore, this paper highlights transformative advancements in diagnostic neuroimaging, specifically the integration of High-Density Electroencephalography (hd-EEG), the Virtual Epileptic Patient (VEP) computational pipeline, and emerging Artificial Intelligence (AI) architectures. Finally, we explore the rapidly expanding therapeutic arsenal, detailing novel precision neuropharmacology (e.g., cenobamate, fenfluramine, XEN1101), bioelectronic neuromodulation, cutting-edge optogenetics, and the overarching shift toward a precision medicine framework modeled after advanced oncology.

Keywords: precision medicine in epilepsy, systems medicine, biomarkers, digital technologies, data science [52]

---

1. Introduction: The Ontological and Clinical Conceptualization of the Epileptic Brain

Epilepsy represents one of the most neurobiologically complex and pervasively

debilitating neurological diseases known to medicine, characterized fundamentally by an enduring, pathological predisposition to generate spontaneous, recurrent seizures.[1] Historically, the conceptualization of epilepsy has evolved from a purely symptom-oriented descriptive framework to a highly sophisticated neurobiological paradigm. The theoretical definition, officially established by the International League Against Epilepsy (ILAE) in 2005, articulated epilepsy as a primary brain disorder defined by the propensity for abnormal, excessive, and hypersynchronous neuronal discharges, alongside the myriad of subsequent cognitive, psychological, and social consequences.[2] However, the transition of this conceptual definition into a framework of clinical utility required the development of a highly rigorous operational architecture.

The subsequent 2014 practical definition reclassified epilepsy as a disease defined by the occurrence of at least two unprovoked (or reflex) seizures separated by an interval of more than 24 hours, or a single unprovoked seizure coupled with a high calculated recurrence probability of at least 60% over the next 10 years, or the definitive diagnosis of an established epilepsy syndrome.[3] This transition from a conceptual to an operational definition underscores a critical paradigm shift in contemporary neuroscience: the recognition of epileptogenesis as a dynamic, temporal continuum rather than a static, binary state.[2] This continuum acknowledges that the "disease" of epilepsy can be considered resolved in individuals who have outgrown age-dependent epilepsy syndromes or who have remained seizure-free for ten years while being unmedicated for at least five.[3]

The underlying neurobiological substrate of this disease involves a profound, progressive disruption in the ionic milieu of the central nervous system, fundamentally altering the intricate chemical and electrical gradients that govern neuronal resting membrane potentials and action potential generation.[4] Seizures manifest through highly diverse, heterogeneous mechanisms—ranging from monogenic channelopathies to widespread post-traumatic architectural remodeling—yet they consistently reflect a catastrophic failure of the delicate homeostatic balance between excitatory and inhibitory synaptic transmission.[4] Understanding this profound pathology requires a multi-scalar investigation, ranging from the sub-cellular alterations in ion channel kinetics and epigenetic transcriptional modifications to the macro-scale topological reorganization of thalamocortical neural networks.[5] The contemporary approach to epilepsy research integrates these deep biological insights with advanced computational modeling, high-density multimodal neuroimaging, and precision target-based pharmacology, paving

the way for targeted interventions that move definitively beyond mere symptom suppression toward disease modification.[8]

2. Global Epidemiology and Socioeconomic Disparities

The epidemiological landscape of epilepsy highlights a staggering global health burden characterized by significant socioeconomic, regional, and demographic disparities. Epilepsy affects more than 50 million individuals worldwide, solidifying its position as one of the most common chronic noncommunicable diseases of the central nervous system globally.[10] The distribution of this disease burden is highly asymmetric, with nearly 80% of affected individuals residing in low- and middle-income countries (LMICs).[10]

A detailed decomposition analysis of the Global Burden of Disease (GBD) data from 1990 to 2021 reveals highly nuanced longitudinal trends across prevalence, incidence, mortality, and disability-adjusted life years (DALYs).[12] While age-standardized mortality and DALY rates have demonstrated a measurable, statistically significant decline over the past three decades—reflecting gradual improvements in pharmacological accessibility, the proliferation of newer antiseizure medications, and broader healthcare infrastructure development—the absolute incidence and prevalence of epilepsy paradoxically continue to rise across all Socio-demographic Index (SDI) levels.[12] In 2021, epidemiological calculations identified approximately 24.22 million active patients with epilepsy globally, translating to an age-standardized prevalence rate of 307.38 per 100,000 persons and an age-standardized incidence rate of 42.82 per 100,000 persons.[14]

Epidemiological and Demographic Metric	High-Income Countries (HIC) / High SDI	Low- and Middle-Income Countries (LMIC) / Low SDI	Global Aggregates (2021 GBD Data)
Annual Incidence	Approximately 49	Ranging up to 139	~42.82 per 100,000 persons

Rate	per 100,000 persons	per 100,000 persons	(Age-standardized)
Prevalence Trends	Stable or Moderately Increasing	Sharply Increasing	~307.38 per 100,000 persons (Age-standardized)
Primary Etiological Risk Drivers	Genetic syndromes, neurodegenerative diseases, stroke	Endemic infections (malaria, neurocysticercosis), perinatal anoxia, TBI	Bimodal age distribution (distinct peaks in pediatric and geriatric populations)
Treatment Gap and Efficacy	Minimal (< 20% untreated)	Extremely High (Up to 75% remain untreated)	Up to 70% of global patients could achieve seizure freedom with proper care
Average Annual Percentage Change (AAPC) in Incidence (Adolescents/Young Adults)	Declining or stable depending on the specific region	Increasing with an AAPC of 0.32% in low SDI regions	Incidence increased globally from 35.88 to 40.48 per 100,000 among youth (1990-2021)

The stark contrast in absolute incidence rates—estimated at 49 per 100,000 persons in high-income nations versus an alarming 139 per 100,000 in certain LMICs—is heavily influenced by the disproportionate prevalence of endemic epileptogenic conditions.[10] Parasitic central nervous system infections such as neurocysticercosis and cerebral malaria, combined with a much higher frequency of road traffic injuries (traumatic brain

injury), birth-related anoxic trauma, and significant variations in the availability of preventive medical infrastructure, systematically elevate the seizure risk in resource-limited settings.[10] Furthermore, the risk of premature mortality among individuals with epilepsy is up to three times higher than that of the general population, a statistic driven by sudden unexpected death in epilepsy (SUDEP), status epilepticus, and seizure-related accidents.[10]

Advanced epidemiological projections utilizing the Bayesian Age-Period-Cohort (BAPC) model forecast a concerning trajectory: the global number of incident cases is predicted to continue climbing steadily, potentially reaching 1.47 million incident cases annually among adolescents and young adults alone by the year 2044.[13] This persistent epidemiological challenge highlights the profound inadequacy of current global healthcare infrastructures and underscores the urgent necessity for highly scalable, cost-effective neuro-interventions, particularly in sub-Saharan Africa and other low-SDI regions that are furthest from the optimal epidemiological frontier.[12]

3. Nosological Architecture and the 2025 ILAE Classification

Standardization in the nomenclature and classification of epileptic seizures is paramount for accurate clinical diagnosis, epidemiological tracking, and the execution of robust, reproducible pharmacological trials. Building upon the foundational 1981 framework and the subsequently refined 2017 operational classification, the International League Against Epilepsy (ILAE) published a highly modernized, updated classification system in 2025.[15] Forged through a rigorous, modified Delphi consensus process involving 37 international experts across all global ILAE regions, the 2025 classification dramatically streamlined the nosological structure, significantly reducing the recognized seizure types from a cumbersome 63 down to a highly practical, clinically actionable 21 types.[17]

The revised architectural framework meticulously preserves the four fundamental overarching seizure classes: Focal, Generalized, Unknown (whether focal or generalized), and Unclassified.[15] However, to enhance scientific clarity and clinical utility, the task force introduced several profound conceptual and structural shifts:

Classification Paradigm	2017 ILAE Framework	2025 ILAE Framework Update	Clinical and Neurobiological Rationale
Nomenclature of Origin	Included "Onset" (e.g., Focal Onset Seizure)	"Onset" explicitly removed (e.g., Focal Seizure)	Shifts focus to the broader network manifestation rather than a strictly localized, presumed origin point.[18]
Diagnostic Stratification	Blended primary and secondary clinical features	Strict distinction between Classifiers and Descriptors	Classifiers dictate biological syndromes and treatments; Descriptors purely detail specific phenomenological features without altering the primary diagnosis.[18]
State of Awareness	"Awareness" utilized as the primary differentiator	"Consciousness" replaces "Awareness" as the classifier	Consciousness is now rigidly and practically defined by the presence of both awareness (recall) and behavioral responsiveness.[15]
Motor Manifestations	Dichotomy of Motor vs.	Observable vs. Non-observable	Acknowledges that critical

	Non-motor	manifestations	semiological features (e.g., profound autonomic shifts, internal sensory auras) lack a distinct somatic motor component.[18]
Semiological Timeline	Classified primarily by the first initial sign	Described via the precise chronological sequence of signs	Provides a much higher-fidelity mapping of the seizure's spatial propagation across distinct cortical and subcortical networks.[15]
Specific Seizure Types	63 distinct types, omitting specific myoclonic variants	21 streamlined types; Epileptic negative myoclonus officially added	Enhances usability while accurately accommodating the specific motor arrest seen in certain developmental and epileptic encephalopathies.[17]

This highly optimized classification hierarchy is explicitly designed to be universally scalable, offering a basic, simplified version that can be reliably applied in resource-limited or primary care environments, alongside an expanded, exhaustive version that details complex temporal sequences and somatotopic data essential for advanced diagnostic and surgical decision-making in highly specialized tertiary epilepsy centers.[15]

4. Cellular and Molecular Neurobiology of Epileptogenesis

Epileptogenesis—the insidious, highly dynamic biological process by which a structurally or functionally normal brain develops the enduring predisposition to generate spontaneous seizures—is driven by a cascade of complex, mutually reinforcing molecular, transcriptional, and cellular aberrations.[5] At the absolute crux of this widespread pathology is an unyielding, pathological imbalance between excitatory glutamatergic signaling and inhibitory GABAergic transmission, a perturbation that compromises the network's ability to self-regulate.[5]

4.1 Ion Channelopathies and Receptor Dysfunction

Genetic epilepsies, including devastating pediatric encephalopathies, frequently emerge from highly specific ion channelopathies that destabilize the neuronal resting membrane potential or action potential kinetics. Voltage-gated sodium channels (NaV) are critical determinants of action potential initiation, upstroke, and propagation. Missense mutations in the *SCN1A* gene, which encodes the alpha subunit of the NaV1.1 channel, are famously responsible for a diverse spectrum of seizure disorders, ranging from relatively benign familial febrile seizures to Generalized Epilepsy with Febrile Seizures Plus (GEFS+) and the catastrophic Dravet syndrome.[5] Because the NaV1.1 channel is disproportionately expressed in fast-spiking, parvalbumin-positive GABAergic interneurons, loss-of-function mutations in this specific channel paradoxically lead to widespread neocortical and hippocampal hyperexcitability due to a profound failure of feed-forward and feedback inhibition.[5]

Conversely, the pathological overexpression or altered gating kinetics of certain channels actively drives focal excitability. For instance, recent neuro-oncological research has demonstrated that tumor necrosis factor-alpha (TNF-α) released from tumor-derived exosomes mediates the specific overexpression of NaV1.6 channels, directly contributing to neuronal hyperexcitability and spontaneous seizures in tumor-associated epilepsy models.[21] Beyond the scope of sodium channels, mutations or acquired dysfunctions in hyperpolarization-activated cyclic nucleotide-gated (HCN) channels, Kv7 potassium channels (often referred to as M-channels), and the Na-K-2Cl cotransporter isoform 1 (NKCC1) dramatically alter the neuronal refractory period and the intracellular chloride gradient, further dismantling the threshold for hypersynchronous firing.[19]

4.2 The Role of Glia, Neuroinflammation, and the Blood-Brain Barrier

The neurobiology of seizures is not solely neuronal; it is fundamentally tripartite, heavily involving complex astrocytic and microglial networks that govern the extracellular milieu.[20] Astrocytes are absolutely indispensable for maintaining the glutamate/GABA-glutamine cycle, which is essential for neurotransmitter homeostasis.[20] During intense periods of synaptic activity, astrocytes absorb excess synaptic glutamate and GABA, enzymatically converting them into glutamine, which is subsequently shuttled back to presynaptic neuronal terminals for neurotransmitter resynthesis and repackaging.[20] In the epileptogenic brain, the downregulation of astrocytic glutamine synthetase or the malfunction of excitatory amino acid transporters (EAATs) results in a catastrophic accumulation of extracellular glutamate, precipitating sustained neuronal depolarization and triggering NMDA receptor-mediated excitotoxicity.[20]

Neuroinflammation acts as both a potent catalyst and a destructive consequence of seizure activity. The purinergic P2X7 receptor, an ATP-gated ion channel, exhibits highly opposing, cell-specific effects during seizure propagation.[21] Its activation on microglia heavily drives the release of pro-inflammatory cytokines (such as interleukin-1 beta and TNF-alpha), exacerbating local network irritability and neurotoxicity, whereas its concurrent activation on certain neuronal populations may serve transient, albeit limited, neuroprotective functions.[21] This pervasive inflammatory milieu directly compromises the tight junctions of the blood-brain barrier (BBB), leading to severe pathophysiologic consequences.[21] Peri-ictal BBB dysfunction—frequently visualized using advanced MRI techniques—permits the extravasation of serum albumin into the brain parenchyma.[21] This albumin binds avidly to astrocytic TGF-beta receptors, initiating a transcriptional cascade that downregulates inward-rectifying potassium channels (Kir4.1) and aquaporins, thereby severely impairing the astrocyte's ability to buffer extracellular potassium, a condition that serves as a potent, direct trigger for local neuronal hypersynchrony.[21]

4.3 Epigenetic Modulation, Kinase Activity, and Mitochondrial Dysfunction

Epileptogenesis also triggers massive, enduring transcriptional reprogramming through diverse epigenetic mechanisms.[6] Epigenetic alterations, including widespread DNA hypermethylation, the activation of histone deacetylases (HDACs), and the differential, pathological expression of non-coding microRNAs, exert far-reaching effects on neuronal gene expression patterns.[6] These epigenetic modifications are tightly coupled to severe

mitochondrial dysfunction and profound oxidative stress.[6] The massive energy demands of sustained seizure activity rapidly deplete local ATP reserves, leading to the excessive generation of reactive oxygen species (ROS).[19] This oxidative stress damages delicate lipid cellular membranes, overwhelms and disrupts endogenous antioxidant buffering systems, and initiates pro-apoptotic cascades leading to neuronal death.[19]

Furthermore, post-translational modifications heavily regulate acute network dynamics. Protein kinases, including protein kinase A (PKA) and protein kinase C (PKC), actively phosphorylate various ion channels, fundamentally altering their conductance and gating properties.[6] This rapid kinase-mediated regulation is essential for physiological synaptic plasticity, but in the context of epileptogenesis, it permanently fine-tunes neuronal excitability toward a pathological set-point.[6] Targets such as the mammalian target of rapamycin (mTOR) pathway and P-glycoprotein (P-gp) further mediate aberrant cellular growth and multidrug resistance, underscoring the vast molecular complexity of the disease.[19]

4.4 Autophagy and Metabolic-Inflammatory Pathways

Recent neurobiological insights highlight the substantial role of autophagy and metabolic-inflammatory pathways in the maintenance of seizure thresholds.[5] When autophagy is impaired, the delicate balance between excitatory and inhibitory cerebral signals is disrupted. Specifically, reduced autophagic activity severely limits the translocation of vital GABA-A receptors to the neuronal plasma membrane, decreasing functional inhibitory input and predisposing the network to unprovoked seizures.[5] Concurrently, targeting these metabolic-inflammatory pathways has emerged as a novel therapeutic strategy; for example, utilizing PPARα agonists like fenofibrate has demonstrated potent, region-specific effects during epileptogenesis, actively reducing cortical inflammasome expression and differentially modulating glutamate receptor subunits across critical hippocampal subfields.[23]

5. Synaptic Plasticity, Circuit Reorganization, and Network Dynamics

The structural and anatomical corollary to molecular epileptogenesis is the profound spatial reorganization of both local microcircuits and global brain networks.[24] In temporal lobe epilepsy (TLE)—the most prevalent form of drug-resistant focal epilepsy in adults—this progressive reorganization is epitomized by a phenomenon known as mossy

fiber sprouting (MFS).[22]

5.1 Mossy Fiber Sprouting and Hilar Cell Loss

Following an initial precipitating injury (e.g., prolonged status epilepticus, severe concussive head trauma, or ischemic stroke), the brain experiences selective, rapid death of highly vulnerable neuronal populations.[26] This particularly affects the hilar polymorphic neurons and mossy cells within the dentate gyrus, as well as the pyramidal cells residing in the CA1 and CA3 regions—a constellation of damage clinically referred to as hippocampal sclerosis.[22] Under normal physiological conditions, the dentate gyrus functions as a high-resistance, restrictive "gate" that prevents excessive, high-frequency neocortical activity from invading the highly epileptogenic hippocampus.[26] However, the death of hilar mossy cells structurally disrupts the crucial excitatory drive onto inhibitory basket cells, resulting in a state of dangerous local disinhibition.[26]

In a misguided biological attempt at homeostatic compensation, the unmyelinated axons of the dentate granule cells (the mossy fibers) sprout and branch aberrantly, migrating out of their normal trajectory and penetrating into the inner molecular layer of the dentate gyrus.[22] While these sprouting fibers may initially attempt to re-innervate the surviving inhibitory interneurons to restore control, as terminal degeneration progresses, they ultimately form pathological, recurrent, monosynaptic excitatory connections directly onto the dendrites and somas of neighboring granule cells.[26] Rigorous computational models of the dentate gyrus—simulating 500 granule cells, 15 mossy cells, 6 basket cells, and 6 hilar perforant path-associated cells—demonstrate a highly significant, near-perfect correlation ($R = 0.95$, $p < 0.0001$) between the specific degree of MFS and explosive, exponential increases in granule cell spike generation.[29] The creation of these pathological recurrent excitatory loops functionally annihilates the dentate gate, allowing seizures to reliably originate, amplify, and reverberate within the mesial temporal structures.[26]

5.2 Long-Term Potentiation (LTP) and Depression (LTD)

The classic kindling model of epilepsy vividly illustrates how the fundamental physiological mechanisms of memory and learning—specifically long-term potentiation (LTP) and long-term depression (LTD)—are maliciously hijacked during the process of epileptogenesis.[30] Discovered in the early 1970s, LTP is characterized by a sustained increase in synaptic strength following brief, high-frequency stimulation.[32] In experimental kindling, repeated, sub-threshold electrical stimulation of limbic structures

induces a permanent, pathological enhancement of synaptic efficacy (a massive form of aberrant LTP), progressively lowering the seizure threshold day by day until spontaneous, unprovoked clinical seizures definitively emerge.[32] This runaway synaptic plasticity is physically associated with the dramatic spatial redistribution of crucial synaptic vesicle proteins, particularly synaptophysin, within the middle molecular layer of the dentate gyrus, fundamentally altering the structural connectome and neurotransmitter release probability of the entire entorhinal-hippocampal circuit.[31]

5.3 Macroscopic Network Synchronization and Seizure Propagation

At the macro-scale, clinical seizures are characterized not by isolated neuronal bursting, but by immense hypersynchronization across widely distributed functional brain networks.[4] Intracranial electrocorticogram (ECoG) recordings captured from patients with refractory epilepsy demonstrate that shifting network states inherently dictate the evolution of a seizure.[24] Seizures typically progress through distinct, mathematically definable dynamical states (S0, representing onset; S1, representing propagation; S2, representing termination).[25] Intriguingly, sophisticated connectivity algorithms reveal that seizure generation is primarily driven by the rapid, volatile reconfiguration of *weak* synaptic connections, which subsequently recruit, entrain, and structurally tighten *stronger* sub-networks as the seizure propagates outward from the onset zone.[25]

The critical transition from a highly localized interictal spike to a fully propagating clinical seizure depends heavily on the "inhibitory in-degree" (the density of inhibitory connections a neuron receives) of the surrounding tissue.[35] Detailed computational neural mass models reveal two distinct possible outcomes when paroxysmal activity attempts to invade a neighboring, healthy brain region:

1. **The Non-Propagative Scenario**: The recipient network successfully "controls" the incoming perturbation. In this state, fast-spiking (FS) inhibitory neurons are robustly recruited and maintain a high-frequency firing plateau (e.g., oscillating at ~15 Hz). This powerful inhibitory tone effectively suppresses the excitatory regular-spiking (RS) neurons, keeping their firing rates extremely low (near ~2 Hz), thereby extinguishing the seizure wave before it can spread.[35]
2. **The Propagative Scenario**: A critical loss of the excitatory/inhibitory balance occurs. The sheer magnitude of the incoming excitation overrides the local inhibitory capacity, causing the recipient network's excitatory population to suddenly discharge at massive rates. This failure allows the network to amplify the seizure and violently

pass it to the next efferent node.[35]

Beyond hypersynchronization, the biochemical mechanisms governing actual seizure termination involve a complex orchestration of ionic and neuromodulatory shifts.[23] Extracellular pH acidification coupled with massive adenosine release collectively act to suppress excitatory synaptic transmission.[23] Acid-sensing ion channels (such as ASIC1a) actively regulate neuronal excitability during these shifts; experimental disruption of ASIC1a exacerbates seizure severity, whereas its overexpression expedites termination.[23] Finally, intense Na+/K+-pump activity aimed at restoring the depleted ionic balance, alongside depolarization blocks caused by severe extracellular potassium accumulation, physically exhaust the network, driving the brain into the profound postictal suppression state.[23]

6. Advanced Diagnostics, High-Density Electrophysiology, and Multimodal Neuroimaging

The historical, largely subjective reliance on the visual inspection of low-resolution, standard scalp electroencephalography (EEG) is rapidly being superseded by an era of high-resolution neuroimaging, advanced signal processing, machine learning classification, and highly individualized computational simulations.[36] The overarching clinical goal of these technologies is the precise, millimeter-accurate delineation of the Seizure Onset Zone (SOZ) and the broader Epileptogenic Zone Network (EZN) to optimize surgical resection outcomes while sparing eloquent cortex.[39]

6.1 High-Density EEG and Electrical Source Imaging (ESI)

High-density EEG (hd-EEG), utilizing tightly spaced arrays of up to 256 electrodes, offers unparalleled temporal and spatial resolution for non-invasive clinical electrophysiology.[41] When combined with advanced, patient-specific structural MRI data, the hd-EEG signals are subjected to Electrical Source Imaging (ESI).[41] Advanced subspace imaging algorithms, such as First-order Inverse Numerical Evaluation (FINE), can localize current density dipoles deep within the 3D brain volume with remarkable precision, effectively mapping the hidden generators of surface potentials.[23] Furthermore, revolutionary sparse imaging algorithms like the Imaging of Rehabilitated EEG Sources (IRES) utilize convex optimization theory to estimate not only the anatomical epicenter of the epileptogenic source but also its objective spatial extent and volume.[23] Extensive

clinical data indicates that calculating ESI precisely at the 50% mark of the rising slope of an interictal spike provides the most accurate spatial localization of the primary epileptogenic generator, heavily influencing successful surgical resection margins, whereas ESI calculated at the spike peak often reflects secondary propagation pathways.[41]

6.2 Multimodal Imaging Integration

The sophisticated integration of hd-EEG with functional MRI (fMRI) and 18F-fluorodeoxyglucose positron emission tomography (18F-FDG-PET) allows researchers to correlate millisecond-level electrical discharges with localized hemodynamic responses (the BOLD signal) and baseline cerebral metabolism.[45] In this highly synchronized setup, interictal epileptiform discharges identified via hd-EEG serve as precise temporal triggers for fMRI General Linear Model (GLM) or Independent Component Analysis (ICA) processing, visually revealing the specific, widespread functional networks that are pathologically perturbed by the isolated epileptic spikes.[45]

6.3 The Virtual Brain and the Virtual Epileptic Patient (VEP)

Perhaps the most revolutionary advancement in the entire field of presurgical diagnostics is the development of the Virtual Epileptic Patient (VEP) pipeline, powered by The Virtual Brain (TVB) simulation technology and hosted on the European EBRAINS digital research infrastructure.[8] The VEP framework constructs a highly personalized, generative mathematical model of an individual patient's brain, designed to simulate personalized seizure dynamics and allow clinicians to conduct predictive "virtual surgeries" entirely *in silico* prior to performing actual tissue resection.[8]

The methodology underpinning the VEP is exhaustively rigorous and mathematically dense [40]:

1. **Structural Scaffolding**: T1-weighted MRI and diffusion-weighted MRI (DW-MRI) tractography are utilized to construct a patient-specific anatomical connectome.
2. **Dynamical Simulation**: The brain is digitally parcellated into distinct regions. Each node is populated with neural mass models driven by the "Epileptor"—a specialized system of nonlinear differential equations explicitly designed to mimic the transition between interictal, pre-ictal, and full ictal states.[40]
3. **Bayesian Inference**: The patient's actual, empirical seizure recordings, obtained via highly invasive stereoelectroencephalography (SEEG), are mapped directly to the structural model via a customized source-to-sensor gain matrix.[40]

4. **Target Identification**: Once optimized, the VEP workflow generates a statistical probability map defining the full Epileptogenic Zone Network (EZN).[40]

Currently, this transformative technology is undergoing rigorous prospective validation in the EPINOV (Improving Epilepsy surgery management and progNOsis using Virtual brain technology) clinical trial, which aims to boost surgical success rates globally far beyond the current, stagnant ~60% threshold.[40] The evaluation phase of the EPINOV trials is slated to conclude by late 2026, after which data analysis will ascertain whether these virtual brain twins can rapidly scale as the new standard of care globally.

7. Artificial Intelligence, Graph Neural Networks, and Transformer Architectures

As of 2025–2026, the integration of Advanced Learning Architectures (ALAs) has revolutionized EEG-based seizure detection and prediction. Traditional deep learning models, while effective at feature extraction, are rapidly being augmented or replaced by foundation model transformers, large language models, and explainable AI (XAI) pipelines designed to handle complex, noisy physiological data streams.

Graph Neural Networks (GNNs) have emerged as particularly powerful diagnostic tools due to their unique ability to model the high-dimensional, non-linear connectivity of the brain by representing EEG and SEEG channels as mathematical graphs. Spatiotemporal GNNs have achieved up to 98% precision and sensitivity in early seizure prediction by precisely capturing the dynamic synchronization and dependencies between distinct brain regions. Furthermore, advanced GNN frameworks are successfully automating the localization of the Seizure Onset Zone (SOZ) by integrating raw interictal SEEG signals directly with spatial MRI coordinates.

In parallel, Transformer architectures—specifically Vision Transformers (ViT) and Spatial-Temporal Hypergraph Attention Transformers—leverage sophisticated self-attention mechanisms to model global dependencies over long temporal sequences. Recent innovations such as Temporal-Patchify Encoding transform raw, 1D EEG signals into image-like patches suitable for segmentation networks like SegFormer or MambaVision. This technique drastically reduces computational latency by over 20% while maintaining exceptional accuracy. Moreover, novel multiclass artificial neural network (ANN) architectures can now simultaneously discriminate between various

neurological conditions (e.g., epilepsy versus schizophrenia) using frequency subbands, achieving 95% sensitivity and paving the way for scalable decision-support tools.

8. Precision Neuropharmacology: Expanding the Antiseizure Arsenal

While approximately 30% of global epilepsy patients remain persistently medically refractory, the pharmacological pipeline for novel antiseizure medications (ASMs) continues to produce highly targeted agents that move decisively beyond the broad-spectrum, non-specific CNS depression of legacy drugs like phenobarbital and valproate.[51] Recent FDA approvals and highly promising late-stage clinical trials highlight a definitive shift toward the precision modulation of specific channel subtypes, extrasynaptic receptors, and entirely novel biochemical pathways.[53]

8.1 Cenobamate: Dual-Action Excitability Dampening

Cenobamate represents a major, paradigm-shifting breakthrough for the treatment of refractory focal-onset seizures in adults.[56] Its remarkable clinical efficacy relies on a highly synergistic, dual mechanism of action that addresses both sides of the excitability equation.[56] Firstly, cenobamate acts as a potent positive allosteric modulator at the GABA-A receptor.[57] Secondly, it acts as a highly selective inhibitor of voltage-gated sodium channels. Crucially, patch-clamp electrophysiology demonstrates that it preferentially inhibits the non-inactivating persistent sodium current (INaP), primarily via direct action at NaV1.6 channels, while safely sparing the transient sodium current (INaT).[58]

8.2 Fenfluramine: Serotonergic and Sigma-1 Receptor Modulation

Originally utilized decades ago as a systemic anorectic agent, fenfluramine has been brilliantly repurposed and formally approved for the management of devastating developmental and epileptic encephalopathies, specifically Dravet syndrome and Lennox-Gastaut syndrome.[60] Fenfluramine stimulates the massive release of extracellular serotonin, acting as a potent agonist at 5HT1A, 5HT2A, and 5HT2C receptors.[60] However, exhaustive *in vitro* pharmacological profiling revealed that fenfluramine also acts as a highly potent positive allosteric modulator of the enigmatic Sigma-1 receptor.[60] The combined, concurrent activation of 5HT receptors and Sigma-1 receptors dramatically alters intracellular calcium dynamics, effectively uncoupling NMDARs from positive control and profoundly reducing the probability of excitotoxic,

calcium-mediated convulsive events.[62]

8.3 XEN1101 (Azetukalner): Targeted Potassium Channel Opening

XEN1101 (azetukalner) is an experimental, highly selective small-molecule opener of the KCNQ2/3 (Kv7.2/7.3) potassium channels.[64] By chemically potentiating the open state of these specific channels, XEN1101 drives a massive, sustained efflux of positively charged potassium ions out of the cell, resulting in profound membrane hyperpolarization.[64] Importantly, while XEN1101 shares a fundamental mechanism of action with the older drug retigabine, it possesses a distinctly engineered molecular structure that completely avoids the formation of chromophoric phenazinium-type dimers, eliminating severe retinal toxicity.[66] As of 2026, XEN1101 is undergoing extensive late-stage evaluation, including the X-TOLE2 and X-TOLE3 Phase 3 clinical trials testing its efficacy as an adjunctive therapy for focal-onset seizures, and the X-ACKT Phase 3 trial assessing its application in primary generalized tonic-clonic seizures (PGTCS).

8.4 Ganaxolone: Neurosteroids and Tonic Inhibition

Ganaxolone is a sophisticated synthetic analog of the naturally occurring endogenous neurosteroid allopregnanolone.[67] Ganaxolone exhibits profound efficacy and affinity for the unique delta-subunit-containing GABA-A receptors located peripherally in the perisynaptic and extrasynaptic spaces.[69] These specific extrasynaptic receptors generate a continuous "tonic" chloride current that provides powerful shunting inhibition.[69] This highly specialized mechanism is uniquely relevant for the treatment of catamenial epilepsy and rare pediatric genetic epilepsies, such as CDKL5 deficiency disorder.[67]

9. Neuromodulation and Bioelectronic Medicine

For the substantial cohort of patients who fall definitively under the umbrella of drug-resistant epilepsy (DRE)—and who are not viable candidates for traditional resective surgery—bioelectronic neuromodulation provides an increasingly sophisticated, hardware-driven alternative.[52] The FDA has formally approved three primary modalities, each offering distinct anatomical targets and operational paradigms: Vagus Nerve Stimulation (VNS), Deep Brain Stimulation (DBS), and Responsive Neurostimulation (RNS).[73]

RNS represents the absolute vanguard of closed-loop, personalized bioelectronic therapy.[72] RNS devices continually monitor local field potentials (LFPs) directly from the cortex via surgically implanted depth or subdural strip electrodes. Utilizing onboard, adaptive machine learning algorithms, the device learns to detect patient-specific pre-ictal electrographic signatures.[72] Upon detection, it delivers highly focal, microsecond electrical bursts designed to disrupt the hypersynchrony and abort the seizure before it can reach clinical manifestation.[72] Cutting-edge neurosurgical research is now actively exploring the placement of RNS depth electrodes within the anterior thalamus to treat patients whose seizures originate independently in both temporal lobes (bitemporal epilepsy), effectively hybridizing the deep anatomical targets of DBS with the highly efficient, battery-saving responsive timing of RNS.[75]

10. Optogenetics, Gene Therapy, and Viral Vector Targeting

The absolute frontiers of modern epilepsy research are currently defined by the radical transition from macro-scale, non-specific electrical modulation to precise, cell-type-specific genetic and optical engineering.[72]

10.1 Optogenetic Interrogation and Seizure Curtailment

Optogenetics permits the unprecedented spatial and temporal manipulation of specific neural circuits *in vivo* via the viral-mediated expression of light-sensitive ion channels and ionic pumps.[20] One of the most recent breakthroughs involves the development of a highly sensitive, moderately K+-selective channelrhodopsin known as HcKCR1-hs. Transcranial activation of HcKCR1-hs has been shown to significantly prolong the time to the first seizure and decrease total seizure activity in status epilepticus models, allowing for completely non-invasive, transcranial optogenetic inhibition of neural hyperactivity.

10.2 Adeno-Associated Virus (AAV) and Precision Gene Therapies

Gene therapy presents the ultimate, highly sought-after disease-modifying strategy for monogenic developmental and epileptic encephalopathies.[9] Adeno-associated viruses (AAVs) are the universally preferred delivery vector due to their remarkable neuronal tropism, low systemic immunogenicity, and immense capacity for capsid engineering to target specific cell layers.[78] However, AAVs possess a strict biological packaging capacity limit of approximately 4.7 kilobases, posing a severe, nearly insurmountable limitation

for treating conditions like Dravet syndrome, where the massive, mutated *SCN1A* gene is far too large to be inserted into a standard viral vector.[80] To circumvent this, researchers have engineered compact gene regulation therapies, such as ETX101, which delivers a custom-built "engineered transcription factor" directly into GABAergic inhibitory interneurons to radically upregulate the endogenous expression of NaV1.1 channels.[80]

11. Future Trajectories: Precision Medicine and the 2030 Horizon

As the global medical paradigm shifts aggressively toward the year 2030, the clinical management of epilepsy is poised to evolve completely from a reactive, symptom-suppression model to a proactive, biology-based, preventative ecosystem.[9] The trajectory of contemporary epileptology is beginning to mirror the precision oncology framework. By integrating Next-Generation Sequencing (NGS) and establishing actionable molecular biomarkers, physicians are moving beyond generalized syndromic treatments toward individualized therapeutic selection based entirely on the patient's underlying genomic architecture.

A major, pervasive factor contributing to drug-resistant epilepsy is the clinical failure to achieve adequate, therapeutic concentrations of ASMs precisely at the specific epileptogenic focus, heavily impeded by the highly restrictive, protective properties of the blood-brain barrier (BBB).[83] Emerging biomedical technologies are actively pioneering the use of targeted intracranial drug delivery devices and highly functionalized, biodegradable nanoparticles to bypass the BBB entirely.[84] Concurrently, non-invasive acoustic techniques utilizing MR-guided focused ultrasound are being optimized to temporarily and highly selectively increase BBB permeability in specific, millimeter-scale brain regions, facilitating the targeted entry of both large-molecule ASMs and massive viral vectors for localized gene therapy without the need for craniotomies.[52]

12. Synthesis and Concluding Perspectives

Neural epilepsy remains an incredibly intricate, multi-dimensional phenomenon, representing a spectacular, highly coordinated systemic failure of the central nervous system's endogenous homeostatic mechanisms. The exhaustive, multi-scalar data surrounding its underlying etiology—from highly specific point mutations in

voltage-gated ion channels and pathological epigenetic reprogramming at the molecular level, to the macroscopic architectural distortion witnessed in mossy fiber sprouting and broad thalamocortical hypersynchrony—illustrates definitively that epileptogenesis is a continuous, relentlessly evolving network disease rather than a static condition.

The trajectory of modern epileptology reveals an undeniable, accelerating pivot away from empirical, trial-and-error clinical pharmacology toward rationally designed, precision-guided medicine. The rapid maturation of computational neurology, spearheaded by revolutionary platforms such as The Virtual Epileptic Patient, provides clinicians with an unprecedented capability to mathematically invert empirical EEG data into causal structural topologies, fundamentally redefining pre-surgical evaluation and network mapping. Simultaneously, the advent of highly selective pharmacological modulators—capable of discerning between specific ion channel states or leveraging entirely novel receptors like Sigma-1—alongside the unprecedented potential of localized gene therapies like engineered transcription factors, promises to directly correct the underlying molecular deficits.

Ultimately, realizing the full, transformative promise of these scientific advancements requires a sustained, highly interdisciplinary global commitment to overcoming the massive translational barriers of targeted drug delivery, while simultaneously scaling these high-fidelity interventions to address the mounting, disproportionate epidemiological burden that continues to heavily afflict populations within the developing world. Through the absolute integration of molecular biology, high-dimensional neuroimaging, bioelectronics, and computational network theory, the historic goal of true disease modification transitions rapidly from theoretical possibility to imminent clinical reality.

13. Acknowledgements and Funding

This research review highlights advancements enabled by extensive international clinical collaborations. The EPINOV clinical trial (Improving Epilepsy surgery management and progNOsis using Virtual brain technology) discussed in this review is supported by a public grant from the French National Research Agency (ANR) under the "Investissements d'Avenir" program (reference: ANR-17-RHUS-0004).

Works cited

1. A practical clinical definition of epilepsy, https://www.ilae.org/files/ilaeGuideline/Definition2014.pdf
2. The 2014 Definition of Epilepsy: A perspective for patients and caregivers, https://www.ilae.org/guidelines/definition-and-classification/the-2014-definition-of-epilepsy-a-perspective-for-patients-and-caregivers
3. ILAE official report: a practical clinical definition of epilepsy - PubMed, https://pubmed.ncbi.nlm.nih.gov/24730690/
4. The Neurobiology of Epilepsy - PMC - NIH, https://pmc.ncbi.nlm.nih.gov/articles/PMC2492886/
5. Advances in understanding the pathogenesis of epilepsy ..., https://pmc.ncbi.nlm.nih.gov/articles/PMC10867297/
6. Unraveling the nexus of age, epilepsy, and mitochondria: exploring the dynamics of cellular energy and excitability - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC11413492/
7. Brain network dynamics codify heterogeneity in seizure propagation - bioRxiv, https://www.biorxiv.org/content/10.1101/2021.06.12.448205.full
8. LEARN: The Virtual Epileptic Patient: Part 1, https://www.thevirtualbrain.org/tvb/zwei/newswire-educase/single/42212-learn-the-virtual-epileptic-patient-part-1
9. Precision Epilepsy: How Gene-Targeted Therapies Are Changing Seizure Treatment in 2026 - GlobalRPH, https://globalrph.com/2026/03/precision-epilepsy-how-gene-targeted-therapies-are-changing-seizure-treatment-in-2026/
10. Epilepsy - World Health Organization (WHO), https://www.who.int/news-room/fact-sheets/detail/epilepsy
11. Epilepsy - World Health Organization (WHO), https://www.who.int/health-topics/epilepsy
12. Global, regional, and national burden of epilepsy, 1990–2021: a Global Burden of Disease study - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC12964325/
13. Global, regional and national burdens of epilepsy in the adolescents and young adults from 1990 to 2021 and its predictions - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC12487120/
14. A population-based analysis of the global burden of epilepsy across all age groups (1990–2021) - Frontiers, https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2024.1448596/full
15. A practical guide to the updated seizure classification 2025 - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC12747708/
16. Updated classification of epileptic seizures: Position paper of the International

League Against Epilepsy, https://www.ilae.org/files/dmfile/updated-classification-of-epileptic-seizures-2025.pdf
17. Updated classification of epileptic seizures (2025) // International League Against Epilepsy, https://www.ilae.org/updated-classification-epileptic-seizures-2025
18. The International League Against Epilepsy Updates Classification of Epileptic Seizures, https://www.ilae.org/news-and-media/news-about-ilae/the-international-league-against-epilepsy-updates-classification-of-epileptic-seizures
19. Biomolecular mechanisms of epileptic seizures and epilepsy: a review - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC11960269/
20. Neural circuit mechanisms of epilepsy: Maintenance of homeostasis at the cellular, synaptic, and neurotransmitter levels - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC12220727/
21. State of the art: glioma-associated epilepsy—bridging tumor biology and epileptogenesis - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC12514801/
22. Mossy Fiber Sprouting in Temporal Lobe Epilepsy: The Impact of Netrin-1, DCC, and Gene Expression Changes - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC11672915/
23. Electrophysiological Source Imaging - Biomedical Functional ..., https://www.cmu.edu/bme/helab/Research/ESI/index.html
24. Phase Synchronization Dynamics of Neural Network during Seizures - PMC - NIH, https://pmc.ncbi.nlm.nih.gov/articles/PMC6205102/
25. Dynamic Network Drivers of Seizure Generation, Propagation and Termination in Human Neocortical Epilepsy - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC4682976/
26. The Role of Synaptic Reorganization in Mesial Temporal Lobe Epilepsy - PMC - NIH, https://pmc.ncbi.nlm.nih.gov/articles/PMC2829602/
27. Is Mossy Fiber Sprouting a Potential Therapeutic Target for Epilepsy? - Frontiers, https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2018.01023/full
28. Role of Mossy Fiber Sprouting and Mossy Cell Loss in Hyperexcitability: A Network Model of the Dentate Gyrus Incorporating Cell Types and Axonal Topography, https://journals.physiology.org/doi/full/10.1152/jn.00777.2004
29. Mechanisms Underlying Hyperexcitability: Combining Mossy Fiber Sprouting and Mossy Cell Loss in Neural Network Model of the Dentate Gyrus - MDPI, https://www.mdpi.com/2227-9059/13/6/1416
30. Impairments of Long-Term Synaptic Plasticity in the Hippocampus of Young Rats during the Latent Phase of the Lithium-Pilocarpine Model of Temporal Lobe Epilepsy - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC8705146/
31. Long Term Potentiation (LTP) and Long Term Depression (LTD) Cause Differential

Spatial Redistribution of the Synaptic Vesicle Protein Synaptophysin in the Middle Molecular Layer of the Dentate Gyrus in Rat Hippocampus | Opera Medica et Physiologica, https://operamedphys.org/OMP_2016_03_0039
32. Interplay of epilepsy and long-term potentiation: implications for memory - PMC - NIH, https://pmc.ncbi.nlm.nih.gov/articles/PMC11760605/
33. Loss of Long-Term Potentiation at Hippocampal Output Synapses in Experimental Temporal Lobe Epilepsy - Frontiers, https://www.frontiersin.org/journals/molecular-neuroscience/articles/10.3389/fnmol.2020.00143/full
34. Network dynamics of the brain and influence of the epileptic seizure onset zone | PNAS, https://www.pnas.org/doi/10.1073/pnas.1401752111
35. A Model for the Propagation of Seizure Activity in Normal Brain ..., https://www.eneuro.org/content/9/6/eneuro.0234-21.2022
36. Advances in Neuroimaging and Deep Learning for Emotion Detection: A Systematic Review of Cognitive Neuroscience and Algorithmic Innovations - MDPI, https://www.mdpi.com/2075-4418/15/4/456
37. Artificial intelligence in electroencephalography analysis for epilepsy diagnosis and management - Frontiers, https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2025.1615120/full
38. LEARN: Modelling Epilepsy with The Virtual Brain, https://www.thevirtualbrain.org/tvb/zwei/newswire-educase/single/42218-learn-modelling-epilepsy-with-the-virtual-brain
39. Combining interictal intracranial EEG and fMRI to compute a dynamic resting-state index for surgical outcome validation - Frontiers, https://www.frontiersin.org/journals/network-physiology/articles/10.3389/fnetp.2024.1491967/full
40. Personalised brain modeling technique may lead to breakthroughs ..., https://www.humanbrainproject.eu/en/follow-hbp/news/2023/01/26/personalised-brain-modeling-technique-may-lead-breakthroughs-clinical-epilepsy-trial/
41. Visual analysis of high density EEG: As good as electrical source imaging? - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC6939057/
42. Feasibility of high‐density electric source imaging in the presurgical workflow: Effect of number of spikes and automated spike detection - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC10472417/
43. Persyst ESI powered by Epilog, https://www.persyst.com/technology/electrical-source-imaging/
44. High density EEG in current clinical practice and opportunities for the future - PMC - NIH, https://pmc.ncbi.nlm.nih.gov/articles/PMC8083969/
45. Multimodal integration of fMRI and EEG data for high spatial and temporal resolution analysis of brain networks - PMC,

https://pmc.ncbi.nlm.nih.gov/articles/PMC5682027/
46. A multimodal imaging approach to the evaluation of post-traumatic epilepsy - PMC - NIH, https://pmc.ncbi.nlm.nih.gov/articles/PMC3458199/
47. Neural network integration from multimodal imaging in epilepsy and healthy controls - eDiss, https://ediss.uni-goettingen.de/bitstream/handle/11858/15070/Daniel_vandeVelden_full_Thesis_v3.pdf?sequence=1&isAllowed=y
48. An Integrated Multi-Channel Deep Neural Network for Mesial Temporal Lobe Epilepsy Identification Using Multi-Modal Medical Data - MDPI, https://www.mdpi.com/2306-5354/10/10/1234
49. Multimodal Image Integration for Epilepsy Presurgical Evaluation: A Clinical Workflow, https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2021.709400/full
50. Virtual epileptic patient brain modeling: Relationships with seizure onset and surgical outcome - PubMed, https://pubmed.ncbi.nlm.nih.gov/35604575/
51. Current state of the epilepsy drug and device pipeline - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC11018510/
52. Precision medicine for epilepsy: challenges and perspectives for an optimized clinical care pathway - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC12460136/
53. Special Report: 2024 Neurology Drug & Device Approvals -, https://practicalneurology.com/diseases-diagnoses/alzheimer-disease-dementias/special-report-2024-neurology-drug-device-approvals/33149/
54. Update on Antiseizure Medications 2025 | Continuum, https://continuum.aan.com/doi/abs/10.1212/cont.0000000000001521
55. Update on Antiseizure Medications 2025 | Continuum, https://continuum.aan.com/doi/10.1212/cont.0000000000001521
56. Cenobamate, a Sodium Channel Inhibitor and Positive Allosteric Modulator of GABAA Ion Channels, for Partial Onset Seizures in Adults: A Comprehensive Review and Clinical Implications - PubMed, https://pubmed.ncbi.nlm.nih.gov/34207493/
57. Cenobamate modulates EEG cortical activity and connectivity in individuals with drug-resistant epilepsy: a pharmaco-EEG study - Frontiers, https://www.frontiersin.org/journals/neurology/articles/10.3389/fneur.2024.1502668/full
58. Cenobamate, a Sodium Channel Inhibitor and Positive Allosteric Modulator of GABAA Ion Channels, for Partial Onset Seizures in Adults: A Comprehensive Review and Clinical Implications - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC8293325/
59. clarification of the mechanism of action of cenobamate - American Epilepsy Society,

https://aesnet.org/abstractslisting/clarification-of-the-mechanism-of-action-of-cenobamate
60. Fenfluramine: A Review of Pharmacology, Clinical Efficacy, and Safety in Epilepsy - MDPI, https://www.mdpi.com/2227-9067/9/8/1159
61. Dual-Activity-of-Fenfluramine-(Fintepla®)-as-a-Serotonin-Receptor-Agonist-and-Positive-Sigma-1-Receptor-Modulator--Implication-for-Disease-Modification-in-Developmental-and-Epileptic-Encephalopathies - American Epilepsy Society, https://aesnet.org/abstractslisting/dual-activity-of-fenfluramine-(fintepla%C2%AE)-as-a-serotonin-receptor-agonist-and-positive-sigma-1-receptor-modulator--implication-for-disease-modification-in-developmental-and-epileptic-encephalopathies
62. Fenfluramine diminishes NMDA receptor-mediated seizures via its mixed activity at serotonin 5HT2A and type 1 sigma receptors | Oncotarget, https://www.oncotarget.com/article/25169/text/
63. Fenfluramine acts as a positive modulator of sigma-1 receptors, https://dravetfoundation.org/fenfluramine-acts-as-a-positive-modulator-of-sigma-1-receptors/
64. TMS as a pharmacodynamic indicator of cortical activity of a novel anti‐epileptic drug, XEN1101 - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC6856596/
65. XEN1101, a Novel Potassium Channel Modulator: Interim Data From an Ongoing, Long-Term, Open-Label Extension of a Phase 2b study (X-TOLE) in Adults with Focal Epilepsy (S44.008) - Neurology.org, https://www.neurology.org/doi/10.1212/WNL.0000000000203823
66. Encukalner - Wikipedia, https://en.wikipedia.org/wiki/Encukalner
67. Neurosteroid regulation of GABAA receptors: Focus on the α4 and δ subunits - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC2657726/
68. Neurosteroids and GABAA Receptor Interactions: A Focus on Stress - Frontiers, https://www.frontiersin.org/journals/neuroscience/articles/10.3389/fnins.2011.00131/full
69. Neurosteroid interactions with synaptic and extrasynaptic GABAa receptors: regulation of subunit plasticity, phasic and tonic inhibition, and neuronal network excitability - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC3832254/
70. Neurosteroid Modulation of Synaptic and Extrasynaptic GABA A Receptors of the Mouse Nucleus Accumbens - MDPI, https://www.mdpi.com/2218-273X/14/4/460
71. Neurosteroids and GABAergic signaling in health and disease - PMC - NIH, https://pmc.ncbi.nlm.nih.gov/articles/PMC5469411/
72. A Comprehensive Review of Emerging Trends and Innovative ..., https://pmc.ncbi.nlm.nih.gov/articles/PMC10527076/
73. The comparison of DBS and RNS for adult drug-resistant epilepsy: a systematic review and meta-analysis - Frontiers, https://www.frontiersin.org/journals/human-neuroscience/articles/10.3389/fnhum.2024.1429223/full

74. A Comprehensive Review of Emerging Trends and Innovative Therapies in Epilepsy Management - Semantic Scholar, https://www.semanticscholar.org/paper/A-Comprehensive-Review-of-Emerging-Trends-and-in-Ghosh-Sinha/0aed4d8c6225bd773a29788fa80141a95f7b6e01
75. Expanding Neurostimulation and Gene Therapy Possibilities | Johns Hopkins Medicine, https://www.hopkinsmedicine.org/news/articles/2025/12/expanding-neurostimulation-and-gene-therapy-possibilities
76. Gene Therapy for Epilepsy - Jasper's Basic Mechanisms of the Epilepsies - NCBI Bookshelf, https://www.ncbi.nlm.nih.gov/books/NBK609899/
77. Gene therapy for epilepsy: An emerging, promising approach for a serious neurological disorder - PubMed, https://pubmed.ncbi.nlm.nih.gov/41366856/
78. Gene therapy for epilepsy: An emerging, promising approach for a serious neurological disorder - PMC, https://pmc.ncbi.nlm.nih.gov/articles/PMC12869011/
79. Adeno-Associated Viral Vectors in the Treatment of Epilepsy - MDPI, https://www.mdpi.com/1422-0067/25/22/12081
80. Gene Therapy for Dravet Syndrome - Dravet Syndrome Foundation, https://dravetfoundation.org/dsf-funded-research/gene-therapy-for-dravet-syndrome/
81. Ian Miller, MD: Adapting AAV Gene Therapy for SCN1A-Positive Epilepsy | NeurologyLive, https://www.neurologylive.com/view/ian-miller-md-adapting-aav-gene-therapy-for-scn1apositive-epilepsy--
82. The Future of Epilepsy Care: 5 Predictions for 2026 and Beyond | Lampsy Health Blog, https://lampsyhealth.com/blog/5-predictions-for-epilepsy-care-2026-and-beyond
83. Intracranial Drug Delivery Devices: A Promising Approach for Future Epilepsy Treatment?, https://read.qxmd.com/read/41863263/intracranial-drug-delivery-devices-a-promising-approach-for-future-epilepsy-treatment
84. Future directions for epilepsy research - PubMed - NIH, https://pubmed.ncbi.nlm.nih.gov/11706087/