

Physics-Guided Multimodal Strategies for WHO Grade IV Tumors: Integrating Precision Energy-Based Therapies, Tumor Biology, Advanced Imaging, and Human-Body Safety Constraints

Author: SAMUELSON G

Independent Researcher

Email: gsamuelsonguna@gmail.com

ORCID: 0009-0005-8744-8178

Structured Abstract

Background: WHO Grade IV neoplasms (e.g. glioblastoma multiforme) are uniformly lethal under standard care. Median survival remains on the order of 12–18 months, and 5-year survival is ~5%. Current guidelines recommend maximal safe resection followed by chemoradiation (e.g. 60 Gy radiotherapy + temozolomide). Physics-based modalities (advanced radiotherapy, particle beams, focused ultrasound, electromagnetic fields, nanotechnology) offer new avenues for tumor ablation, but their integration remains experimental.

Objective: To propose an extremely aggressive, multi-modality treatment protocol (“cure at any cost”) for Grade IV tumors, combining physics-based therapies, microenvironment modulation, and novel interventions.

Methods: We review recent literature on advanced radiation techniques (protons, carbon ions, FLASH), thermal and non-thermal ablation (high-intensity focused ultrasound (HIFU), nanophotonic photothermal therapy, magnetic hyperthermia), alternating electric fields (TTFields), and tumor microenvironment-targeting strategies. We outline a combined protocol (maximal surgery + dose-escalated conformal RT with nanoparticles, simultaneous hyperthermia and TTFields, immune modulation) and design preclinical/clinical studies. Ethical/legal issues (e.g. excessive dose, experimental agents) are flagged.

Expected Results: We hypothesise dramatically improved local control and overall survival, at the expense of increased toxicity.

Conclusions: Although radical, the proposed integrated strategy is informed by recent primary studies. Safety concerns, ethical constraints, and practical limitations are acknowledged.

Keywords

Grade IV tumors, glioblastoma, radiotherapy, proton/carbon ion therapy, FLASH radiotherapy, focused ultrasound, hyperthermia, tumor-treating fields, nanotechnology, tumor microenvironment.

Executive Summary

- Grade IV malignancies (e.g. glioblastoma) are rapidly fatal under standard care. Recent data (NICE guidelines) report median survival ~15 months. Five-year survival remains <10%.
- Standard care (surgery + 60 Gy photon RT + concurrent and adjuvant temozolomide) is limited by radioresistance and tumor microenvironment (hypoxia, diffusion barriers).
- We propose an extreme combinatorial protocol: maximal cytoreductive surgery followed by *dose-escalated conformal RT* (including protons/heavy ions, and FLASH dosing) with *concurrent sensitisation* using **nanoparticle radioenhancers** (e.g. hafnium oxide), **thermal ablation** (focused ultrasound and magnetic hyperthermia), **electrical field therapy** (200 kHz TTFs), and **photothermal nanomedicine**. Treatment is guided by advanced physiological modelling of the tumor microenvironment.
- **Tumor microenvironment analysis:** We will characterise tumor hypoxia, acidity, vasculature, immune infiltrates and molecular markers (IDH, MGMT, etc.) to personalise therapy. For brain tumors, strategies to bypass the blood–brain barrier (e.g. MR-guided FUS-induced opening) are included. Normal tissue dose constraints are monitored by advanced imaging and real-time thermometry.
- **Preclinical studies:** Mouse models (orthotopic xenografts and GEMMs) will test safety/efficacy of the combined regimen. Phase I human trials will then escalate dose and modality combinations, focusing on toxicity endpoints.
- **Safety/Risk:** Major risks include acute toxicity (radiation necrosis, CNS injury, organ failure), long-term sequelae (neurocognitive impairment, secondary cancers), and off-target effects of nanoparticles or viruses. We will include stringent monitoring (e.g. MR-thermography, blood biomarkers) and planned dose de-

escalation rules. Ethical concerns (e.g. extreme interventions, uncertain benefit) are explicitly flagged.

- **Implementation:** A multidisciplinary team (neurosurgery, radiation physics, interventional radiology, oncology) will synchronise therapies (see flowchart). A tumor board will review each case.
- **Outcomes and Future Work:** We expect improved local control and survival, potentially converting lethal stage IV disease into chronic illness. Limitations include untested toxicity of combinations, regulatory hurdles, and high cost. Future work will refine the protocol via adaptive trial design and might incorporate AI-driven planning.

1. Introduction

Grade IV tumors (e.g. glioblastoma multiforme) are among the most lethal human cancers. These neoplasms exhibit highly aberrant (undifferentiated) cells that grow rapidly and invasively. Standard therapy (maximal surgical resection followed by radiotherapy with concurrent and adjuvant temozolomide) extends median survival to only ~15–20 months. Five-year survival is below 10%. Thus, novel approaches are urgently needed. Physics-based modalities—leveraging ionising radiation, sound waves, electromagnetic fields, and nanotechnology—offer new mechanisms to attack tumors. Recent advances (proton and carbon-ion therapy, ultrahigh-dose-rate (“FLASH”) radiotherapy, focused ultrasound, tumor-treating fields) show promise. However, individual modalities have plateaued, and no single method cures all patients. Here we propose an **extreme multi-modal protocol** aimed at curing Grade IV tumors “at any cost,” combining aggressive physics-based ablation with systemic modulation. We assume no fixed legal or ethical constraints, but explicitly discuss their implications.

1.1 Clinical Background and Standard of Care

WHO Grade IV gliomas (glioblastomas) are defined by necrosis and rapid mitotic activity. Patients typically present with neurological deficits and MRI evidence of heterogeneously enhancing masses. Prognosis is dismal: one recent report notes median overall survival ~15 months despite trimodal therapy. As NICE guidelines state: “Offer radiotherapy 60 Gy in 30 fractions with concomitant temozolomide, followed by up to 6 cycles of adjuvant temozolomide” for eligible newly diagnosed Grade 4 glioma. Even with this aggressive regimen, recurrence is nearly universal. Factors contributing to treatment failure include radioresistant tumor cells, hypoxic microregions, and infiltrative spread. Chemotherapy (alkylators, targeted agents) has limited penetration across the blood-brain barrier. In short, Grade IV tumors remain incurable by current standards.

1.2 Tumor Microenvironment and Physiological Barriers

The **tumor microenvironment (TME)** critically shapes therapy response. It comprises abnormal vasculature, cancer-associated fibroblasts (CAFs), immune cells, and

extracellular matrix. Tumor vessels are tortuous and leaky, causing hypoxia, acidosis and high interstitial pressure. These conditions impair drug delivery and render cells more radioresistant. Ionising radiation itself can induce vascular damage and fibrosis, exacerbating hypoxia. Moreover, irradiated TMEs often become immunosuppressive ($\uparrow$ TGF- β , regulatory T cells). Successful eradication thus requires not only killing tumor cells but also overcoming these physiological barriers.

In brain tumors, the blood–brain barrier (BBB) adds an additional challenge. Many drugs cannot penetrate unless actively transported. Failing to address the BBB results in subtherapeutic drug levels in infiltrating tumor cells. Focused ultrasound is one method to transiently open the BBB for enhanced delivery. Brain physiology also limits radiation dose: normal brain parenchyma is sensitive to radiation necrosis and cognitive injury. Thus, techniques that spare healthy brain or precisely target only tumor tissue are desirable.

1.3 Survey of Physics-Based Modalities

Radiotherapy (Conventional and Advanced): Photon RT (X-rays) is the mainstay. Conformal IMRT/VMAT can sculpt dose around critical structures, but exit dose and penumbra are unavoidable. Proton and heavy-ion beams offer better conformality (Bragg peak) and higher relative biological effectiveness (especially carbon ions). For example, recent NRG Oncology data suggest that proton dose-escalation to 75 Gy improved 2-year survival compared to standard photon RT. The absolute 2-year survival improvement was 6.8% (49.9% vs 43.1%). This signals possible benefit of particle therapy in GBM.

FLASH Radiotherapy: Ultra-high dose-rate (≥ 40 Gy/s) “FLASH” RT spares normal tissue compared to conventional RT, while retaining tumor control. Preclinical studies show significantly reduced neurocognitive decline in animals receiving electron-, photon-, or proton-FLASH. FLASH is poised for clinical translation and may be incorporated to deliver ablative doses while protecting healthy brain.

Tumor Treating Fields (TTFields): Alternating electric fields (~ 200 kHz) applied via scalp electrodes disrupt mitosis in glioma cells. In the EF-14 trial, adding TTFields to maintenance temozolomide reduced risk of death by $\sim 37\%$ and extended median survival by 5 months. These low-intensity fields (200 kHz) kill dividing glioma cells by interfering with microtubule spindle formation. TTFields are well tolerated; skin reactions are mild and compliance is high (patients wear the device >18 h/day). We will include TTFields as an adjuvant to chemoradiation in our protocol (noting that optimal tumor coverage is limited to ~ 7 cm depth from skull).

Focused Ultrasound (FUS): HIFU devices (often MR-guided) can thermally ablate deep tumors noninvasively. MRgFUS focuses ultrasound waves to raise target temperature $>60^\circ\text{C}$, causing coagulative necrosis within the focal zone. Because the beam is focused, surrounding tissue is largely spared. MDPI review notes that MRgFUS “induces deep coagulation necrosis within an elliptical focal area, effectively encompassing the entire tumor site”. Clinically, MRgFUS has been used to ablate metastatic bone lesions and

uterine fibroids; brain applications include thalamotomy and early trials in tumor ablation. Importantly, FUS can also open the BBB when used at low intensity with microbubbles. The Focused Ultrasound Foundation highlights its advantages: non-invasiveness, MR imaging guidance, repeated reversible BBB opening, and immune modulation. We incorporate FUS both as an ablation tool (for accessible portions of tumor) and as a drug delivery enhancer.

Nanophotonics and Photothermal Therapy: Nanomaterials (e.g. gold nanoshells, carbon nanotubes, or carbon dots) can convert near-infrared light to heat. Photothermal therapy uses injected nanoparticles that accumulate in tumors (via EPR effect) and then irradiates with NIR laser to locally heat and kill tumor cells. This approach has shown efficacy in preclinical tumor models. For example, gold nanorods raise tumor temperature selectively, inducing cell death without ionising radiation. Although deep penetration is limited, interstitial or fiber-optic delivery can reach internal tumors. We propose trials of nanoparticle-mediated photothermal ablation, especially in combination with other modalities. (No definitive clinical references yet in brain cancer, but early studies exist in prostate and breast).

Thermal Ablation – Magnetic Hyperthermia: An emerging modality uses magnetic nanoparticles and alternating magnetic fields (AMF). Iron-oxide nanoparticles (IONPs) are delivered IV or intratumorally; when exposed to AMF, they heat to ~45–50 °C (limited by coating) and thermally ablate cancer cells. The first US installation of a clinical AMF hyperthermia system (New Phase Ltd.) occurred in 2025. In that setup, IONPs accumulate in tumor vasculature and an induction coil heats them like an “induction cooktop”. This method is being trialled for metastatic solid tumors (excluding brain so far). Its advantages include the ability to treat deep, widespread lesions simultaneously. We include magnetic hyperthermia for extracranial Grade IV metastases and investigate adapting it for brain (via convection-enhanced infusion of IONPs).

Cryoablation: Extreme cooling (liquid nitrogen/argon probes) can also ablate tumor tissue. Cryotherapy produces an “ice ball” around the probe, causing direct cell kill and vascular stasis. This is well-established for liver, kidney and bone lesions. For brain tumors, stereotactic cryoprobes have been trialled, although precision is limited and risks include edema. Cryo will be reserved as a salvage approach for surgically accessible metastases, and is considered an “extreme” adjunct here.

Immunotoxins and Oncolytic Agents (Extremes): In cases of disseminated disease, we consider “biological extreme measures.” Examples might include intratumoural injection of potent immunotoxins (e.g. IL-13–pseudomonas exotoxin fusion), or engineered oncolytic viruses (HSV, adenovirus) to lyse tumor cells. While not physics-based, they represent “at any cost” interventions. Such agents often are lethal if misdirected, so we will flag them as ethically and legally challenging. (Not approved outside trials, but we discuss their theoretical integration).

Tumor Treating Fields (TTFields) and EM Fields: See above. We also consider low-intensity pulsed electric/magnetic fields beyond TTFields. For instance, experimental data suggest weak magnetic fields may affect cellular signalling. However, TTFields have the strongest evidence.

Integrated Physiology/TME Analysis: Our protocol requires detailed patient-specific modelling. Pre-treatment imaging (MRI, PET, perfusion scans) and biopsies will map oxygenation, interstitial pressure and molecular markers. We will use radiobiological models (e.g. TCP/NTCP) adjusted for FLASH effects and incorporate TME data (hypoxia fraction, pH) to predict treatment efficacy. For example, severely hypoxic regions may trigger combined hyperthermia + RT to improve oxygenation. Immune profiling (TILs, PD-L1 expression) will guide addition of immune checkpoint inhibitors as needed.

Comparative Modalities (see Table 1): We compare key therapies in terms of mechanism, benefits and risks. For instance, carbon ions (high LET) may overcome radioresistance, but availability and cost are limiting. FLASH RT spares normal brain (cognitive function) but requires special hardware. TTFields improve survival by ~5 months but require continuous patient compliance. FUS gives real-time imaging and can open BBB, yet skull heating limits power. A comparative table is provided below.

Table 1: Comparison of Advanced Therapeutic Modalities for Grade IV Tumors.

Modality	Mechanism	Advantages	Disadvantages / Key Risks	Example References
Photon Radiotherapy	DNA double-strand breaks; conventional fractionation	Widely available; proven efficacy with chemo	Normal tissue toxicity, limited by dose tolerance (cognitive decline)	Standard-of-care
Proton Therapy	Bragg-peak energy deposition	Reduced exit dose spares normal tissue; allows dose escalation	Limited availability; higher cost	NRG-BN001 phase II
Carbon/Ion Therapy	High-LET particle kill	Increased RBE, better kill of hypoxic/resistant cells	Very limited access; uncertain CNS tolerance	(theoretical, see reviews)
FLASH Radiotherapy	Ultra-high dose-rate photons/particles	Markedly spares normal tissue (e.g. cognitive function) while	Experimental; equipment not widely available	Preclinical studies

Modality	Mechanism	Advantages	Disadvantages / Key Risks	Example References
		maintaining tumor kill		
Tumor-Treating Fields (TTFields)	200 kHz alternating electric fields disrupt mitosis	Improves median survival by ~5 months in GBM; non-invasive once device worn	Requires continuous wear (≥ 18 h/day); local skin irritation	EF-14 trial
MR-guided FUS (HIFU/LIFU)	Focused ultrasound heating (or low-intensity with microbubbles)	Noninvasive focal ablation; MR-imaging guidance; BBB opening; immune stimulation	Skull attenuation; thermal spread risk; edema; incomplete penetration	Review; Trials underway
Nanophotonic PTT	Nanoparticles (e.g. Au) + NIR laser	Highly localized heat; minimal radiation dose	Light penetration limited; heating non-uniform; experimental	(preclinical studies)
Magnetic Hyperthermia	IV iron-oxide NPs + AMF (100–500 kHz)	Deep tumor heating; can treat multiple lesions simultaneously	Need NP delivery; potential systemic heating; first-in-human stage	Mayo Phase I (2025)
Cryoablation	Local freezing (-196°C)	Precise for accessible lesions; clear demarcation (“ice ball”)	Pain, swelling; freeze thaw damage to surrounding tissue	Clinical use (liver, prostate)
Histotripsy (non-thermal)	Mechanical cavitation disruption	Antigen-preserving ablation; strong immune activation	Experimental; requires special ultrasound pulses	Emerging research

Modality	Mechanism	Advantages	Disadvantages / Key Risks	Example References
Oncolytic Viruses	Engineered viruses infecting/killing tumor	Potential systemic anti-tumor effect; addresses micrometastases	Safety concerns (infection of normal cells); immune clearance	Clinical trials (e.g. T-VEC)
Immune Modulation	Checkpoint inhibitors, vaccines	May sustain long-term control; synergistic with ablation	Toxicity (autoimmunity); often ineffective in “cold” GBM	Trials ongoing

Table 1: Summary of modalities: mechanisms, potential benefits, and main risks. Examples of clinical evidence or review references are indicated.

2. Methods – Integrated Treatment Protocol

We propose a **multimodal curative protocol** structured as follows:

1. **Diagnosis and Profiling:** Standard workup (MRI/CT, biopsy). Perform molecular profiling (IDH1/2, MGMT methylation) and advanced imaging (PET for hypoxia, perfusion MRI). Generate a patient-specific physiological model (tumor geometry, vascular map, dose constraints).
2. **Maximal Cyto-reduction:** Surgical resection of all safely resectable tumor bulk. Use intraoperative MRI/fluorescence for maximal clearance. (Ethical note: in “no constraint” scenario, consider extended resection even at functional cost, but must discuss patient consent.)
3. **Concomitant Radiotherapy and Sensitisation:**
 - **Photon/Proton/Carbon RT:** Deliver a biologically intensified dose. Standard is 60 Gy, but we plan dose-escalation (e.g. 75 Gy) using protons to spare normal tissue. Optionally use carbon ions for their higher RBE in hypoxic regions. Where available, apply **FLASH** RT for portions of dose, interleaving with normal fractionation. (For example, 50 Gy via conventional RT + 25 Gy via FLASH).
 - **Nanoparticle Radioenhancers:** Inject hafnium-oxide nanoparticles (NBTXR3) into or around the tumor (by stereotactic delivery). Preclinical and Phase I studies show these NPs absorb X-ray/proton energy and release

low-range electrons, amplifying local dose. This allows higher effective dose to tumor without raising photon dose.

- **Concurrent Chemotherapy:** Continue temozolomide per Stupp protocol. Consider adding angiogenesis inhibitors (bevacizumab) or radiosensitisers (PARP inhibitors) off-label as “extreme measures,” noting limited evidence.

4. **Physical Ablations:** Interleaved with RT, perform focused ablation sessions:

- **MRgFUS Sessions:** Using MRI guidance, target residual tumor regions unreachable by surgery or too radioresistant. For each session, apply high-intensity focused ultrasound to create thermal necrosis within a precise volume. Use real-time MR-thermometry to monitor. Typical sonications raise target to $>60^{\circ}\text{C}$ in seconds. Multiple overlapping spots are used to cover irregular tumor shape.
- **Sonodynamic Therapy:** Administer a sonosensitising agent (e.g. 5-ALA) systemically, then apply low-intensity FUS with microbubbles to activate it. This can induce localized cytotoxic free radicals and immune effects. Early trials (e.g. NAVIFUS) suggest feasibility.
- **Laser Interstitial Thermal Therapy (LITT):** For deep-seated tumor foci, insert a fiberoptic laser probe under stereotactic guidance to thermally ablate tumor (as done in recurrent GBM). Monitor via MRI.

5. **Adjunctive Hyperthermia:** After RT sessions, apply whole or regional hyperthermia:

- **Magnetic nanoparticle hyperthermia:** Systemic IV infusion of iron-oxide nanoparticles (magnetic fluid). Wait for tumor uptake ($\sim 1\text{--}3$ h). Then place patient in AMF coil (as per Mayo study) to heat all nanoparticle-laden tissue to $45\text{--}50^{\circ}\text{C}$. Cooling blankets protect the body. This is repeated in multiple sessions (e.g. weekly) to damage residual tumor and sensitize to RT.
- **Whole-body hyperthermia:** In extreme cases (disseminated disease), consider extracorporeal perfusion heating of the patient to $\sim 42^{\circ}\text{C}$ (used in some melanoma protocols). High risk of organ stress.

6. **Tumor-Treating Fields (TTFields):** Patients will begin daily TTFields therapy (Optune device) as early as possible post-RT. Arrays are worn on the scalp (or body for extracranial tumors) delivering 200 kHz fields. Aim for ≥ 18 h/day use. TTFields will continue for months or until intolerable. This continual anti-mitotic therapy “plugs” remaining tumor cells.

7. **Immunotherapy and Immune Adjuvants:** Concurrent with all above, we aggressively modulate the immune response:

- **Checkpoint blockade:** Given that Grade IV tumors are immunosuppressive, we will trial combining PD-1 or CTLA-4 inhibitors. Radiotherapy and ablations release tumor antigens, which may synergise with ICI therapy.
 - **Vaccine/Oncolytic Virus:** If available, introduce personalised peptide vaccines or oncolytic viruses (e.g. engineered poliovirus) into the resection cavity. These extreme measures could convert the ablation site into an in situ cancer vaccine.
8. **Physiological Support:** Intensive nursing care is needed to support organ systems under this assault. Thermoregulation, hydration, and neurological monitoring are critical. Consider prophylactic steroids for brain edema, antiemetics, and antibiotics as needed.
9. **Monitoring and Adaptation:** Frequent MRI/PET imaging (weekly then monthly) assesses response. If tumors shrink insufficiently, additional focused ablations or dose boosts are applied. A flowchart of the decision process is shown below.

flowchart LR

```

A[Newly diagnosed Grade IV patient] --> B[Genomic and Physiological Profiling]
B --> C[Surgical Resection (maximal safe debulking)]
C --> D{Residual Tumor?}
D -->|Yes| E[Advanced Radiotherapy + Radiosensitisers]
D -->|No| F[Skip to E]
E --> G[Nanoparticle Radioenhancers (NBTXR3)]
G --> H[FLASH/Proton/Heavy-ion RT + Chemotherapy]
H --> I[Simultaneous Ablation Sessions]
I --> J(MRgFUS & LITT \n [Image-guided Thermal Ablation])
I --> K[Magnetic Hyperthermia]
I --> L[TTFields Device Therapy (continuous)]
I --> M[Immune Modulation (ICI, Oncolytic Virus)]
J & K & L & M --> N[Ongoing Assessment (MRI/PET, Biomarkers)]
N --> O{Tumor Response Adequate?}
O -->|No| P[Escalation: re-irradiation or experimental agents]
O -->|Yes| Q[Maintenance therapy \n (continue TTFields, chemo, follow-up)]

```




Figure 1: Proposed workflow for integrated treatment of Grade IV tumors.

After surgical debulking, patients receive dose-escalated RT with nanoparticle sensitizers, concurrent chemotherapy, and adjunctive physical therapies (focused ultrasound ablation, magnetic hyperthermia, and TTFs). Immune therapy is interwoven. Responses are evaluated iteratively to adapt the regimen (escalation or consolidation as needed).

Ethical and Legal Considerations

We note upfront that some proposed interventions border on unethical experimentation. Examples include: delivering **very high radiation doses** beyond standard constraints (risking brain necrosis), using **unapproved nanoparticles or viruses**, and depriving patients of potential quality of life for extended intensive therapy. In real life, these would require exceptional consent, institutional review board (IRB) oversight, and possibly crisis regulations. For this exercise we assume no legal limits, but we will *flag* all ethically problematic steps for discussion. Key issues: informed consent (patients must understand lethal risk), risk–benefit ratio, resource use, and long-term follow-up burden.

3. Preclinical Study Design

Prior to human trials, we design robust preclinical studies to evaluate safety and efficacy of the combined protocol:

- **Animal Models:** Use immunocompetent rodent models bearing orthotopic Grade IV gliomas (e.g. GL261 murine glioma in mice). Also consider genetically engineered mouse models (GEMMs) of GBM. For extracranial targets, use orthotopic xenografts in rabbit or swine for FUS/hyperthermia studies (due to skull size).
- **Treatment Groups:** Randomize to (1) standard RT only, (2) RT + one novel modality, (3) RT + multi-modal combo. Examples: RT+TTFields, RT+FUS, RT+hyperthermia, RT+all combined.
- **Endpoints:** Tumor control (MRI volumetrics), survival, cognitive function (rodent maze tests), and histology. Toxicities: brain edema, necrosis (histological sections), organ damage (bloodwork, pathology).
- **Mechanistic Studies:** Assess tumor oxygenation (pimonidazole staining), immune infiltration (flow cytometry for TILs), and genomic damage markers. Confirm that nanoparticle injection amplifies RT damage in vivo. Study FLASH vs conventional dose in animals (cognitive tests).
- **Safety:** Escalate each modality separately in dose-finding studies (e.g. find maximum tolerated ultrasound intensity). Evaluate co-toxicity: e.g. combining radiation with hyperthermia may increase blood–brain barrier permeability and neurotoxicity; quantify via serum biomarkers.

These preclinical trials will iteratively refine dosages and sequences. Endpoints will guide which combinations show synergy versus unacceptable toxicity. For example, if combining FUS+RT causes 100% mortality from edema in animals, that arm is re-designed (lower FUS energy or steroids).

4. Clinical Trial Design

Following encouraging preclinical results, we plan a multi-phase clinical development:

1. **Phase I (Safety and Feasibility):** A small cohort ($n \approx 10-20$) of patients with recurrent or newly diagnosed Grade IV glioma. Dose-escalation of one integrated regimen: e.g. proton-RT to 75 Gy + weekly magnetic hyperthermia + TTFields + standard chemo. Primary endpoints: dose-limiting toxicities (DLTs), maximum tolerated dose (MTD), feasibility of applying all modalities. Secondary endpoints: radiographic response, survival, neurocognitive function.
2. **Phase II (Signal-Finding):** Larger cohort ($n \approx 50$) to evaluate efficacy signals. Patients stratified by MGMT status, age, resection extent. Possibly randomize to (A) standard Stupp protocol vs (B) Stupp + one or more novel modalities from

Phase I. Endpoints: progression-free survival (PFS) at 6 months, overall survival (OS), quality of life. Biomarkers: serial liquid biopsy (cfDNA) to detect residual tumor, imaging of hypoxia pre- and post-treatment.

3. **Phase III (Comparative Efficacy):** A definitive trial if Phase II suggests benefit. Example design: standard chemo-RT vs experimental multi-modal arm. Large n (100+), with OS as primary outcome. (We expect this stage only if extreme protocol shows dramatically improved survival in Phase II).

Treatment Schedule (Example): We envision the following timeline for a patient (see Figure 2): Surgery (week 0). Weeks 1–2: planning (imaging, profiling). Weeks 3–10: concurrent proton FLASH-RT (25 fx of 3 Gy each) with temozolomide. Interdigitate: two hyperthermia sessions per week, and two FUS sessions (targeting residual tumor). TTFields device applied continuously from week 3 onward. Immune therapy (e.g. nivolumab) started week 4. Weeks 11–20: adjuvant chemotherapy (temozolomide cycles) with continued TTFields and monthly hyperthermia boosts. MRI surveillance every 8 weeks thereafter.




Figure 2: Example timeline of a combined treatment protocol.

After resection, the patient undergoes 8–10 weeks of intensified therapy (concurrent RT, TTFields, hyperthermia, etc.), followed by adjuvant treatment and long-term follow-up.

5. Safety and Risk Assessment

The proposed protocol carries high risk. We construct a risk matrix (Table 2) and address key concerns:

Table 2: Key safety risks and mitigation strategies for each major intervention.

Intervention	Major Risks	Mitigation/Safeguards
High-dose RT (≥ 75 Gy)	Neurocognitive decline, radionecrosis, leukoencephalopathy	Use protons/FLASH to spare normal tissue; strict dosimetry
FLASH RT	Unknown late effects, therapy machine failure	Begin with hybrid regimens; extensive QA on equipment
Carbon-ion RT	Unexpected severe CNS toxicity	Only in few-fraction settings; monitor carefully
Nanoparticles (NBTXR3)	Infarction if particles obstruct vessels, off-target radiation	Stereotactic injection; use tracers to confirm location
TTFields	Scalp dermatitis, patient compliance issues	Topical steroids for rash; patient education/support
MRgFUS (HIFU)	Brain edema, bleeding, off-target heating	Real-time MR-thermometry; ICU-level post-care
Magnetic Hyperthermia	Burns from induction coil; multi-organ stress; NP accumulation in liver/spleen	Cooling blankets; exclude patients with metal implants; start at low power
Combined therapies	Synergistic toxicity (e.g. radiation+heat causing widespread necrosis)	Stagger treatments; intensive monitoring; dose-escalation trial
Immunotherapy	Autoimmune reactions, cytokine storm	Standard immunosuppression protocols; close monitoring
Experimental biologics	Virus escape, off-target effects	Use attenuated vectors; compartmental administration

Key safety notes: We will use MRI/MR thermometry during FUS to prevent overheating. After each ablation or hyperthermia session, patients are observed in ICU to manage edema (steroids as needed). Neurocognitive function is tested serially; any decline prompts protocol review. All novel agents (NPs, viruses) require first-in-human safety trials before this combined protocol. In the unlikely event of catastrophic toxicity, we have defined stopping rules (e.g. any Grade 4 neurological event in $>10\%$ of patients halts escalation).

6. Expected Outcomes, Limitations, and Future Work

Expected Outcomes: Based on literature, we hypothesise dramatic tumor control improvement. For example, combining TTFields (5 mo OS gain) with proton dose-escalation (6–7% 2yr OS gain) and tumor-ablation could raise median survival well beyond current ~15 months. In an optimistic scenario, we might achieve 2–3 year median survival and 20–30% long-term survivors. We also expect reduction in tumor volume on MRI, and possibly remissions never seen with standard therapy alone.

Limitations: This extreme approach may incur severe side-effects. Neurocognitive damage from radiation, radiation necrosis requiring re-operation, organ dysfunction from systemic hyperthermia, and autoimmune disease from immunotherapy are all possible. Even if tumor is eradicated, quality of life may be poor. There is also the logistical limitation: assembling all modalities (particle centre, FUS unit, hyperthermia machine) at one centre is challenging. High cost and resource intensity are barriers. Technological feasibility (e.g. delivering FLASH with protons, which is under development) may not yet exist for routine use. Finally, we may encounter unknown biological interactions (e.g. combining certain therapies could promote an “antigen sink” effect or suppress immune response).

Future Work: Pending initial successes, future work includes: (1) integrating real-time imaging biomarkers (e.g. MRI spectroscopy, PET tracers) to adapt therapy mid-course; (2) developing “digital twin” patient models for personalized dose planning; (3) exploring gene editing (CRISPR) for tumor-specific targets in parallel; and (4) extending to other Grade IV cancers (e.g. high-grade sarcomas). If this protocol shows efficacy, it could shift the paradigm from palliation to potential cure in aggressive cancers.




Figure 3: Physics Modalities and Human-Body Constraints

7. Conclusion

We have outlined an ultra-aggressive, physics-driven multimodal treatment strategy for WHO Grade IV tumors. By combining advanced radiation (protons/FLASH), targeted hyperthermia (FUS, magnetic, photothermal), electrical fields, and immunomodulation, we aim to overcome the intrinsic resistance of these tumors. This integrative approach is rooted in recent primary evidence: for example, FLASH sparing normal tissue, nanoparticle radioenhancement, and TTFields extending survival. However, the strategy pushes the boundaries of safety and ethics. If carefully tested, it could provide a foundation for curing cancers that are currently considered incurable. All findings will need rigorous peer review and ethical scrutiny before any clinical adoption.

Declarations

Author contributions: Samuelson G. conceived the study, performed the literature review, and wrote the manuscript.

Funding: No funding was received for this work.

Conflict of interest: The author declares no conflicts of interest. Except my father diagnosed with glioblastoma.

Ethical approval: Not applicable.

Consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and material: Not applicable.

References

1. NICE. *Brain tumors (primary) and brain metastases in over 16s: management*. NICE Guideline (NG99), 2018. Recommendations 1.2.19–1.2.20: “Offer radiotherapy 60 Gy... with concomitant temozolomide”.
2. Green LM. *TTFields therapy shows clear survival benefit for glioblastoma*. Oncol Nurs News. April 3, 2017. (Describing EF-14 trial: “TTFields reduced risk of death by 37% and extended OS by a median of 5 months”).
3. Weiss J. *How nanoparticles enhance radiation therapy in lung cancer*. Targeted Oncology, Apr 22, 2025. (On NBTXR3 hafnium nanoparticles: “hafnium nanoparticles... absorb a photon and release ~9 electrons... localised amplification of radiation’s cytotoxic effect”).
4. Leporace M et al. *The Thermal Ablation with MRgFUS: From Physics to Oncological Applications*. Cancers 2025;17(1):36. (Review: “MRgFUS induces deep coagulation necrosis... encompassing the entire tumor site”; “In neuro-oncology... LIFU plus microbubbles to open BBB and enhance drug delivery”).
5. Focused Ultrasound Foundation. *Glioblastoma*. (Patient-oriented summary: lists advantages of focused ultrasound: noninvasive, image-guided BBB opening.)
6. Mayo Clinic News. Cordeiro B. *Mayo installs first magnetic nanoparticle hyperthermia system...* Feb 17, 2026. (Describes IV iron-oxide NPs heated by alternating magnetic fields; “The tumor absorbs the nanoparticles, effectively turning it into an induction pan... alternating magnetic fields pulse... creates heat”).
7. Ma X, Zhang NZ. *Tumor microenvironment as a key regulator of radiotherapy response*. Front. Immunol. 2026;17:1776636. (Review: “Tumor microenvironment encompasses... stromal and immune components... abnormal vasculature contributes to hypoxia and acidosis, which leads to radioresistance”).
8. Xiang S et al. *Ultrasound-assisted immunotherapy in cancer treatment*. Front. Immunol. 2025;16:1670527. (Review: “HIFU can generate localized hyperthermia... releasing tumor-associated antigens and HSPs... leading to expression of pro-inflammatory cytokines... activating adaptive immune responses”).

9. Knol M, et al. *FLASH radiotherapy against medulloblastoma*. 2026 preprint. (Background note: FLASH “minimizes complications... e-/photon/proton FLASH have demonstrated reduced cognitive decrements while maintaining anti-tumor efficacy”.)
10. Mehta MP et al. *NRG-BN001 trial (proton vs IMRT in GBM)*. ASTRO 2025. (Signal-finding: Proton 75 Gy arm had improved OS vs 60 Gy IMRT; 2-year OS 49.9% vs 43.1%.)