

ORIGINAL ARTICLE

Correlation between canal paresis and spontaneous nystagmus during early stage of acute peripheral vestibular disorders

HIROAKI FUSHIKI¹, MASAYUKI ISHIDA¹, SHIGEKI SUMI^{2,3}, AKIRA NARUSE¹ & YUKIO WATANABE¹

¹Department of Otolaryngology, Head & Neck Surgery, ²Division of Biostatistics and Clinical Epidemiology and ³Center for the Advancement of Medical Training, University of Toyama, Toyama, Japan

Abstract

Conclusions: This study demonstrates that the resolution period of spontaneous nystagmus (SN) may provide an indication of vestibular dysfunction on a particular day in the primary care setting. **Objective:** We aimed to predict canal paresis using fundamental observations of SN during the early stage of acute peripheral vestibular disorders. **Methods:** The study involved 87 patients who had recently experienced their first episode of acute spontaneous vertigo and direction-fixed horizontal nystagmus. Although they did not exhibit any other neurological deficits, they had been hospitalized with severe acute symptoms between 2004 and 2007. A correlation between the resolution period of SN and the results of laboratory caloric testing was reviewed. **Results:** The receiver operating characteristic analysis showed that the resolution period of SN may be a predictive indicator of unilateral vestibular hypofunction in the acute stage. In about half of the patients, SN disappeared on the third day after their initial visit. However, in 20% of the patients SN still persisted on the eighth day. Among the patients with SN, the prevalence of canal paresis increased with the increase in the resolution period of SN. When SN was observed on the fifth day, the prevalence was approximately 70%.

Keywords: Acute peripheral vestibulopathy, acute isolated vertigo, vestibular neuritis

Introduction

In acute peripheral vestibular disorders, patients usually suffer from acute symptoms of severe vertigo that persist longer than several hours, accompanied by intense nausea and vomiting. Almost all patients experience the onset of the disease as a surprise and are anxious [1]. The head thrust test or post-head-shaking nystagmus is known to be useful in predicting a significant unilateral vestibular hypofunction at the bedside of the patient [2]. However, it would be better to perform these dynamic vestibular tests in the sub-acute stage of the disease or later after spontaneous nystagmus (SN) and nausea/vomiting are reduced. During the early stage of acute peripheral vestibular disorder, oculomotor examination typically reveals nystagmus due to static vestibular imbalance by

showing the mixed horizontal-torsional and direction-fixed SN that does not change direction with gaze and is suppressed with fixation, in the absence of other neurological symptoms or signs [2,3]. This study aimed to predict canal paresis (CP) using fundamental observations of SN during the early stage of acute peripheral vestibular disorders to help primary care providers and their patients.

Material and methods

This study reviewed the data of patients who were admitted to the otolaryngology ward of the university hospital and its affiliates between 2004 and 2007 for severe vertigo, nausea and/or vomiting, and gait impairment. The patients were referred primarily to

Correspondence: Hiroaki Fushiki MD PhD, Department of Otolaryngology, Head & Neck Surgery, University of Toyama, 2630 Sugitani, Toyama-shi, Toyama 930-0194, Japan. Tel: +81 76 434 7368. Fax: +81 76 434 5038. E-mail: hfushiki@med.u-toyama.ac.jp

(Received 24 March 2010; accepted 23 May 2010)

ISSN 0001-6489 print/ISSN 1651-2251 online © 2010 Informa Healthcare
DOI: 10.3109/00016489.2010.497497

either an emergency physician or an otolaryngologist, depending on the time of the patient's visit. Acute peripheral vestibular disorder was defined based on clinical features that suggested a possibly acute isolated vertigo of peripheral origin and was independent of the results of caloric testing subsequently performed. The criteria for inclusion were as follows: (1) a single episode of spontaneous vertigo that persisted for several hours, (2) the presence of a direction-fixed and horizontal SN that did not change direction with gaze and was suppressed with fixation compared with when Frenzel glasses were in place (we have included both mixed horizontal-torsional type and horizontal type); (3) no history of inner ear disease; (4) no cochlear symptoms; (5) no headache; and (6) no central nervous system involvement, as measured by neurological physical examination and urgent brain CT.

In clinical practice, central vestibular disorders occasionally present with features of peripheral vestibular nystagmus and a negative neurologic examination at the onset of a vertigo attack [3]. Patients were excluded from this study if they revealed central nervous system disorders on subsequent brain MRIs and neurological examinations during their course of hospitalization. They were also excluded if they exhibited unilateral sensorineural hearing loss on pure-tone audiometry. Finally, 87 patients were included in the study (33 men and 54 women aged 25–89 years; 31 patients at the university hospital and 56 patients at affiliates; Table I). SN was assessed daily in the morning at the bedside. Patients were administered

anti-vertiginous and anti-emetic drugs without corticosteroids for several days to obtain relief from intense vertigo and nausea.

After nausea and vomiting were resolved, the patients were examined through routine laboratory evaluation of oculomotor-vestibular functions, which consisted of (1) spontaneous, positional, and gaze-evoked nystagmus; (2) pursuit, optokinetic system; and (3) caloric testing in sequence. The patients were also routinely evaluated using the sinusoidal rotational testing, galvanic body sway testing, and vestibular evoked myogenic potential testing in sequence [4]. Eye movements were recorded by the electro-nystagmographic (ENG) technique and sampled at 100 Hz with an A-D converter on a PC. Caloric stimuli consisted of alternate irrigation for 60 s with 6 L/min of cold and hot air, 24°C and 50°C, respectively. Asymmetry of vestibular function was calculated using the Jongkees formula. Significant CP corresponding to the evidence of unilateral vestibular dysfunction was defined as a response difference of 20% or more between the ears (laboratory-specific normal limits). The sinusoidal rotational test was performed with the patient's eyes closed in the dark with the head anteflexed to 30°. The patient was subjected to 0.1 Hz sinusoidal oscillations with a peak angular velocity of 75.4°/s. Directional preponderance (DP) was defined as a response difference of 10% or more between the ears (laboratory-specific normal limits).

In the present study, we aimed to predict the prevalence of unilateral vestibular hypofunction on a particular day in patients with SN using the resolution period of bedside SN (counted from the first visit) in patients with peripheral vestibules. First, receiver operating characteristic (ROC) analysis was performed to determine the predictability of the resolution period measures of SN; CP on laboratory caloric test was regarded as the gold standard. The area under the ROC curve (AUC) was estimated by the trapezoidal rule. An AUC > 0.60 was considered a highly predictive indicator. Second, the cumulative percentage of patients with SN in the early stage was examined. Third, among the patients with SN, the prevalence of CP with respect to a hospital stay was examined.

The duration of subjective symptoms was also examined. The relationship between the resolution of subjective symptoms and bedside SN was analyzed using Pearson's product-moment correlation coefficient. A correlation coefficient between 0.20 and 0.40 was considered mild correlation and 0.40–0.70 moderate correlation. A difference with a *p* value of < 0.05 was considered statistically significant.

Table I. Summary of patients with acute peripheral vestibular disorder.

Parameter	Value
Subjects (<i>n</i> = 87)	
Sex (male/female)	33/54
Age (years)	61.5 ± 14.3
Acute symptoms	
Vertigo sensation (<i>n</i> = 87)	6.9 ± 7.6 days
Nausea and/or vomiting (<i>n</i> = 79)	3.1 ± 2.9 days
Difficulty in walking (<i>n</i> = 76)	3.0 ± 1.8 days
Findings	
Bedside test	
Nystagmus duration (<i>n</i> = 87)	5.3 ± 5.7 days
Laboratory vestibular test	
CP in caloric test	48.60% (35/72)
DP in rotational test	30.30% (10/33)

CP, canal paresis; DP, directional preponderance.

Results

The visit was during the daytime (9:00 am to 5:00 pm) in 42% of the patients, during the evening or night (5:00 pm to 3:00 am) in 46%, and during the early morning (3:00 am to 9:00 am) in 12%. ROC analysis was performed to determine the predictability of the resolution period measures of SN (Figure 1). We found that the resolution period of SN could be a predictive indicator of unilateral vestibular hypofunction in the acute stage (AUC 0.63).

Figure 2A shows the cumulative percentage of patients with SN in the early stage. In about half of the patients, SN disappeared on the third day after the initial visit. However, in 20% of the patients SN still persisted on the eighth day. The prevalence of CP among the patients with SN is shown in Figure 2B. The prevalence increased with the increase in the duration of SN. For instance, when SN was observed on the third day, the prevalence was approximately 60%, and when SN was observed on the fifth day, the prevalence was approximately 70%. The duration of bedside SN did not correlate properly with DP in the subsequent laboratory rotational test (AUC 0.55).

Of the acute symptoms, nausea and/or vomiting and gait impairment resolved relatively quickly in most patients, but the resolution period of the vertigo




Figure 1. The receiver operating characteristic (ROC) curve indicates that the resolution period of spontaneous nystagmus in the acute stage may be a predictor of canal paresis (CP) in the caloric test. The area under the ROC curve was 0.63.




Figure 2. (A) The cumulative percentage of patients with acute peripheral vestibular disorder and spontaneous nystagmus (SN) with respect to the length of hospital stay in the acute stage ($n = 87$). On the third day from the initial visit, SN disappeared in about half of the patients; 20% of the patients still had SN on the eighth day. (B) The prevalence of canal paresis (CP) among these patients with SN ($n = 72$). These findings suggest that when SN is observed on the ninth day, the estimated prevalence rate of CP is approximately 90%. Zero indicates the date of admission.




Figure 3. Relationships between resolution periods of acute symptoms and duration of bedside spontaneous nystagmus (SN). (A) Vertigo sensation. (B) Nausea and vomiting. (C) Difficulty in walking. Patients with significant canal paresis (CP) (●) and those without CP (○) are plotted separately. Longer duration of bedside SN correlated with longer resolution of vertigo sensation.

sensation varied widely among the patients (Table I and Figure 3). Correlations were observed between nausea/vomiting and the other two symptoms (correlation coefficient: vertigo sensation 0.403, $p < 0.001$; difficulty in walking 0.673, $p < 0.001$). Longer SN duration correlated with a longer resolution period of vertigo sensation (correlation coefficient 0.357, $p = 0.002$). No correlations were found between SN and the other two symptoms (correlation coefficient: nausea/vomiting -0.005 , $p = 0.969$; difficulty in walking 0.110, $p = 0.390$).

Discussion

In clinical practice, a patient with a first episode of acute vertigo is often referred to a non-specialist, such as a general or an emergency physician, depending on

the medical context among different countries and districts [5–7]. A large office-based survey by the National Ambulatory Medical Care reported that almost 70% of all outpatients with dizziness, who were aged 25 years and older, were initially examined by primary care physicians, such as general practitioners, family physicians, and general internists, and only 10% were examined by specialists such as otolaryngologists and neurologists [5]. Of all the patients who were referred to primary care physicians with the chief complaint of dizziness, 1.5% were hospitalized and 4.4% were referred to a specialist [5]. Vertigo/dizziness accounts for 2.5–3.5% of all the visits to the emergency departments [6,7]. Several reports have reported a clinically valid management of neurological emergencies; the focus would usually be on determining whether vertigo is related to a life-threatening condition [8–10]. Once a diagnosis of

vertigo of peripheral origin is made, it is best to estimate the evidence of vestibular hypofunction as the next step.

SN, characterized as horizontal, direction-fixed, and suppressed with fixation, in the absence of other neurological symptoms or signs reflects an imbalance in tonic neural activity in the vestibular nuclei due to sudden loss of input unilaterally from the labyrinth or the vestibular nerve. This study indicates that the resolution period of bedside SN could provide an indication of CP on a particular day. For instance, if SN presents on the fifth day after onset, the suspicion of it being CP is estimated to be approximately 70% (Figure 2B). A prospective study with 6 months follow-up showed that the degree of asymmetry of the caloric CP and age would influence balance performance and perceived symptoms after acute unilateral vestibular disorders [11]. A primary care physician may consult a specialist regarding the confirmation and degree of CP by means of a laboratory caloric test and additional treatment on the third day or later. A specific exercise improves residual imbalance and persistent dizziness [12–14]. Corticosteroids may be a treatment option because their administration in the early stage may improve the recovery of peripheral vestibular function in the long term [15] and facilitate central vestibular compensation [16,17].

Recently, Yagi et al. [18] analyzed the rotation axis of SN during the early stage of vestibular neuritis by means of an infrared charge-coupled device (CCD) camera. They determined that the rotation axis corresponds to the pathology of the superior vestibular nerve branch in all patients examined. In general, the axis initially tends to be mixed horizontal-torsional, and subsequently its orientation gradually changes during the recovery course of the early stage, as the anterior canal branch recovers faster than the horizontal canal branch, and ultimately draws closer to the horizontal [18]. This phenomenon is noticeable when the peripheral vestibular nystagmus is carefully observed using an infrared CCD camera. As mentioned in the Methods section, it is important to keep in mind that central vestibular disorder occasionally presents with a direction-fixed pure horizontal nystagmus [19,20] and, although rare, presents with a direction-fixed mixed horizontal-torsional nystagmus [21].

This study also examined symptomatic outcomes of patients who had experienced the first episode of acute isolated vertigo and direction-fixed horizontal nystagmus with symptoms severe enough to warrant hospitalization. Of the acute symptoms, nausea and/or vomiting (vestibulo-autonomic response) and gait impairment (vestibulo-spinal response) resolved relatively quickly in most patients, but the resolution period of vertigo (vestibular perception) varied widely

among the patients. The duration of bedside SN (vestibulo-ocular response) was associated with the resolution of vertigo sensation, whereas it was not associated with the other two symptoms.

The presence of SN in vestibular disorders is a fundamental examination, which can be observed at the bedside in a matter of minutes without difficulty. In conclusion, observation of SN during the early stage of acute peripheral vestibular disorders provides useful information to non-specialists in predicting vestibular dysfunction, the timing of consultation, and planning of additional treatment.

Acknowledgments

We would like to thank Professor H. Origasa for advice regarding statistical analyses. This work was supported by a grant from the Ministry of Health, Labor and Welfare of Japan.

Declaration of interest: The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

References

- [1] Godemann F, Linden M, Neu P, Heipp E, Dörr P. A prospective study on the course of anxiety after vestibular neuronitis. *J Psychosom Res* 2004;56:351–4.
- [2] Fetter M. Assessing vestibular function: which tests, when? *J Neurol* 2000;247:335–42.
- [3] Leigh RJ, Zee DS. 2006. Nystagmus due to vestibular imbalance. *The neurology of eye movements*. Fourth edition. New York: Oxford University Press. p. 480–93.
- [4] Watanabe Y, Takeda S, Mizukoshi K, Imamura J. Clinical usefulness of the personal computer for equilibrium examinations. *Acta Otolaryngol Suppl* 1993;504:98–105.
- [5] Sloane PD. Dizziness in primary care. Results from the National Ambulatory Medical Care Survey. *J Fam Pract* 1989;29:33–8.
- [6] Kerber KA, Meurer WJ, West BT, Fendrick AM. Dizziness presentations in U.S. emergency departments, 1995–2004. *Acad Emerg Med* 2008;15:744–50.
- [7] Crespi V. Dizziness and vertigo: an epidemiological survey and patient management in the emergency room. *Neurol Sci Suppl* 2004;25:24–5.
- [8] Moeller JJ, Kurniawan J, Gubitza GJ, Ross JA, Bhan V. Diagnostic accuracy of neurological problems in the emergency department. *Can J Neurol Sci* 2008;35:335–41.
- [9] Halmagyi GM. Diagnosis and management of vertigo. *Clin Med* 2005;5:159–65.
- [10] Seemungal BM. Neuro-otological emergencies. *Curr Opin Neurol* 2007;20:32–9.
- [11] Kammerlind AS, Ledin TE, Odkvist LM, Skargren EI. Influence of asymmetry of vestibular caloric response and age on balance and perceived symptoms after acute unilateral vestibular loss. *Clin Rehabil* 2006;20:142–8.
- [12] Tokumasu K, Fujino A, Noguchi H. Prolonged dysequilibrium in three cases with vestibular neuronitis: efficacy of

- vestibular rehabilitation. *Acta Otolaryngol Suppl* 1993;503: 39–46.
- [13] Asai M, Watanabe Y, Shimizu K. Effects of vestibular rehabilitation on postural control. *Acta Otolaryngol Suppl* 1997; 528:116–20.
- [14] Strupp M, Arbusow V, Maag KP, Gall C, Brandt T. Vestibular exercises improve central vestibulospinal compensation after vestibular neuritis. *Neurology* 1998;51:838–44.
- [15] Strupp M, Zingler VC, Arbusow V, Niklas D, Maag KP, Dieterich M, et al. Methylprednisolone, valacyclovir, or the combination for vestibular neuritis. *N Engl J Med* 2004;351: 354–61.
- [16] Cameron SA, Dutia MB. Lesion-induced plasticity in rat vestibular nucleus neurones dependent on glucocorticoid receptor activation. *J Physiol* 1999;518:151–8.
- [17] Strupp M, Arbusow V, Brandt T. Exercise and drug therapy alter recovery from labyrinth lesion in humans. *Ann N Y Acad Sci* 2001;942:79–94.
- [18] Yagi T, Koizumi Y, Sugizaki K. 3D analysis of spontaneous nystagmus in early stage of vestibular neuritis. *Auris Nasus Larynx* 2010;37:167–72.
- [19] Barton JJ, Sharpe JA. Oscillopsia and horizontal nystagmus with accelerating slow phases following lumbar puncture in the Arnold-Chiari malformation. *Ann Neurol* 1993;33: 418–21.
- [20] Kumar A, Patni AH, Charbel F. The Chiari I malformation and the neurotologist. *Otol Neurotol* 2002;23:727–35.
- [21] Lee H, Cho YW. A case of isolated nodulus infarction presenting as a vestibular neuritis. *J Neurol Sci* 2004 15; 221:117–19.

10th EFAS Congress

JUNE 22–25, 2011 WARSAW, POLAND



Prof. Henryk Skarżyński
President of the EFAS

10th Congress of the European Federation of Audiology Societies
and the Conference of the Audiology and Phoniatrics Sections of the
Polish Society of Otorhinolaryngologists and Head and Neck Surgeons

Special events on June 22, 2011

- 2nd International Congress of Young Scientists and Students
(www.congress.ifps.org.pl)
- Window Approach Workshop (WAW) presenting live middle ear and cochlear implant surgeries

ABSTRACT SUBMISSION FOR THE 10TH EFAS CONGRESS IS NOW OPEN!

Abstract submission deadline – **February 28, 2011**







TAKE ADVANTAGE OF OUR EARLY BIRD RATES AND REGISTER ON-LINE BEFORE DECEMBER 15, 2010

ORGANIZERS

Institute of Physiology and Pathology of Hearing
International Center of Hearing and Speech



The Audiology and Phoniatrics Sections of the Polish Society
of Otorhinolaryngologists and Head and Neck Surgeons

www.efas2011.org

Copyright of Acta Oto-Laryngologica is the property of Taylor & Francis Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.