Wiqaya: Wildfire Susceptibility Mapping and Alerts for the Forests of Northern Morocco

Sanae Belfrouh · Wiame Kane · Yassine Azmou · Mohammed Yassine Sof · Hajar Elouatiq

Team Wiqaya — Morocco

Arab Youth Space Hackathon 2026 · Proof of Concept

Theme: Climate & Disaster Resilience

October 2026 · github.com/SanaeBelfrouh/wiqayagit

The western Rif burns almost every summer, and warnings arrive after ignition

- **Recent large fires:** Bouhachem 2022 (> 7,500 ha), Larache 2022 (> 10,500 ha), Derdara 2021 and Bouhachem 2020 (~1,100 ha each), two fires in August 2025
- **Drivers:** hot, dry chergui winds, steep slopes, villages (douars) inside the forest
- **Gap:** danger bulletins are province-scale and satellite detections come after ignition; there is no daily, cell-level view of risk
- **Why satellite data:** NASA FIRMS gives the only consistent 20-year fire record of these remote mountains (MODIS since 2005, VIIRS 375 m since 2017)




Fig. 1. Study area (25,900 ha) and NASA FIRMS detections by period. Population: WorldPop 2017.

10,600
people living inside the forest

264,000
people within 10 km of it

445
FIRMS detections inside the forest since 2005

Business use case: where to patrol today, and whether to warn residents


End users	Forest managers (national forestry agency, ANEF), Civil Protection, provinces and communes; residents of the villages in and around the forest
Decision supported	Each summer morning: where to send patrols and first-intervention teams, and whether the day is dangerous enough to warn residents and restrict burning
Current practice	Province-scale danger bulletins, field knowledge, and fire detections (FIRMS, EFFIS) that arrive after ignition; no alerts in Arabic and French
Contribution of Wiqaya	250 m risk map with alert zones and a per-cell explanation, daily weather danger index, 20-year fire history, SMS / email alerts in three languages, open GeoJSON API
Business model	Subscription per protected area for institutions; free citizen alerts funded by these contracts or CSR partners; open data, so no imagery cost

Data: open Earth-observation and ancillary datasets

Dataset	Product and provider	Period	Use
Active fires	NASA FIRMS: MODIS C6.1 and VIIRS S-NPP 375 m	2005–2026 (1,176 points)	Labels, fire history
Terrain	Digital elevation model, ≈30 m	static	Elevation, slope, aspect
Population	WorldPop mar_ppp_2017 (CC BY 4.0)	2017	Exposure per cell
Weather	Open-Meteo: ERA5 archive and forecast (CC BY 4.0)	2017–today	Weather danger index
Optical imagery *	Sentinel-2 L2A, Landsat C2 L2 (Planetary Computer)	2005–2025	NDVI, NDMI, NBR, dNBR
Land cover, OSM *	2021 classification; OpenStreetMap 2022	2021–2022	Fuel, human pressure

* *Research notebook only.* **Hyperspectral data is not used in this proof of concept;** EnMAP or Arab Satellite 813 scenes are planned to map live fuel moisture and vegetation species.

Approach: from satellite detections to alert zones



Reproducible: notebooks/02_main_analysis.ipynb runs end to end in about 5 minutes from the committed sample inputs (fixed seeds, relative paths).

Risk classes on 1 August 2025: the 2025 fires fell outside the alert zones




Fig. 2. Left: five risk classes; outlined cells are alert zones (riskiest 20 %); blue points are the 2025 FIRMS fires (blind test). Right: years since the last FIRMS fire (20 = none since 2005). Notebook output results/example_output.png.

Population, terrain and fire history drive the selected model




Fig. 3. Absolute standardised coefficients of the logistic regression.

- **Selected model:** L2 logistic regression (C = 0.5, balanced classes)
- **Strongest predictors:** population, elevation, east-facing aspect, then fire history (years since the last fire, burned in the last 5 years)
- **Weak predictor:** slope
- **Missing drivers:** vegetation dryness and human access, available only in the research notebook

The same model is delivered to users through a web platform




Fig. 4. Daily risk map with the weather danger of the day and the five-class legend.




Fig. 5. Fire history: 25 July 2022, start of the Bouhachem fire; FIRMS shows 5 detections, 4 inside the forest.

Access for residents: Arabic, French and English, on any phone




Fig. 6. Right-to-left Arabic interface.




Fig. 7. Fire history and SMS / email alert sign-up on a phone.

Validation: informative on large-fire years, but the 2025 blind test fails

Test	Result
Leave-one-year-out 2017–2024, logistic regression (selected)	Mean ROC-AUC per year 0.62; global 0.65; PR lift 1.64
Leave-one-year-out 2017–2024, random forest	Mean ROC-AUC per year 0.56; global 0.60
ROC-AUC per held-out year	2018: 0.99 · 2019: 0.88 · 2020: 0.80 · 2021: 0.68 · 2023: 0.23 · 2024: 0.06
Blind test, 2025 fires (39 cells)	**ROC-AUC 0.25; no fire cell in the alert zones**

- **Works** on the large fire years 2018–2021
- **Fails** on 2023–2025: the 2025 fires started in cells rated low risk
- **Cause:** predictors describe exposure, terrain and history, not vegetation dryness or human access
- **Few positives:** 345 fire cell-years; per-year scores are unstable
- **No leakage:** history strictly before each date; 2025 never used for training or selection

Expected impact and strategic alignment

Goal	Contribution
SDG 13 Climate action (13.1)	Resilience to heatwave- and wind-driven fire seasons
SDG 15 Life on land (15.1, 15.2)	Patrols and firebreak work targeted at the riskiest cells
SDG 11 Sustainable communities (11.5)	Early warning for ~10,600 people living inside the forest

- **Morocco:** supports the strategy Forêts du Maroc 2020–2030, ANEF and Civil Protection
- **Arab region:** open data only; replicable in Mediterranean forests of Algeria, Tunisia, Lebanon and Syria
- **Inclusion:** Arabic-first interface; SMS for residents without smartphones

Next steps for incubation

1. **Better predictors:** land cover, OpenStreetMap distances and daily Sentinel-2 NDVI, NDMI, NBR
2. **Hyperspectral:** EnMAP / Arab Satellite 813 maps of live fuel moisture and species
3. **Better reference data:** validation on dNBR burned areas, not only FIRMS points
4. **Operations:** SMS gateway, daily batch processing, alert workflow with forest managers
5. **Pilot:** one fire season in Bouhachem, Derdara and Jbel Sougna, then other Mediterranean forests

Reproduce the headline result

github.com/SanaeBelfrouh/wiqayagit

```
pip install -r requirements.txt
jupyter lab notebooks/02_main_analysis.ipynb
```

About 5 minutes on a standard laptop

References and data sources

Assali, F. et al. (2016). Forest-fire risk mapping in the Chefchaouen-Ouazzane region, Morocco (topoclimate index).

Hersbach, H. et al. (2020). The ERA5 global reanalysis. Q. J. R. Meteorol. Soc., 146, 1999–2049.

NASA FIRMS. MODIS Collection 6.1 and VIIRS S-NPP 375 m active fire detections. https://earthdata.nasa.gov/firms

WorldPop (2018). Morocco 2017 population counts (mar_ppp_2017). University of Southampton. CC BY 4.0.

Zippenfeld, P. Open-Meteo.com Weather API. CC BY 4.0. https://open-meteo.com

Copernicus Sentinel-2 L2A data (2015–2025) and USGS Landsat Collection 2 Level-2, accessed via Microsoft Planetary Computer.

© OpenStreetMap contributors (ODbL). Basemaps: Esri World Topo Map; OpenTopoMap (CC BY-SA).

Software: GeoPandas, Rasterio, rasterstats, scikit-learn, FastAPI, Leaflet. Code released under the MIT licence.