

Week 8

Isoperimetric Inequalities and Applications

LECTURER: SIVA ATHREYA

SCRIBE: ATREYA CHOUDHURY, SRIVATSA BALAJI

The focus of this chapter is to look at how the geometry of weighted graph affects the properties of the corresponding random walk on it.

Definition 8.0.1 (Isoperimetric Inequality). *Let $A, B \subseteq V$, $\mu_E(A, B) = \sum_{x \in A} \sum_{y \in B} \mu_{xy}$ and $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be an increasing function.*

(Γ, μ) is said to satisfy the ψ -isoperimetric inequality if $\exists c_0 > 0$ such that

$$\frac{\mu_E(A, V \setminus A)}{\psi(\mu(A))} \geq \frac{1}{c_0} \quad \forall A \subseteq V \text{ and } |A| < \infty$$

If a weighted graph satisfies the ψ -isoperimetric inequality, we say it has the I_ψ property.

A graph is said to have the property I_α for $\alpha \in [0, \infty)$ when $\psi(t) = t^{1-\frac{1}{\alpha}}$ and said to have the property I_∞ when $\psi(t) = t$

Example. \mathbb{R}^d . We look at $A = B(0, r)$

$$S_B \equiv \text{surface area of } A = c_d r^{d-1}$$

$$V_B \equiv \text{volume of } A = \tilde{c}_d r^d$$

$$\therefore \frac{S_B}{V_B^{\frac{d-1}{d}}} \geq \frac{1}{c_0}$$

We can take $\psi(t) = t^{1-\frac{1}{d}}$

Show that \mathbb{R}^d has the I_d property for all such A such that $|A| < \infty$

Example. \prod_2 , the binary tree has the I_∞ property with $c_0 = 3$

Observations. *If (Γ, μ) satisfies $I_{\alpha+\delta}$, then it satisfies I_α*

Definition 8.0.2 (Nash Inequality). $\alpha \in [1, \infty)$, (Γ, μ) is said to have the property N_α if $\forall f \in \mathbb{L}^1(V) \cap \mathbb{L}^2(V)$,

$$\mathcal{E}(f, f) \geq C_N \|f\|_1^{-\frac{4}{\alpha}} \|f\|_2^{2+\frac{4}{\alpha}}$$

Remark. 1. (Γ, μ) satisfies I_α for $\alpha \in [1, \infty) \implies (\Gamma, \mu)$ satisfies N_α
 2. \mathbb{Z}^d satisfies N_α

Theorem 8.0.1. *Let $\alpha \geq 1$. Then the following are equivalent*

1. (Γ, μ) satisfies N_α
2. $\exists C_H > 0$ such that

$$p_n(x, x) \leq \frac{C_H}{(n \vee 1)^{\frac{\alpha}{2}}} \quad \forall n \geq 0 \text{ and } x \in V$$

3. $\exists C'_H > 0$ such that

$$p_n(x, y) \leq \frac{C'_H}{(n \vee 1)^{\frac{\alpha}{2}}} \quad \forall n \geq 0 \text{ and } x, y \in V$$

Corollary 8.0.1. 1. Suppose (Γ, μ) satisfies I_α . Then, $\exists C > 0$ such that

$$p_n(x, y) \leq \frac{C}{(n \vee 1)^{\frac{\alpha}{2}}} \quad \forall n \geq 0 \text{ and } x, y \in V$$

2. Let V be infinite and $\mu_{xy} \geq c_0 > 0 \ \forall x \sim y$. Then, $\exists C_1 > 0$ such that

$$p_n(x, y) \leq \frac{C_1}{(n \vee 1)^{\frac{1}{2}}} \quad \forall n \geq 0 \text{ and } x, y \in V$$

Remark. 1. $p_n(x, x) \equiv$ on-diagonal bounds

2. Theorem provides global upper bounds
3. part b of corollary 8.0.1 applied to $V = \mathbb{Z}$
 \implies the shortest possible on-diagonal upper bounds with natural weights
4. Let $\Gamma = \mathbb{Z}^d$ have natural weights $\mu_{xy}^{(0)}$ and $\Gamma' = \mathbb{Z}^d$ have natural weights $\mu_{xy}^{(1)}$ such that $\mu_{xy}^{(1)} \geq c_0 \mu_{xy}^{(0)}$ Let $(\Gamma, \mu^{(0)})$ satisfy N_d
 $\implies (\Gamma', \mu^{(1)})$ satisfies N_d
 \implies the upper bound of the theorem holds
5. $\Gamma = \mathbb{Z}^d \cup_{(0, \dots, 0)} \mathbb{Z}^d$
 $\implies \Gamma$ also satisfies N^d
6. 8.0.1 does not give us any information on upper bounds when we fix $n \geq 0$ and let $d(x, y)$ get large.

Theorem 8.0.2. *Let (Γ, μ) be a weighted graph. Then,*

$$p_n(x, y) \leq \frac{2}{\sqrt{\mu_x \mu_y}} e^{-\frac{d(x, y)^2}{2n}} \quad \forall x, y \in V \text{ and } n \geq 1$$

Example. Consequences for \mathbb{Z}^d

We expect

$$p_n(x, y) \leq \frac{c_1}{n^{\frac{d}{2}}} e^{-c_2 \frac{d(x, y)^2}{n}}$$

$$\mathbb{Z}^d \text{ satisfies } I_d \implies \mathbb{Z}^d \text{ satisfies } N_d \xRightarrow{8.0.1} p_n(x, y) \leq \frac{c}{n^{\frac{d}{2}}} \quad \forall x, y \in V \text{ and } n \geq 1$$

$$\therefore p_n(x, y) \leq \frac{c}{n^{\frac{d}{2}}} \leq \frac{c}{n^{\frac{d}{2}}} e^{-\frac{d(x, y)^2}{n}} \quad \text{when } d(x, y) \leq \sqrt{n}$$

When, $d(x, y) \geq \sqrt{2dn \log n}$,

$$p_n(x, y) \leq c_1 e^{-\frac{d(x, y)^2}{n}} = c_1 e^{-\frac{2c_2}{4} \frac{d(x, y)^2}{n}} e^{-\frac{2c_2}{4} \frac{d(x, y)^2}{n}} \leq \frac{\tilde{c}_1}{n^{\frac{d}{2}}} e^{-\frac{c_2^2 d(x, y)^2}{n}}$$

Definition 8.0.3. (Γ, μ) is said to have **polynomial volume growth** if $\exists C_V$ and θ such that

$$\max\{|B(x, r)|, \mu(B(x, r))\} \leq C_V r^\theta \quad \forall x \in V \text{ and } r \geq 1$$

Lemma 8.0.1. (Γ, μ) has polynomial volume growth with index θ . Then,

$$\mathbf{P}^x(d(x, X_n) > r) \leq cr^\theta e^{-\frac{r^2}{4n}}$$

This implies $\exists c_2 > 0$ such that

$$d(x, X_n) \leq c_2 \sqrt{n \log n} \quad \forall \text{ large } n \text{ w.p. } 1$$

Proof. We define $\mathcal{D}_k = B(x, 2^k r) \setminus B(x, 2^{k-1} r)$

$$\begin{aligned} \mathbf{P}^x(d(x, X_n) > r) &\stackrel{Ex}{=} \sum_{k=1}^{\infty} \sum_{y \in \mathcal{D}_k} p_n(x, y) \mu_x \\ &\leq \sum_{k=1}^{\infty} \sum_{y \in \mathcal{D}_k} \frac{2}{\sqrt{\mu_x}} \sqrt{\mu_y} e^{-\frac{(2^{k-1} r)^2}{2n}} \\ &= \sum_{k=1}^{\infty} \frac{2}{\sqrt{\mu_x}} e^{-\frac{(2^{k-1} r)^2}{2n}} \sum_{y \in \mathcal{D}_k} \sqrt{\mu_y} \\ &\leq \sum_{k=1}^{\infty} \frac{2}{\sqrt{\mu_x}} e^{-\frac{(2^{k-1} r)^2}{2n}} \sqrt{|\mathcal{D}_k|} \sqrt{\mu(\mathcal{D}_k)} \\ &\leq \sum_{k=1}^{\infty} \frac{2}{\sqrt{\mu_x}} e^{-\frac{(2^{k-1} r)^2}{2n}} c(2^k r)^\theta \end{aligned}$$

□