

S

Spatial Applications of Matrices

TABLE OF CONTENTS

		Page
§S.1.	Points, Planes	S-3
§S.2.	Lines	S-3

In this Appendix we summarize some geometric applications of matrices in 3D space. Indexed homogeneous Cartesian coordinates¹ $\{x_0, x_1, x_2, x_3\}$ are used. To pass to physical coordinates, divide x_1, x_2 and x_3 by x_0 : $\{x_1/x_0, x_2/x_0, x_3/x_0\}$. If $x_0 = 0$ the physical coordinates are at infinity.

§S.1. Points, Planes

Points in 3D space will be identified by X, Y, P, Q , etc. Their coordinates are put in the 4-vectors

$$\mathbf{x} = [x_0 \ x_1 \ x_2 \ x_3]^T, \quad \mathbf{y} = [y_0 \ y_1 \ y_2 \ y_3]^T, \quad \mathbf{p} = [p_0 \ p_1 \ p_2 \ p_3]^T, \text{ etc} \quad (\text{S.1})$$

Planes in 3D space will be identified by A, B, C , etc. The equation of plane A is written $a_0x_0 + a_1x_1 + a_2x_2 + a_3x_3 = 0$ or

$$\mathbf{a}^T \mathbf{x} = 0, \quad \text{or} \quad \mathbf{x}^T \mathbf{a} = 0 \quad (\text{S.2})$$

The plane A is thus defined by the 4-vector \mathbf{a} .

Example S.1. Find the coordinates of the point where line joining points P and Q intersects plane A .

Solution. Any point of PQ is R where

$$\mathbf{r} = \lambda \mathbf{p} + \mu \mathbf{q} \quad (\text{S.3})$$

If R is on plane A , then $\mathbf{a}^T \mathbf{x} = 0$ so that $\lambda \mathbf{a}^T \mathbf{p} + \mu \mathbf{a}^T \mathbf{q} = 0$. Absorbing a suitable multiplier into the coordinates of R we obtain its vector in the form

$$\mathbf{r} = (\mathbf{a}^T \mathbf{q}) \mathbf{p} - (\mathbf{a}^T \mathbf{p}) \mathbf{q}. \quad (\text{S.4})$$

§S.2. Lines

Let \mathbf{x} and \mathbf{y} be the coordinates of points X and Y on a given line L . The 4×4 antisymmetric matrix

$$\mathbf{L} = \mathbf{xy}^T - \mathbf{yx}^T \quad (\text{S.5})$$

is called the *coordinate matrix* of the line. It can be shown that \mathbf{L} determines the line L to within a scale factor.

Let A and B be two planes through L . The 4×4 antisymmetric matrix

$$\mathbf{L}^* = \mathbf{ab}^T - \mathbf{ba}^T \quad (\text{S.6})$$

is called the *dual coordinate matrix* of line L . It can be shown that²

$$\mathbf{L}^* \mathbf{L} = \mathbf{0}. \quad (\text{S.7})$$

¹ These coordinates were independently invented in 1827 by Møbius and Feuerbach, and further developed in 1946 by E. A. Maxwell at Cambridge. See E. A. Maxwell, *General Homogeneous Coordinates in Space of Three Dimensions*, Cambridge University Press, 1951.

² See E. A. Maxwell, loc. cit., page 150.

Example S.2. Find where line L defined by two points X and Y meets a plane A .

Solution. Consider the vector

$$\mathbf{p} = \mathbf{L}\mathbf{a} = (\mathbf{xy}^T - \mathbf{yx}^T)\mathbf{a} = (\mathbf{y}^T\mathbf{a})\mathbf{x} - (\mathbf{x}^T\mathbf{a})\mathbf{y} \quad (\text{S.8})$$

From the last form the point P of coordinates \mathbf{p} must lie on the line that joins X and Y . Moreover since \mathbf{L} is antisymmetrical, $\mathbf{a}^T\mathbf{L}\mathbf{a} = 0$ so that $\mathbf{a}^T\mathbf{p} = 0$. Thus P is the intersection of line L and plane A .

Example S.3. Find the plane A that joins line L to a point X .

Solution.

$$\mathbf{a} = \mathbf{L}^* \mathbf{x} \quad (\text{S.9})$$

where \mathbf{L}^* is the dual of \mathbf{L} . The demonstration is trivial.

Two immediate corollaries: (i) Line L lies in the plane A if $\mathbf{L}\mathbf{a} = \mathbf{0}$; (ii) Line L passes through the point X if $\mathbf{L}^*\mathbf{x} = \mathbf{0}$.

Example S.4.

Consider two lines L_1 and L_2 with coordinate matrices $\mathbf{L}_1, \mathbf{L}_2$ and dual matrices $\tilde{\mathbf{L}}_1$ and $\tilde{\mathbf{L}}_2$, respectively. Find the conditions for the lines to intersect.

Solution. Any of the four equivalent conditions

$$\mathbf{L}_1\mathbf{L}_2^*\mathbf{L}_1 = \mathbf{0}, \quad \mathbf{L}_1^*\mathbf{L}_2\mathbf{L}_1^* = \mathbf{0}, \quad \mathbf{L}_2\mathbf{L}_1^*\mathbf{L}_2 = \mathbf{0}, \quad \mathbf{L}_2^*\mathbf{L}_1\mathbf{L}_2^* = \mathbf{0}. \quad (\text{S.10})$$

For the proof see E. A. Maxwell, loc. cit, page 154.