

# Adversarial LQ Mean-Field Games over Multigraphs

Muhammad Aneeq uz Zaman, Sujay Bhatt and Tamer Başar



I L L I N O I S

UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN

# Opinion Dynamics on a multigraph

# Opinion Dynamics on a multigraph




Figure: Multigraph of 6 agents

# Opinion Dynamics on a multigraph




Figure: Multigraph of 6 agents

- Vertices represent agents, edges represent interactions.

# Opinion Dynamics on a multigraph




Figure: Multigraph of 6 agents

- Vertices represent agents, edges represent interactions.
- Multigraph composed of two underlying graphs.

# Opinion Dynamics on a multigraph




Figure: Multigraph of 6 agents

- Vertices represent agents, edges represent interactions.
- Multigraph composed of two underlying graphs.
- Blue edges represent virtual interactions (global graph).

# Opinion Dynamics on a multigraph




Figure: Multigraph of 6 agents

- Vertices represent agents, edges represent interactions.
- Multigraph composed of two underlying graphs.
- Blue edges represent virtual interactions (global graph).
- Red edges represent physical interactions (local graph/neighborhood).

# Opinion Dynamics on a multigraph




Figure: Multigraph of 6 agents

- Vertices represent agents, edges represent interactions.
- Multigraph composed of two underlying graphs.
- Blue edges represent virtual interactions (global graph).
- Red edges represent physical interactions (local graph/neighborhood).
- Adversary manipulates mass opinion (Media Houses).



# Opinion Dynamics on a multigraph




Figure: Multigraph of 6 agents

- Vertices represent agents, edges represent interactions.
- Multigraph composed of two underlying graphs.
- Blue edges represent virtual interactions (global graph).
- Red edges represent physical interactions (local graph/neighborhood).
- Adversary manipulates mass opinion (Media Houses).
- Assumption: Local graphs are non-overlapping.

# Opinion Dynamics on a multigraph

---

<sup>1</sup>F. Delarue, "Mean field games: A toy model on an Erdős-Rényi graph." ESAIM: Proceedings and Surveys, vol. 60, pp. 1–26, 2017.

<sup>2</sup>S. Gao, P. E. Caines, and M. Huang, "LQG graphon MFGs," arXiv preprint arXiv:2004.00679, 2020.

<sup>3</sup>H. Tembine, Q. Zhu, and T. Başar, "Risk-sensitive mean-field games," IEEE Transactions on Automatic Control, vol. 59, no. 4, pp. 835–850, 2013.

# Opinion Dynamics on a multigraph

- Can be modeled as a game over multigraphs.

---

<sup>1</sup>F. Delarue, "Mean field games: A toy model on an Erdős-Rényi graph." ESAIM: Proceedings and Surveys, vol. 60, pp. 1–26, 2017.

<sup>2</sup>S. Gao, P. E. Caines, and M. Huang, "LQG graphon MFGs," arXiv preprint arXiv:2004.00679, 2020.

<sup>3</sup>H. Tembine, Q. Zhu, and T. Başar, "Risk-sensitive mean-field games," IEEE Transactions on Automatic Control, vol. 59, no. 4, pp. 835–850, 2013.

# Opinion Dynamics on a multigraph

- Can be modeled as a game over multigraphs.
- Nash Equilibrium (NE) hard to find so consider Mean-Field Games (MFG),  $N \rightarrow \infty$ .

---

<sup>1</sup>F. Delarue, “Mean field games: A toy model on an Erdős–Rényi graph.” ESAIM: Proceedings and Surveys, vol. 60, pp. 1–26, 2017.

<sup>2</sup>S. Gao, P. E. Caines, and M. Huang, “LQG graphon MFGs,” arXiv preprint arXiv:2004.00679, 2020.

<sup>3</sup>H. Tembine, Q. Zhu, and T. Başar, “Risk-sensitive mean-field games,” IEEE Transactions on Automatic Control, vol. 59, no. 4, pp. 835–850, 2013.

# Opinion Dynamics on a multigraph

- Can be modeled as a game over multigraphs.
- Nash Equilibrium (NE) hard to find so consider Mean-Field Games (MFG),  $N \rightarrow \infty$ .
- MFGs over Erdős–Rényi graphs<sup>1</sup> and Graphon MFGs<sup>2</sup> studied in literature.

---

<sup>1</sup>F. Delarue, “Mean field games: A toy model on an Erdős–Rényi graph.” ESAIM: Proceedings and Surveys, vol. 60, pp. 1–26, 2017.

<sup>2</sup>S. Gao, P. E. Caines, and M. Huang, “LQG graphon MFGs,” arXiv preprint arXiv:2004.00679, 2020.

<sup>3</sup>H. Tembine, Q. Zhu, and T. Başar, “Risk-sensitive mean-field games,” IEEE Transactions on Automatic Control, vol. 59, no. 4, pp. 835–850, 2013.

# Opinion Dynamics on a multigraph

- Can be modeled as a game over multigraphs.
- Nash Equilibrium (NE) hard to find so consider Mean-Field Games (MFG),  $N \rightarrow \infty$ .
- MFGs over Erdős–Rényi graphs<sup>1</sup> and Graphon MFGs<sup>2</sup> studied in literature.
- First to investigate the local(physical) vs global(virtual) graph dichotomy in multigraphs.

---

<sup>1</sup>F. Delarue, “Mean field games: A toy model on an Erdős–Rényi graph.” ESAIM: Proceedings and Surveys, vol. 60, pp. 1–26, 2017.

<sup>2</sup>S. Gao, P. E. Caines, and M. Huang, “LQG graphon MFGs,” arXiv preprint arXiv:2004.00679, 2020.

<sup>3</sup>H. Tembine, Q. Zhu, and T. Başar, “Risk-sensitive mean-field games,” IEEE Transactions on Automatic Control, vol. 59, no. 4, pp. 835–850, 2013.

# Opinion Dynamics on a multigraph

- Can be modeled as a game over multigraphs.
- Nash Equilibrium (NE) hard to find so consider Mean-Field Games (MFG),  $N \rightarrow \infty$ .
- MFGs over Erdős–Rényi graphs<sup>1</sup> and Graphon MFGs<sup>2</sup> studied in literature.
- First to investigate the local(physical) vs global(virtual) graph dichotomy in multigraphs.
- Risk-sensitive and robust MFGs<sup>3</sup> considers adversarial agents but does not consider sparse graph structure.

---

<sup>1</sup>F. Delarue, “Mean field games: A toy model on an Erdős–Rényi graph.” ESAIM: Proceedings and Surveys, vol. 60, pp. 1–26, 2017.

<sup>2</sup>S. Gao, P. E. Caines, and M. Huang, “LQG graphon MFGs,” arXiv preprint arXiv:2004.00679, 2020.

<sup>3</sup>H. Tembine, Q. Zhu, and T. Başar, “Risk-sensitive mean-field games,” IEEE Transactions on Automatic Control, vol. 59, no. 4, pp. 835–850, 2013.

# Outline



# Outline

- Formulate the  $N$ -agent Adversarial LQ game over Multigraphs.

# Outline

- Formulate the  $N$ -agent Adversarial LQ game over Multigraphs.
- Introduce the Adversarial LQ-MFG over Multigraphs.

# Outline

- Formulate the  $N$ -agent Adversarial LQ game over Multigraphs.
- Introduce the Adversarial LQ-MFG over Multigraphs.
- Characterize the MFE

# Outline

- Formulate the  $N$ -agent Adversarial LQ game over Multigraphs.
- Introduce the Adversarial LQ-MFG over Multigraphs.
- Characterize the MFE
  - Characterize the equilibrium policies of adversary and generic agent.

# Outline

- Formulate the  $N$ -agent Adversarial LQ game over Multigraphs.
- Introduce the Adversarial LQ-MFG over Multigraphs.
- Characterize the MFE
  - Characterize the equilibrium policies of adversary and generic agent.
  - Characterize the equilibrium aggregate behavior.

# Outline

- Formulate the  $N$ -agent Adversarial LQ game over Multigraphs.
- Introduce the Adversarial LQ-MFG over Multigraphs.
- Characterize the MFE
  - Characterize the equilibrium policies of adversary and generic agent.
  - Characterize the equilibrium aggregate behavior.
- Numerical Results.

# Outline

- Formulate the  $N$ -agent Adversarial LQ game over Multigraphs.
- Introduce the Adversarial LQ-MFG over Multigraphs.
- Characterize the MFE
  - Characterize the equilibrium policies of adversary and generic agent.
  - Characterize the equilibrium aggregate behavior.
- Numerical Results.
- Future directions.

## Formulation (finite population)

- Agent  $i \in [N]$  has linear dynamics (uncoupled),

$$Z_{t+1}^i = AZ_t^i + BU_t^i + CV_t + W_t^i, t \in (0, \dots, T-1),$$



# Formulation (finite population)

- Agent  $i \in [N]$  has linear dynamics (uncoupled),

$$Z_{t+1}^i = AZ_t^i + BU_t^i + CV_t + W_t^i, t \in (0, \dots, T-1),$$

- Adversary input ( $V_t$ ) affects dynamics and cost of agents,

# Formulation (finite population)

- Agent  $i \in [N]$  has linear dynamics (uncoupled),

$$Z_{t+1}^i = AZ_t^i + BU_t^i + CV_t + W_t^i, t \in (0, \dots, T-1),$$

- Adversary input ( $V_t$ ) affects dynamics and cost of agents,
- $Y_t^i$  is local effect on agent  $i$ ,  $Y_t^i = \sum_{j \in \mathcal{N}(i)} Z_t^j / |\mathcal{N}(i)|$ .

# Formulation (finite population)

- Agent  $i \in [N]$  has linear dynamics (uncoupled),

$$Z_{t+1}^i = AZ_t^i + BU_t^i + CV_t + W_t^i, t \in (0, \dots, T-1),$$

- Adversary input ( $V_t$ ) affects dynamics and cost of agents,
- $Y_t^i$  is local effect on agent  $i$ ,  $Y_t^i = \sum_{j \in \mathcal{N}(i)} Z_t^j / |\mathcal{N}(i)|$ .
- Agent  $i$  coupled through *consensus-like* terms in cost,

$$J_N^i(\pi^i, \pi^{-i}, V) =$$

$$\begin{aligned} & \sum_{t=0}^{T-1} \mathbb{E} \left[ \|Z_t^i\|_{Q_t}^2 + \left\| Z_t^i - \frac{1}{N} \sum_{j=1}^N Z_t^j \right\|_{\bar{Q}_t}^2 + \|Z_t^i - Y_t^i\|_{\bar{Q}_t}^2 + \|U_t^i\|_{R_t}^2 - \|V_t\|_{S_t}^2 \right] \\ & + \mathbb{E} \left[ \|Z_T^i\|_{Q_T}^2 + \left\| Z_T^i - \frac{1}{N} \sum_{j=1}^N Z_T^j \right\|_{\bar{Q}_T}^2 + \|Z_T^i - Y_T^i\|_{\bar{Q}_T}^2 \right]. \end{aligned}$$

# Formulation (finite population)

- Agent  $i \in [N]$  has linear dynamics (uncoupled),

$$Z_{t+1}^i = AZ_t^i + BU_t^i + CV_t + W_t^i, t \in (0, \dots, T-1),$$

- Adversary input ( $V_t$ ) affects dynamics and cost of agents,
- $Y_t^i$  is local effect on agent  $i$ ,  $Y_t^i = \sum_{j \in \mathcal{N}(i)} Z_t^j / |\mathcal{N}(i)|$ .
- Agent  $i$  coupled through *consensus-like* terms in cost,

$$J_N^i(\pi^i, \pi^{-i}, V) =$$

$$\sum_{t=0}^{T-1} \mathbb{E} \left[ \|\mathbf{Z}_t^i\|_{Q_t}^2 + \left\| Z_t^i - \frac{1}{N} \sum_{j=1}^N Z_t^j \right\|_{\bar{Q}_t}^2 + \|Z_t^i - Y_t^i\|_{\bar{Q}_t}^2 + \|U_t^i\|_{R_t}^2 - \|V_t\|_{S_t}^2 \right] \\ + \mathbb{E} \left[ \|\mathbf{Z}_T^i\|_{Q_T}^2 + \left\| Z_T^i - \frac{1}{N} \sum_{j=1}^N Z_T^j \right\|_{\bar{Q}_T}^2 + \|Z_T^i - Y_T^i\|_{\bar{Q}_T}^2 \right].$$

# Formulation (finite population)

- Agent  $i \in [N]$  has linear dynamics (uncoupled),

$$Z_{t+1}^i = AZ_t^i + BU_t^i + CV_t + W_t^i, t \in (0, \dots, T-1),$$

- Adversary input ( $V_t$ ) affects dynamics and cost of agents,
- $Y_t^i$  is local effect on agent  $i$ ,  $Y_t^i = \sum_{j \in \mathcal{N}(i)} Z_t^j / |\mathcal{N}(i)|$ .
- Agent  $i$  coupled through *consensus-like* terms in cost,

$$J_N^i(\pi^i, \pi^{-i}, V) =$$

$$\begin{aligned} & \sum_{t=0}^{T-1} \mathbb{E} \left[ \|Z_t^i\|_{Q_t}^2 + \left\| Z_t^i - \frac{1}{N} \sum_{j=1}^N Z_t^j \right\|_{\bar{Q}_t}^2 + \|Z_t^i - Y_t^i\|_{\bar{Q}_t}^2 + \|U_t^i\|_{R_t}^2 - \|V_t\|_{S_t}^2 \right] \\ & + \mathbb{E} \left[ \|Z_T^i\|_{Q_T}^2 + \left\| Z_T^i - \frac{1}{N} \sum_{j=1}^N Z_T^j \right\|_{\bar{Q}_T}^2 + \|Z_T^i - Y_T^i\|_{\bar{Q}_T}^2 \right]. \end{aligned}$$

# Formulation (finite population)

- Agent  $i \in [N]$  has linear dynamics (uncoupled),

$$Z_{t+1}^i = AZ_t^i + BU_t^i + CV_t + W_t^i, t \in (0, \dots, T-1),$$

- Adversary input ( $V_t$ ) affects dynamics and cost of agents,
- $Y_t^i$  is local effect on agent  $i$ ,  $Y_t^i = \sum_{j \in \mathcal{N}(i)} Z_t^j / |\mathcal{N}(i)|$ .
- Agent  $i$  coupled through *consensus-like* terms in cost,

$$J_N^i(\pi^i, \pi^{-i}, V) =$$

$$\begin{aligned} & \sum_{t=0}^{T-1} \mathbb{E} \left[ \|Z_t^i\|_{Q_t}^2 + \left\| Z_t^i - \frac{1}{N} \sum_{j=1}^N Z_t^j \right\|_{\bar{Q}_t}^2 + \|Z_t^i - Y_t^i\|_{\bar{Q}_t}^2 + \|U_t^i\|_{R_t}^2 - \|V_t\|_{S_t}^2 \right] \\ & + \mathbb{E} \left[ \|Z_T^i\|_{Q_T}^2 + \left\| Z_T^i - \frac{1}{N} \sum_{j=1}^N Z_T^j \right\|_{\bar{Q}_T}^2 + \|Z_T^i - Y_T^i\|_{\bar{Q}_T}^2 \right]. \end{aligned}$$

# Formulation (finite population)

- Agent  $i \in [N]$  has linear dynamics (uncoupled),

$$Z_{t+1}^i = AZ_t^i + BU_t^i + CV_t + W_t^i, t \in (0, \dots, T-1),$$

- Adversary input ( $V_t$ ) affects dynamics and cost of agents,
- $Y_t^i$  is local effect on agent  $i$ ,  $Y_t^i = \sum_{j \in \mathcal{N}(i)} Z_t^j / |\mathcal{N}(i)|$ .
- Agent  $i$  coupled through *consensus-like* terms in cost,

$$J_N^i(\pi^i, \pi^{-i}, V) =$$

$$\sum_{t=0}^{T-1} \mathbb{E} \left[ \|Z_t^i\|_{Q_t}^2 + \left\| Z_t^i - \frac{1}{N} \sum_{j=1}^N Z_t^j \right\|_{\bar{Q}_t}^2 + \|Z_t^i - Y_t^i\|_{\bar{Q}_t}^2 + \|U_t^i\|_{R_t}^2 - \|V_t\|_{S_t}^2 \right] \\ + \mathbb{E} \left[ \|Z_T^i\|_{Q_T}^2 + \left\| Z_T^i - \frac{1}{N} \sum_{j=1}^N Z_T^j \right\|_{\bar{Q}_T}^2 + \|Z_T^i - Y_T^i\|_{\bar{Q}_T}^2 \right].$$

# Formulation (finite population)

- Agent  $i \in [N]$  has linear dynamics (uncoupled),

$$Z_{t+1}^i = AZ_t^i + BU_t^i + CV_t + W_t^i, t \in (0, \dots, T-1),$$

- Adversary input ( $V_t$ ) affects dynamics and cost of agents,
- $Y_t^i$  is local effect on agent  $i$ ,  $Y_t^i = \sum_{j \in \mathcal{N}(i)} Z_t^j / |\mathcal{N}(i)|$ .
- Agent  $i$  coupled through *consensus-like* terms in cost,

$$J_N^i(\pi^i, \pi^{-i}, V) =$$

$$\sum_{t=0}^{T-1} \mathbb{E} \left[ \|Z_t^i\|_{Q_t}^2 + \left\| Z_t^i - \frac{1}{N} \sum_{j=1}^N Z_t^j \right\|_{\bar{Q}_t}^2 + \|Z_t^i - Y_t^i\|_{\bar{Q}_t}^2 + \|U_t^i\|_{R_t}^2 - \|V_t\|_{S_t}^2 \right] \\ + \mathbb{E} \left[ \|Z_T^i\|_{Q_T}^2 + \left\| Z_T^i - \frac{1}{N} \sum_{j=1}^N Z_T^j \right\|_{\bar{Q}_T}^2 + \|Z_T^i - Y_T^i\|_{\bar{Q}_T}^2 \right].$$



# Formulation (finite population)

## Formulation (finite population)

- $|\mathcal{N}(i)| = |\mathcal{N}(j)|, i, j \in [N]$

## Formulation (finite population)

- $|\mathcal{N}(i)| = |\mathcal{N}(j)|, i, j \in [N]$
- Agent  $i$  has access to local history, NE hard to find.

## Formulation (finite population)

- $|\mathcal{N}(i)| = |\mathcal{N}(j)|, i, j \in [N]$
- Agent  $i$  has access to local history, NE hard to find.
- Cost function of adversary,

$$J_N^0(V, \pi^{(N)}) = -\frac{1}{N} \sum_{i=1}^N J_N^i(\pi^i, \pi^{-i}, V)$$

## Formulation (finite population)

- $|\mathcal{N}(i)| = |\mathcal{N}(j)|, i, j \in [N]$
- Agent  $i$  has access to local history, NE hard to find.
- Cost function of adversary,

$$J_N^0(V, \pi^{(N)}) = -\frac{1}{N} \sum_{i=1}^N J_N^i(\pi^i, \pi^{-i}, V)$$

- Adversary aims to disrupt/delay consensus.

# Formulation (finite population)

- $|\mathcal{N}(i)| = |\mathcal{N}(j)|, i, j \in [N]$
- Agent  $i$  has access to local history, NE hard to find.
- Cost function of adversary,

$$J_N^0(V, \pi^{(N)}) = -\frac{1}{N} \sum_{i=1}^N J_N^i(\pi^i, \pi^{-i}, V)$$

- Adversary aims to disrupt/delay consensus.

## Definition (Nash Equilibrium)

The set  $\pi^* = (\pi^{1*}, \dots, \pi^{N*})$  and adversary policy  $V^* = (V_0^*, \dots, V_{T-1}^*)$  constitute Nash equilibrium if,

$$J_N^i(\pi^{i*}, \pi^{-i*}, V^*) \leq J_N^i(\pi^i, \pi^{-i*}, V^*), \quad \pi^i \in \Pi^i, i \in [N]$$

$$J_N^0(\pi^*, V^*) \leq J_N^0(\pi^*, V), \quad V \in \mathcal{V}$$

# Formulation (mean-field game)

## Formulation (mean-field game)

- $N \rightarrow \infty$  but finite local graph.



## Formulation (mean-field game)

- $N \rightarrow \infty$  but finite local graph.
- Generic agent dynamics

$$Z_{t+1} = AZ_t + BU_t + CV_t + W_t.$$

# Formulation (mean-field game)

- $N \rightarrow \infty$  but finite local graph.
- Generic agent dynamics

$$Z_{t+1} = AZ_t + BU_t + CV_t + W_t.$$

- Generic agent cost function

$$\begin{aligned} J(\mu, \bar{Z}, V) = & \sum_{t=0}^{T-1} \mathbb{E} [\|Z_t\|_{Q_t}^2 + \|Z_t - Y_t\|_{\tilde{Q}_t}^2 + \|Z_t - \bar{Z}_t\|_{\tilde{Q}_t}^2 + \|U_t\|_{R_t}^2 - \|V_t\|_{S_t}^2] \\ & + \mathbb{E} [\|Z_T\|_{Q_T}^2 + \|Z_T - Y_T\|_{\tilde{Q}_T}^2 + \|Z_T - \bar{Z}_T\|_{\tilde{Q}_T}^2] \end{aligned}$$

# Formulation (mean-field game)

- $N \rightarrow \infty$  but finite local graph.
- Generic agent dynamics

$$Z_{t+1} = AZ_t + BU_t + CV_t + W_t.$$

- Generic agent cost function

$$\begin{aligned} J(\mu, \bar{Z}, V) = & \sum_{t=0}^{T-1} \mathbb{E} [\|Z_t\|_{Q_t}^2 + \|Z_t - Y_t\|_{\hat{Q}_t}^2 + \|Z_t - \bar{Z}_t\|_{\bar{Q}_t}^2 + \|U_t\|_{R_t}^2 - \|V_t\|_{S_t}^2] \\ & + \mathbb{E} [\|Z_T\|_{Q_T}^2 + \|Z_T - Y_T\|_{\hat{Q}_T}^2 + \|Z_T - \bar{Z}_T\|_{\bar{Q}_T}^2] \end{aligned}$$

- $\bar{Z} = (\bar{Z}_0, \dots, \bar{Z}_T)$  represents global mean-field,  
 $Y = (Y_0, \dots, Y_T)$  represents local mean-field.

# Formulation (mean-field game)

---

<sup>4</sup>P. Turnes Jr and L. Monteiro, "An epidemic model to evaluate the homogeneous mixing assumption," Communications in Nonlinear Science and Numerical Simulation, vol. 19, no. 11, pp. 4042–4047, 2014.

<sup>5</sup>B. O. Baumgaertner, P. A. Fetros, S. M. Krone, and R. C. Tyson, "Spatial opinion dynamics and the effects of two types of mixing," Physical Review E, vol. 98, no. 2, p. 022310, 2018

# Formulation (mean-field game)

## Assumption (Homogeneous Mixing)

*The local mean-field  $Y_t$  is assumed to be an exogenous noise process with mean  $\mathbb{E}[Y_t] = \bar{Z}_t$ .*

---

<sup>4</sup>P. Turnes Jr and L. Monteiro, "An epidemic model to evaluate the homogeneous mixing assumption," Communications in Nonlinear Science and Numerical Simulation, vol. 19, no. 11, pp. 4042–4047, 2014.

<sup>5</sup>B. O. Baumgaertner, P. A. Fetros, S. M. Krone, and R. C. Tyson, "Spatial opinion dynamics and the effects of two types of mixing," Physical Review E, vol. 98, no. 2, p. 022310, 2018

# Formulation (mean-field game)

## Assumption (Homogeneous Mixing)

*The local mean-field  $Y_t$  is assumed to be an exogenous noise process with mean  $\mathbb{E}[Y_t] = \bar{Z}_t$ .*

- Hypothesis standard in infection<sup>4</sup> and opinion dynamics<sup>5</sup> literature.

---

<sup>4</sup>P. Turnes Jr and L. Monteiro, "An epidemic model to evaluate the homogeneous mixing assumption," Communications in Nonlinear Science and Numerical Simulation, vol. 19, no. 11, pp. 4042–4047, 2014.

<sup>5</sup>B. O. Baumgaertner, P. A. Fetros, S. M. Krone, and R. C. Tyson, "Spatial opinion dynamics and the effects of two types of mixing," Physical Review E, vol. 98, no. 2, p. 022310, 2018

# Formulation (mean-field game)

## Assumption (Homogeneous Mixing)

*The local mean-field  $Y_t$  is assumed to be an exogenous noise process with mean  $\mathbb{E}[Y_t] = \bar{Z}_t$ .*

- Hypothesis standard in infection<sup>4</sup> and opinion dynamics<sup>5</sup> literature.
- Adversary's cost function,  $J^0(V, \mu, \bar{Z}) = \lim_{N \rightarrow \infty} J_N^0(V, \mu^{(N)})$ .

---

<sup>4</sup>P. Turnes Jr and L. Monteiro, "An epidemic model to evaluate the homogeneous mixing assumption," Communications in Nonlinear Science and Numerical Simulation, vol. 19, no. 11, pp. 4042–4047, 2014.

<sup>5</sup>B. O. Baumgaertner, P. A. Fetros, S. M. Krone, and R. C. Tyson, "Spatial opinion dynamics and the effects of two types of mixing," Physical Review E, vol. 98, no. 2, p. 022310, 2018

# Formulation (mean-field game)

## Assumption (Homogeneous Mixing)

*The local mean-field  $Y_t$  is assumed to be an exogenous noise process with mean  $\mathbb{E}[Y_t] = \bar{Z}_t$ .*

- Hypothesis standard in infection<sup>4</sup> and opinion dynamics<sup>5</sup> literature.
- Adversary's cost function,  $J^0(V, \mu, \bar{Z}) = \lim_{N \rightarrow \infty} J_N^0(V, \mu^{(N)})$ .

## Definition (Mean-Field Equilibrium)

The tuple  $(\mu^*, V^*, \bar{Z}^*, Y^*)$  is a mean-field equilibrium if,

**Optimality:**  $\mu^* = \operatorname{argmin}_{\mu} J(\mu, \bar{Z}^*, V^*)$ ,  $V^* = \operatorname{argmin}_V J^0(V, \mu^*, \bar{Z}^*)$ ,

**Consistency:**  $\bar{Z}^*$  and  $Y^*$  are aggregate behaviors *consistent* with  $\mu^*$  and  $V^*$ .

---

<sup>4</sup>P. Turnes Jr and L. Monteiro, "An epidemic model to evaluate the homogeneous mixing assumption," Communications in Nonlinear Science and Numerical Simulation, vol. 19, no. 11, pp. 4042–4047, 2014.

<sup>5</sup>B. O. Baumgaertner, P. A. Fetros, S. M. Krone, and R. C. Tyson, "Spatial opinion dynamics and the effects of two types of mixing," Physical Review E, vol. 98, no. 2, p. 022310, 2018



# Form of generic agent's equilibrium control

# Form of generic agent's equilibrium control

## Theorem

*The generic agent's equilibrium control adapted to filtration  $\mathcal{F}_t^Z \vee \mathcal{F}_t^Y$  is given by*

$$U_t^* = -R_t^{-1} B^T \zeta_{t+1}$$

*where  $\zeta_t$  can be constructed as*

$$\zeta_t = A^T \zeta_{t+1} + (Q_t + \bar{Q}_t + \tilde{Q}_t) Z_t - \bar{Q}_t \bar{Z}_t - \tilde{Q}_t Y_t - M_t^\zeta,$$

$$\zeta_T = (Q_T + \bar{Q}_T + \tilde{Q}_T) Z_T - \bar{Q}_T \bar{Z}_T - \tilde{Q}_T Y_T,$$

$$M_t^\zeta = A^T \zeta_{t+1} - A^T \mathbb{E}[\zeta_{t+1} \mid \mathcal{F}_t^Z \vee \mathcal{F}_t^Y].$$

*where  $M_t^\zeta$  is a martingale difference sequence adapted to filtration  $\mathcal{F}_t^Z \vee \mathcal{F}_t^Y$ .*

# Form of adversary's equilibrium control

# Form of adversary's equilibrium control

## Theorem

*If the following condition is satisfied*

$$S_t - C^T \hat{P}_{t+1} C > 0, \quad (1)$$

*where the matrix  $\hat{P}_t$  is defined recursively by*

$$\hat{P}_t = Q_t + A^T \hat{P}_{t+1} A + A^T \hat{P}_{t+1} C (S_t - C^T \hat{P}_{t+1} C)^{-1} C^T \hat{P}_{t+1} A, \quad \hat{P}_T = Q_T,$$

*then the equilibrium control policy of the adversary is,*

$$V_t^* = -S_t^{-1} C^T \bar{\zeta}_{t+1}^0, \bar{\zeta}_t^0 = A^T \bar{\zeta}_{t+1}^0 - Q_t \bar{Z}_t, \bar{\zeta}_T^0 = -Q_T \bar{Z}_T, \quad (2)$$

*where  $\bar{\zeta}_t^0$  is the adversary's co-state and  $\bar{Z}$  is the global MF of the agents.*

# Equilibrium Global MF dynamics

# Equilibrium Global MF dynamics

## Theorem

*If the preceding assumptions are satisfied, then equilibrium global MF follows linear dynamics,*

$$\bar{Z}_{t+1}^* = E_t^{-1} A \bar{Z}_t^* = \bar{F}_t \bar{Z}_t^*, \quad (3)$$

*where  $E_t = (I + BR_t^{-1}B^T\bar{P}_{t+1} - CS_t^{-1}C^T\bar{P}_{t+1})$  and  $\bar{P}_t$  is given by the Riccati equation,*

$$\bar{P}_t = A^T \bar{P}_{t+1} E_t^{-1} A + Q_t, P_T = Q_T$$

*and the equilibrium adversarial policy,*

$$V_t^* = S_t^{-1} C^T \bar{P}_t \bar{F}_t \bar{Z}_t^*. \quad (4)$$

*is linear in the equilibrium global MF.*

# Equilibrium Local MF dynamics

# Equilibrium Local MF dynamics

## Theorem

*If the preceding assumptions are satisfied, then local equilibrium MF has linear Gaussian dynamics driven by the equilibrium global MF:*

$$Y_{t+1}^* = \tilde{F}_t^1 Y_t^* + \tilde{F}_t^2 \bar{Z}_t^* + \tilde{E}_t^{-1} \tilde{W}_t \quad (5)$$

where  $\tilde{W}_t = \sum_{j \in \mathcal{N}} W_t^j / |\mathcal{N}|$ ,

$$\tilde{E}_t = (I + BR^{-1}B^T \tilde{P}_{t+1} - CS_t^{-1}C^T \tilde{P}_{t+1}), \quad \tilde{F}_t^1 = \tilde{E}_t^{-1}A,$$

$$\tilde{F}_t^2 = \tilde{E}_t^{-1}(BR_t^{-1}B^T - CS_t^{-1}C^T) \sum_{i=0}^{T-t} \prod_{j=1}^i \tilde{H}_{t+j} \bar{Q}_{t+i} \bar{F}_{t+j}$$

$$\tilde{P}_t = A^T \tilde{P}_{t+1} \tilde{E}_t^{-1} A + Q_t + \bar{Q}_t, \quad \tilde{P}_T = Q_T + \bar{Q}_T,$$

and  $\tilde{H}_t = A^T(I - \tilde{P}_{t+1} \tilde{E}_t^{-1} BR^{-1}B^T)$ .



# MFE Characterization

- Generic agent's problem can be cast as time varying linear quadratic regulator (TV-LQR) problem.

$$U_t^* = \mu_t^*(Z_t, Y_t^*, \bar{Z}_t^*) = \bar{K}_t^* \begin{pmatrix} Z_t \\ Y_t^* \\ \bar{Z}_t^* \end{pmatrix}$$

# MFE Characterization

**Conclusion:**

# MFE Characterization

## Conclusion:

- Then global mean-field (at equilibrium) has deterministic dynamics, so can be computed offline.

# MFE Characterization

## Conclusion:

- Then global mean-field (at equilibrium) has deterministic dynamics, so can be computed offline.
- The adversary actions (at equilibrium) depend *only* on the global mean-field.


# MFE Characterization

## Conclusion:

- Then global mean-field (at equilibrium) has deterministic dynamics, so can be computed offline.
- The adversary actions (at equilibrium) depend *only* on the global mean-field.
- The agent's actions (at equilibrium) have linear dependence on,
  - the agent's state,
  - the local mean-field,
  - the global mean-field.

# Numerical Results (Affect of Adversary)

- Number of agents  $N = 10,000$
- Neighborhood size  $|\mathcal{N}(i)| = 1000$ .



# Future Directions

- $\epsilon$ -Nash analysis of the MFE.
- More general local network structure given *homogenous mixing hypothesis*,
  - Overlapping local graphs,
  - Agents can have types specifying local connectivity.
- Multiple adversaries.