

AI Crop Diagnostics Report

Detailed disease prediction report with AI insights, recommended treatment, and probability analytics.

Report Generated:	2026-05-28 09:41:20 UTC
Farmer / User:	sandeep
Crop:	Corn
Disease Detected:	Gray_Leaf_Spot
Prediction Time:	2026-05-28 09:40:56




Prediction Overview

Confidence Score	83.86%
Severity	Moderate
Fertilizer Recommendation	Use a balanced fertilizer for corn with moderate nitrogen and adequate potassium to support recovery.
Dosage	Apply a light corrective dose according to soil test results or label guidance. Avoid over-fertilizing.
Application Timing	Apply after removing infected tissue and when foliage is dry, preferably early morning or late afternoon.
Spraying Interval	Repeat application every 7-10 days during recovery, or as directed by product label.
Organic Alternative	Use well-decomposed compost, vermicompost, seaweed extract, or fermented plant-based bio-fertilizers.

Prevention Tips	Maintain spacing, rotate crops, remove debris, keep tools clean, and monitor new symptoms closely.
-----------------	--

Fertilizer Guidance

Fertilizer Name	Balanced NPK 10-10-10 fertilizer for Corn
NPK Values	10-10-10
Dosage	Apply a light corrective dose according to soil test results or label guidance. Avoid over-fertilizing.
Application Timing	Apply after removing infected tissue and when foliage is dry, preferably early morning or late afternoon.
Spraying Interval	Repeat application every 7-10 days during recovery, or as directed by product label.
Organic Alternative	Use well-decomposed compost, vermicompost, seaweed extract, or fermented plant-based bio-fertilizers.



AI Remedy Summary

Disease Summary
Gray_Leaf_Spot has been detected on Corn with 83.86% confidence. This recommendation is designed to help slow spread, support recovery, and protect nearby plants.

Causes
Likely contributors include excess humidity, leaf wetness, poor airflow, infected planting material, or delayed field sanitation.

Treatment
Remove severely affected leaves, improve canopy airflow, and follow approved crop-protection practices recommended by local agricultural officers.

Fertilizer Name
Balanced NPK 10-10-10 fertilizer for Corn

NPK Values

10-10-10

Fertilizer Recommendation

Use a balanced fertilizer for corn with moderate nitrogen and adequate potassium to support recovery without forcing weak new growth.

Dosage

Apply a light corrective dose according to soil test results or label guidance. Avoid over-fertilizing stressed plants.

Application Timing

Apply after removing infected tissue and when foliage is dry, preferably early morning or late afternoon.

Spraying Interval

Repeat application every 7-10 days during recovery, or as directed by product label.

Organic Alternatives

Use well-decomposed compost, vermicompost, seaweed extract, or fermented plant-based bio-stimulants as safer support options.

Prevention Tips

Maintain spacing, rotate crops, remove debris, keep tools clean, and monitor new symptoms closely for 7 days.

7-Day Recovery Plan

Day 1-2: remove infected tissue and sanitize tools. Day 3-4: apply the first support treatment. Day 5-7: monitor new growth and repeat only if symptoms continue to spread.

Recovery & Prevention

Day 1-2: remove infected tissue and sanitize tools. Day 3-4: apply the first support treatment. Day 5-7: monitor new growth and repeat only if symptoms continue to spread.

AI Diagnostics Lab | AgriDetect™
Certified AI Crop Health Reporting

This document is generated by AgriDetect AI for agricultural diagnosis and advisory. Refer to local agricultural experts for further treatment.