

University
of Basel

**Fully quantum description of a three level
maser, driven by a thermal bath**

Master-project

Sander Stammbach
Prof. Patrick Potts
Dr. Mateo Brunelli
Marcelo Janovitch

12 November 2022

Contents

1	Introduction	2
2	System and model	3
2.1	3-level-Maser model	3
2.2	Master-Equation	4
3	Methods	5
3.1	Implementation of the three-level-system in qutip	5
4	Lasing transition	6
4.1	Wigner function and phase-averaged coherent states	6
4.2	Double threshold behaviour and population inversion	8
5	Thermodynamics	12
5.1	Heat currents	12
5.2	Entropy production	14
6	Conclusion and outlook	16
7	Appendix	18
7.1	Software	18

1 Introduction

One of the most important questions in thermodynamics is how to convert thermal energy into work. For such tasks there exist many classical engines, for example the steam machines or gasoline engines. Those ideas also generalize to quantum systems. In this master-project, a three-level maser, driven by the coupling of a hot and a cold bath is quantified. The three-level maser is a quantum heat engine (QHE). The work extraction from a classical heat engine is often a moving piston. But in this case it is a driving field. In the year 1916 Albert Einstein already discussed three ways of light-matter-interaction (spontaneous emission, absorption, and stimulated emission)[2]. In the paper from Scovil and Schulz-DuBois 1959 [5] investigated whether a laser is a heat engine. In this paper, they took a maser as a device to transform heat into coherent radiation, because heat can make a population inversion. In their thermodynamic analysis, they used a single-atom laser. They made a groundwork for emerging theory of quantum thermodynamics. In practice and also for the calculations, two different reservoirs are necessary. The high-temperature reservoir can be realized by a fast and accurate estimation of the thermal occupation of propagating microwave modes [4].

2 System and model

2.1 3-level-Maser model




Figure 1: Schematic representation of a three-level maser heat engine continuously coupled to two reservoirs of temperatures T_h ; and T_c . The three energy levels are described by E_1, E_2, E_3 . The system is interacting with a quantized single mode field. λ represents the strength of matter-field coupling.

A maser/laser consists of two elements. One of them is a gain medium and the other one is an optical resonator. A gain medium is always a material with an atomic transition between two atomic states. When an atom decays from an energetically higher state to an energetically lower state, a photon is created. In a three level system the three energy levels are E_1, E_2, E_3 shown in Fig. 1. In a first part the system gets pumped from the lowest E_1 level to the highest level E_3 . The resonator should have a longer decay time, so that a population inversion can build up. This means that the system is in the energetically higher state. From this state they come almost exclusively

through stimulated emission into the lower state E_1 . The cavity is build of two mirrors. One of the cavity has a small leaking, so that a small part of the photons can leave the cavity. This leaking is quantified by a constant κ . Stimulated emission is a necessary condition for coherent light. Coherent light means all the photons have the same phase and same frequency [2]. Here we consider that higher level will be reached with a interaction of a hot bath. We denote the transition frequencies of $\omega_h = (E_3 - E_1)/\hbar$, $\omega_c = (E_2 - E_1)/\hbar$ and $\omega_{cav} = (E_2 - E_1)/\hbar$. Our cavity is in resonance, therefore we set $\omega_{cav} = \omega_f$. In this three-level system each thermal photon from the bath trigger a lasing photon. J_h iast the heat flow from the hot bath, and J_{cav} is the heat flow in the cavity. The efficiency of the three-level system is then given by the following formula:

$$\frac{J_{cav}}{J_h} = \eta_{maser} = \frac{\omega_{cav}}{\omega_h} < 1 - \frac{T_c}{T_h}. \quad (1)$$

2.2 Master-Equation

An arbitrary state of a cavity and a atom can be described by a total density operator $\hat{\rho}(t)$. The complete description of the total system's state at a given time t is encoded in this density operator $\hat{\rho}$ [2]. The master equation is:

$$\dot{\rho}(t) = \frac{1}{i\hbar}[H, \rho] + \mathcal{L}_h\rho + \mathcal{L}_c\rho + \mathcal{L}_{cav}\rho. \quad (2)$$

To derive the $\hat{\rho}$ we can solve a specific differential equation. This equation is called Lindblad-master-equation (2). The first part of eq (2) is the von Neuman equation, the analogue of the Schrödinger equation for density matrices. This part of the equation is unitary and therefore the process is reversible. H is the the Hamiltonian operator. The Hamiltonian describes the energy. The atomic field system is composed of two crucial parts; the atomic states, the cavity field, and the interaction between the cavity and the state of the system. The total Hamiltonian,

$$H = H_{free} + H_{int}, \quad (3)$$

consist of two parts; The part of the free Hamiltonian, where i describe the different states,

$$H_{free} = \sum_{i=1}^3 \hbar\omega_i |i\rangle \langle i| + \hbar\omega_{cav} a^\dagger a, \quad (4)$$

and the interaction Hamiltonian or Jaynes-Cummings Hamiltonian:

$$H_{int} = \hbar g(\sigma_{12}a^\dagger + \sigma_{21}a). \quad (5)$$

g is the coupling constant between the system and the photons.

The interaction with the various environmental heat baths is described by the non-unitary Liouvillian. \mathcal{L}_h describe the interaction with the hot bath, \mathcal{L}_c is the contribution from the interaction with the cold bath coupled with the atom. \mathcal{L}_{cav} describe the interaction with the photons which are in the cavity. Those $\mathcal{L}_{h,c,cav}$ are also called superoperators, which act on the density operator. A superoperator is a linear operator acting on a vector space of linear operator. The interaction with the various environmental heat baths is described by the Liouvillian:

$$\begin{aligned}\mathcal{L}_h\hat{\rho} &= \frac{\gamma_h}{2}(n(\omega_h, T_h) + 1) \cdot \mathcal{D}[\sigma_{13}]\rho + \frac{\gamma_h}{2}n(\omega_h, T_h) \cdot \mathcal{D}[\sigma_{31}]\rho \\ \mathcal{L}_c\hat{\rho} &= \frac{\gamma_c}{2}(n(\omega_c, T_c) + 1) \cdot \mathcal{D}[\sigma_{23}]\rho + \frac{\gamma_c}{2}n(\omega_c, T_c) \cdot \mathcal{D}[\sigma_{32}]\rho \\ \mathcal{L}_{cav}\hat{\rho} &= \kappa((\omega_{cav}, T_f) + 1) \cdot \mathcal{D}[a]\rho + \kappa n(\omega_{cav}, T_f) \cdot \mathcal{D}[a^\dagger]\rho.\end{aligned}\tag{6}$$

σ_{12} is the transition operator and defined as $|1\rangle\langle 2|$. Similar for $\sigma_{13} = |1\rangle\langle 3|$, $\sigma_{23} = |2\rangle\langle 3|$. The transition operator describe the transition between one atomic state to a other. \mathcal{D} is defined with following formula

$$\mathcal{D}[A]\rho = (2A\rho A^\dagger - A^\dagger A\rho - \rho A^\dagger A).\tag{7}$$

Where A represents one of the transition operators σ_{ab} and a ladder operator. The Bose-Einstein occupation number is the mean number of excitations in the reservoir damping the oscillator. It describes the mean occupation number $\langle n(E) \rangle$ of a quantum state of energy E , in thermodynamic equilibrium at absolute temperature T for identical bosons as occupying particles. n depends on the temperature and the frequency. n is defined as: $n_i(\omega_i, T) = \frac{1}{\exp[\frac{\hbar\omega_i}{k_b T_i}] - 1}$.

κ describes the rate of photons which leave the cavity. The prefactor γ_c, γ_h describes the spontaneous decay rates and are set to one in this calculation. In this case the master equation is solved for the steady states. A steady state is a state or condition of a system or process, that the density matrix of the state does not change in time, or the changes are negligibly. Therefore all observables do not change in time either. It contains the whole description of the three-level system and the cavity.

3 Methods

3.1 Implementation of the three-level-system in qutip

The master equation is solved numerically. For that it is necessary to define the constants first. In our case, only the transition between $|1\rangle$ and $|2\rangle$

interact with the light. Defined as constants are the three different Bose-Einstein occupation n_h , n_c and n_f . The transition-operators σ_{ab} are made by qutip implementation of the tensor product of $|a\rangle\langle b|$ and the identity matrices of size of maximum photon number. $|1\rangle$ and $|2\rangle$ are basisstates. For our purpose, it will always suffice to truncate the photon's Hilbertspace of $n = 30$. So we get 90 x 90 matrices. The projectors are implemented similarly, with the matrix $|a\rangle\langle a|$. With those it is easy to construct the Hamiltonions, H_{free} and H_{int} , as in Eq. (4), Eq. (5). To calculate the density matrices for steady states we can also use a qutip function. This function needs the total Hamiltonian and a list of the non-unitary operators as arguments. As output of the function steady-state we get the density-matrices for steady-states. [3]

4 Lasing transition

4.1 Wigner function and phase-averaged coherent states

The output of a laser is coherent light. The quantum description of coherent light is a coherent state. The photon number distribution of coherent light is a Poisson distribution. In our case the output is a phase-averaged coherent state (PHAV). That's a coherent state which have a averaged phase ϕ around 2π . Randomizing the phase of a coherent state doesn't change the photon-number distribution.

A Wigner function is a representation of a general quantum state of light [1]. The function describes the probability density in phase space.

$$w(x, p) = \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} d\xi e^{\frac{-ip\xi}{\hbar}} \langle x + \frac{1}{2}\xi | \rho | x - \frac{1}{2}\xi \rangle. \quad (8)$$

The Wigner function of a coherent and a phase-average-state is non-Gaussian, because p_{nn} of Eq. (10) is a Poisson distribution. So we get a new term of $\exp(i\pi\phi)$ in it. The standard coherent state $|\alpha\rangle$ can be represented by following formula:

$$|\alpha\rangle = e^{\frac{-|\alpha|^2}{2}} \sum_{n=0}^{\infty} \frac{|\alpha|^n e^{in\phi}}{\sqrt{n!}} |n\rangle. \quad (9)$$

Get the PHAV state will be obtained by integrating over the phase ϕ .

$$\rho_{PHAV} = \int_0^{2\pi} \frac{d\phi}{2\pi} |\alpha(\phi)\rangle \langle \alpha(\phi)| = \sum_{n=0}^{\infty} p_{nn} |n\rangle \langle n|, \quad (10)$$

thus is equal to

$$\rho_{PHAV} = \sum_{n=0}^{\infty} \exp(-|\alpha|^2) \frac{|\alpha|^{2n}}{n!} |n\rangle \langle n|. \quad (11)$$

Eq. (11) is implemented in python to visualize a photon number distribution and Wigner plot Eq (8), shown in Fig 4.1. A Fock plot shows the occupation probability p_{nn} . So the goal of this work is to find a n_h which yield a PHAV state, as in [1] For the Wigner function (8) we finally get the following equation

$$w(x, p) = 2 \exp[-2(|\alpha|^2 + |x + ip|^2)] I_0(4|\alpha||x + ip|). \quad (12)$$

I_0 is the first Bessel function. This function is plotted in Fig. 2




Figure 2: The Wigner-Fock plot of the density matrix of ρ_{PHAV} with the parameter $\alpha = 3.5$.

4.2 Double threshold behaviour and population inversion

When we work with density matrices, it is common to work with expectation values with $\langle A \rangle = \text{Tr}[A\rho]$. A is an operator and describe a measurement. With this, we can calculate the expectation value from an Operator. The population inversion is seen by calculating the probability for the occupation number ($\text{Tr}[|i\rangle\langle i|\rho]$) and i stands for the different occupation of the state. Fig. 3a. With high n_h the probability for the system to stay in a the state $|2\rangle$ increases as well. Because the transition between the E_2 and E_1 happened by photon emission and this emission rate is slower as the decay rate from E_3 to E_2 . Lasing starts once a threshold for emission is attained, however, increasing the heat flow too much will end up producing thermal light instead of lasing. This phenomena is called a double threshold behaviour.[2] To investigate this double threshold behaviour, we study the steady states of a three-level system. This double threshold behaviour can be explained by the Zeno effect. The Zeno effect describes that the transition from one state to the other can be stopped by rapid successive measurements. The system behaves similarly when sufficient stimulation is applied quickly. With the plot $\text{Tr}[n\rho] = \langle n \rangle$ against different n_h Fig. 3b we see this double threshold behaviour because, when the hot bath temperature n_h is too low ($n_h \approx n_c$) we have also a low photon number. With increasing n_h , population inversion happens, the photon number increase and the output looks like a PHAV state, see Fig. 2. With too high n_h the expected photon number decrease again and this system shows a double threshold behaviour, the excitation is too weak, and we say that the system is below the lasing threshold. This is another critical point, where the output light becomes a thermal state.




Figure 3: In Fig. 3a the probability for an atom to stay in a state $|0\rangle, |2\rangle$ or $|3\rangle$ vs n_h with the parameters. With a increasing n_h $P2$ increase as well. We call it population inversion. The parameters for the first plot are $n_c = 0.02, n_f = 0.02, \kappa = 0.01\gamma_h$. The blue lines mark the the values for n_h for which the Wigner functions are plotted. In Fig. 3b we see, the the expectation value of the photon number versus different n_h

A other way to see the double threshold behaviour of the three-level system is to analyse the Wigner function plot. For that the reduce density matrices ρ_{free} will be used to make Wigner and Fock plots numerically. For the calculation of the coherent state, the information about the state are not necessary anymore and the system get traced out. It is possible to make the partial trace of ρ , to trace out the reduced density matrices ρ_{free} . The first result are plots from the photon number distribution and the Wigner function. For all calculations, I set the parameters γ_h, γ_c are set to one. And $\kappa = 0.028\gamma_h$ I calculated Wigner functions for this set of parameters, shown in Fig. 4 and Fig. 5.

Those rings which are shown in Fig. 4 and Fig. 5, are similar to the plot of Fig. 2. If we have a small κ means less photons will not leave and stay in the cavity. we see that in the plot of the photon number distribution. In the first plot I set a high leaking-parameter $\kappa = \frac{1}{\gamma_h}$. Shown in Fig. 4. This means that many photons leave the cavity, and only a few remain in the cavity. We see, that the occupation-number the photon number distribution is most zero and the probability for one photon is just 0.1.




Figure 4: The parameters for the first plot are $n_h = 2.6, n_c = 0.001, n_f = 0.02, \kappa = 1\gamma_h$. With the factor of $\kappa = 1\gamma_h$ the leaking is too high for a lasing state.

In the plots of Fig 5 I took the same parameters again, but with a lower κ . We get more photons in the cavity. We see the distribution in the photon number distribution plot and a PHAV state in the Wigner function.




Figure 5: For all of the four the parameters are $n_c = 0.001n_f = 0.02, \kappa = 0.1\gamma_h$. Cavity with a small leaking. **a)** The parameters for n_h are $n_h = 0.2$. **b)** $n_h = 2.6$ **c)** $n_h = 5.5$ **d)** $n_h = 20$. **4b,4c** looks similar to a PHAV state from Fig 2. The blue lines from Fig 4 mark the values of n_h where the plots from Fig 4 are done for.

We see in Fig 5a that start to get a state which looks like PHAV state.

If we increase $2n_h$, then the cavity photon number increases again when increasing n_h at the high temperature regime, as shown in Fig. 5b, Fig 5c. If we further increase $n_h > 10$, we see the double threshold behavior. The state of Fig. 5d is done in a very high temperature region with the value of $n_h = 20$. In this case the output is a thermal state, so no coherent state anymore and the output is below the threshold.

5 Thermodynamics

5.1 Heat currents

The first law of Thermodynamics says: "The change in internal energy of a closed system is equal to the sum of the change in heat and the change in work." That is equal to the equation

$$\Delta U = \Delta Q + \Delta W. \quad (13)$$

In the case of this calculation we have no change in the work. Therefore it is possible to calculate the entropy production(19) only with the heat flow divide through the temperature. To calculate the expected heat flow we can take the trace from $Tr[H_{free}\mathcal{L}[\rho]]$.

$$\langle J_{tot} \rangle = Tr[H_{free} \cdot \mathcal{L}_h[\rho]] + Tr[H_{free} \cdot \mathcal{L}_c[\rho]] + Tr[H_{free} \cdot \mathcal{L}_{cav}[\rho]]. \quad (14)$$

A part of my work is to calculate the occupation number analytically. The calculation is made in two steps. For the hot and the cold bath, we have a transition-operators in the trace. The equation gave the following result:

$$\langle J_h \rangle = Tr[H_{free} \cdot \mathcal{L}_h[\rho]] = \hbar\omega_h\gamma_h(2n_h + 1)(P1 - P3), \quad (15)$$

The same calculation can be done for the interaction with the cold bath.

$$\langle J_c \rangle = Tr[H_{free} \cdot \mathcal{L}_c[\rho]] = \hbar\omega_c\gamma_c(2n_c + 1)(P2 - P3), \quad (16)$$

For the calculation the $Tr[H_{free} \cdot \mathcal{L}_{cav}]$, I get the following result

$$\langle J_{cav} \rangle = Tr[H_{free} \cdot \mathcal{L}_{cav}[\rho]] = 2\hbar\omega_{cav}\kappa(\kappa - \langle a^\dagger a \rangle). \quad (17)$$

The calculations are shown in Fig. 8 and Fig. 9 in the Appendix. With the density matrices it is possible to calculate the heat flux by taking $Tr[H\mathcal{L}\rho]$. and plot this for different g .

In the second step of the calculation we look for $\langle J_{tot} \rangle$ on different coupling constants g . I calculated the expected heat flow. The cold interaction, the hot bath and the interaction with the cavity. Fig. 6 shows for the parameters $n_h = n_c = 2.6, n_{cav} = 0.02, \kappa = 0.01$ is shown below.




Figure 6: Energy flux vs g with the parameters The parameters for the first plot are $n_h = 2.6, n_c = n_f = 0.02, \kappa = 0.01$

5.2 Entropy production

Another relevant concept is the entropy. It is also a physical property. Entropy is described by the second law of thermodynamics. Entropy is defined as the quotient of an infinitesimal amount of heat ΔQ_{rev} to the instantaneous temperature.

$$\Delta S = \frac{Q_{rev}}{T} \quad (18)$$

The entropy production is given by the formula

$$\dot{\sigma} = \frac{J(n_h)}{T(n_c)} + \frac{J(n_c)}{T(n_h)} + \frac{J(n_{cav})}{T(n_{cav})}. \quad (19)$$

It is the time derivative of the entropy. As already seen in Fig. 3b we have a optimum parameter n_h . in Fig. 3a we see the probability for $P1, P2, P3$ is in the region of $n_h = 5$ well distributed. We see also that this corresponds with the average of the photon number. The higher the entropy, the more the state of the three-level system changes and the more photons are emitted. We see that the entropy production correlate with the average of photon number in Fig 3b.




Figure 7: The entropy production for different n_h . The parameters for the first plot are $n_c = 0.001n_f = 0.02, \kappa = 1$. As in paper [2] is used the values for the frequencies $\omega_f = 30\frac{1}{\gamma}, \omega_h = 150\frac{1}{\gamma}, \omega_c = 120\frac{1}{\gamma}$.

6 Conclusion and outlook

In those numerical calculations are found states which look similar to PHAV states, driven by the coupling of two different thermal bath. We found the double threshold behaviour from the system, by calculating different energy flux for different n_h . As in the paper [2]. With the condition that n_c, n_{cav} is almost zero, the leaking κ is small too, and the hot bath received a value between 1 and 5.5. When we compare the Wigner functions of the numerical calculated states from a our three-level system Fig 5b,5c with the Wigner plot of a PHAV state Fig. 2, we see, that they are pretty similar. As conclusion; it's possible to get a phase average coherent state. In Fig. 6 we see the average of the current, calculated with the formula Eq. (14) for different coupling constants g . The conclusion from Fig. 6 is, that the current increase at most between the $0 < g < 0.25$. In Fig. 5a we see the probability in which state the system is. If we increase the temperature from the hot bath, we see that the occupation probability of $P2$ increase as well. A population inversion is visible. Fig. 3 helps to get some conclusion about the system starts lasing and when it stops. With a too small temperature, It has not enough energy in the system and the photon number is low Fig 5a. If n_h increase more, a maximum lasing output is reached Fig 5b,5c. If the temperature increasing further, the system is in the highest level, therefore the Rabi oscillation between E_0 and E_2 does not exist anymore and we get a thermal state Fig 5d. At higher heat of the warm bath, dephasing of the system occurs. Thus, it interacts more with the environment. This has the same effect as if the system is measured often. The Zeno effect then make that the system no longer oscillates between two different states. The consistent numerical and theoretical results we have obtained in the characterization of both PHAV states. In further calculation we will look at the thermodynamic uncertainty relation (TUR) and a second laser as input.

References

- [1] Alessia Allevi et al. “Characterization of phase-averaged coherent states”. In: *Journal of the Optical Society of America B* 30.10 (2013), p. 2621. ISSN: 0740-3224. DOI: 10.1364/josab.30.002621. arXiv: 1302.2011.
- [2] Sheng Wen Li et al. “Quantum statistics of a single-atom Scovil-Schulz-DuBois heat engine”. In: *Physical Review A* 96.6 (2017), pp. 1–10. ISSN: 24699934. DOI: 10.1103/PhysRevA.96.063806. arXiv: 1710.00902. URL: <http://arxiv.org/abs/1710.00902><http://dx.doi.org/10.1103/PhysRevA.96.063806>.

- [3] P. D. Nation and J. R. Johansson. “QuTiP: Quantum Toolbox in Python 4.7.0”. In: *QuTip* (2022), p. 241. URL: <http://qutip.org/documentation.html>.
- [4] Marco Scigliuzzo et al. “Primary Thermometry of Propagating Microwaves in the Quantum Regime”. In: *Physical Review X* 10.4 (2020), p. 41054. ISSN: 21603308. DOI: 10.1103/PhysRevX.10.041054. arXiv: 2003.13522. URL: <https://doi.org/10.1103/PhysRevX.10.041054>.
- [5] H. E.D. Scovil and E. O. Schulz-Dubois. “Three-level masers as heat engines”. In: *Physical Review Letters* 2.6 (1959), pp. 262–263. ISSN: 00319007. DOI: 10.1103/PhysRevLett.2.262. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.2.262>.

7 Appendix

The hole Master equation without any substitution:

$$\begin{aligned}
\mathcal{L}\hat{\rho} = & \frac{\gamma_h}{2} \left[\frac{1}{\exp[\frac{\hbar\omega_h}{k_b T_h}] - 1} + 1 \right] \cdot \left(2\sigma_{13} \cdot \rho \cdot \sigma_{13}^\dagger - \sigma_{13}^\dagger \sigma_{13} \rho - \rho \sigma_{13}^\dagger \sigma_{13} \right) \\
& + \frac{\gamma_h}{2} \left[\frac{1}{\exp[\frac{\hbar\omega_h}{k_b T_H}] - 1} \right] \cdot \left(2\sigma_{31} \cdot \rho \cdot \sigma_{31}^\dagger - \sigma_{31}^\dagger \sigma_{31} \rho - \rho \sigma_{31}^\dagger \sigma_{31} \right) \\
& + \frac{\gamma_c}{2} \left[\frac{1}{\exp[\frac{\hbar\omega_c}{k_b T_c}] - 1} + 1 \right] \cdot \left(2\sigma_{23} \cdot \rho \cdot \sigma_{23}^\dagger - \sigma_{23}^\dagger \sigma_{23} \rho - \rho \sigma_{23}^\dagger \sigma_{23} \right) \\
& + \frac{\gamma_c}{2} \left[\frac{1}{\exp[\frac{\hbar\omega_c}{k_b T_c}] - 1} \right] \cdot \left(2\sigma_{32} \cdot \rho \cdot \sigma_{32}^\dagger - \sigma_{32}^\dagger \sigma_{32} \rho - \rho \sigma_{32}^\dagger \sigma_{32} \right) \\
& + \kappa \left[\frac{1}{\exp[\frac{\hbar\omega_f}{k_b T_f}] - 1} + 1 \right] \cdot \left(2a\rho a^\dagger - a^\dagger a \rho - \rho a^\dagger a \right) \\
& + \kappa \left[\frac{1}{\exp[\frac{\hbar\omega_f}{k_b T_f}] - 1} \right] \cdot \left(2a^\dagger \rho a - a a^\dagger \rho - \rho a a^\dagger \right)
\end{aligned} \tag{20}$$

7.1 Software

For the implementation of the three-level-system in a cavity, I used qutip. Qutip is a python library, which allows to solving master equation fast. Further calculations and methods were easily applied in python.

$$\begin{aligned}
& \text{Tr} [H_{\text{free}} \cdot \mathcal{L}_h g] \\
&= \text{Tr} \left[\sum_{i=0}^3 \hbar \omega_i |i\rangle \langle i| + \hbar \omega_2 a^\dagger a \right] \left(\frac{\gamma_h}{2} [n(\omega_h, T_h) + 1] \cdot (2 \bar{\sigma}_{13} g \bar{\sigma}_{13}^\dagger - \bar{\sigma}_{13}^\dagger \bar{\sigma}_{13} g - g \bar{\sigma}_{13}^\dagger \bar{\sigma}_{13}) + \right. \\
&\quad \left. + \frac{\gamma_h}{2} [\bar{n}(\omega_h, T_h)] (2 \bar{\sigma}_{31} g \bar{\sigma}_{31}^\dagger - \bar{\sigma}_{31}^\dagger \bar{\sigma}_{31} g - g \bar{\sigma}_{31}^\dagger \bar{\sigma}_{31}) \right) \\
&= \text{Tr} \left[\hbar \omega_h \frac{\gamma_h}{2} (n(\omega_h, T_h) + 1) (|3\rangle \langle 3| 2 \bar{\sigma}_{13} g \bar{\sigma}_{13}^\dagger - |3\rangle \langle 3| \bar{\sigma}_{13}^\dagger \bar{\sigma}_{13} g - |3\rangle \langle 3| g \bar{\sigma}_{13}^\dagger \bar{\sigma}_{13}) \right. \\
&\quad + \text{Tr} \left[\hbar \omega_h \frac{\gamma_h}{2} n(\omega_h, T_h) (|3\rangle \langle 3| 2 \bar{\sigma}_{31} g \bar{\sigma}_{31}^\dagger - |3\rangle \langle 3| \bar{\sigma}_{31}^\dagger \bar{\sigma}_{31} g - |3\rangle \langle 3| g \bar{\sigma}_{31}^\dagger \bar{\sigma}_{31}) \right. \\
&\quad + \text{Tr} \left[\hbar \omega_c \frac{\gamma_h}{2} (n(\omega_h, T_h) + 1) (|1\rangle \langle 1| 2 \bar{\sigma}_{13} g \bar{\sigma}_{13}^\dagger - |1\rangle \langle 1| \bar{\sigma}_{13}^\dagger \bar{\sigma}_{13} g - |1\rangle \langle 1| g \bar{\sigma}_{13}^\dagger \bar{\sigma}_{13}) \right. \\
&\quad + \text{Tr} \left[\hbar \omega_c \frac{\gamma_h}{2} n(\omega_h, T_h) (|1\rangle \langle 1| 2 \bar{\sigma}_{31} g \bar{\sigma}_{31}^\dagger - |1\rangle \langle 1| \bar{\sigma}_{31}^\dagger \bar{\sigma}_{31} g - |1\rangle \langle 1| g \bar{\sigma}_{31}^\dagger \bar{\sigma}_{31}) \right. \\
&\quad \left. + \text{Tr} [\hbar \omega_2 |2\rangle \langle 2| \cdot \mathcal{L}_h g] \right] = 0 \\
&\quad + \text{Tr} [\hbar \omega_h a \cdot \mathcal{L}_h g] = 0 \quad // \text{Kommutiert} \\
&= \text{Tr} \left[\hbar \omega_h \frac{\gamma_h}{2} (n(\omega_h, T_h) + 1) (2 \cdot |3\rangle \langle 3| \bar{\sigma}_{13} g \bar{\sigma}_{13}^\dagger - |3\rangle \langle 3| \bar{\sigma}_{13}^\dagger \bar{\sigma}_{13} g - |3\rangle \langle 3| g \bar{\sigma}_{13}^\dagger \bar{\sigma}_{13}) \right. \\
&\quad + \text{Tr} \left[\hbar \omega_h \frac{\gamma_h}{2} n(\omega_h, T_h) \cdot (2 |3\rangle \langle 3| 3\rangle \langle 1| g \bar{\sigma}_{13}^\dagger) = 4 (2 \cdot \bar{\sigma}_{31} g \bar{\sigma}_{13}^\dagger) = 4 P_1 \right. \\
&\quad + \text{Tr} \left[\hbar \omega_c \frac{\gamma_h}{2} n(\omega_h, T_h) \cdot |1\rangle \langle 1| (2 \cdot \bar{\sigma}_{13} g \bar{\sigma}_{13}^\dagger) = 2 |1\rangle \langle 1| 1\rangle \langle 3| g |3\rangle \langle 1| = 2 P_3 \right. \\
&\quad \left. + \text{Tr} \left[\hbar \omega_c \frac{\gamma_h}{2} n(\omega_h, T_h) \cdot |1\rangle \langle 1| 2 \bar{\sigma}_{31} g \bar{\sigma}_{31}^\dagger - |1\rangle \langle 1| g \bar{\sigma}_{31}^\dagger \bar{\sigma}_{31} g = -2 P_1 \right. \right. \\
&\quad \left. \left. - |3\rangle \langle 3| g \bar{\sigma}_{13}^\dagger \bar{\sigma}_{13} = g |3\rangle \langle 1| 1\rangle \langle 3| = P_3 \right. \right. \\
&\quad \left. \left. \begin{aligned} &= -\hbar \omega_h \gamma_h \cdot n(\omega_h, T_h) + 1 \cdot P_3 \\ &+ \hbar \omega_h \gamma_h \cdot n(\omega_h, T_h) \cdot P_3 \end{aligned} \right\} = -\hbar \omega_h \gamma_h P_3 \right. \\
&\quad \left. \left. \begin{aligned} &+ \hbar \omega_c \gamma_h \cdot (n(\omega_h, T_h) + 1) \cdot P_1 \\ &- \hbar \omega_c \gamma_h \cdot n(\omega_h, T_h) \cdot P_2 \end{aligned} \right\} = +\hbar \omega_c \gamma_h P_1 \right. \\
&= \hbar \gamma_h \cdot (\omega_c P_1 - \omega_h P_3)
\end{aligned}$$

Figure 8: Berechnung-Tr[HL]

$$\begin{aligned}
& \text{Tr} [H_{\text{tree}} \cdot \mathcal{L}_{\text{cav}} g] \\
&= \left(\sum_{i \leftarrow d} \hbar \omega_i |i\rangle \langle i| + \hbar \omega a^\dagger a \right) \cdot [k(\bar{n}+1) \cdot (2 a g a^\dagger - a^\dagger a g - g a^\dagger a)] \\
&+ \left(\sum_{i \leftarrow d} \hbar \omega_i |i\rangle \langle i| + \hbar \omega a^\dagger a \right) \cdot [k \bar{n} \cdot (2 a^\dagger g a - a a^\dagger g - g a a^\dagger)] \\
&= \text{Tr} [\hbar \omega k(\bar{n}+1) [2 a^\dagger a a g a^\dagger - a^\dagger a a^\dagger a g - a^\dagger a g a^\dagger a] \\
&+ \hbar \omega k \bar{n} [2 a^\dagger a a^\dagger g a - a^\dagger a a^\dagger a g - a^\dagger a g a a^\dagger] \\
&= \hbar \omega k(\bar{n}+1) [2 \text{Tr} a^\dagger a^\dagger a a g] \ominus 2 \text{Tr} [a^\dagger a a^\dagger a g] \quad \begin{matrix} \nearrow 2 \text{Tr} [a^\dagger a g] - \text{Tr} [a^\dagger a a a] \\ 0_a \quad a a^\dagger - a^\dagger a = 1 \end{matrix} \\
&+ \hbar \omega k \bar{n} \cdot [2 \text{Tr} [a a^\dagger a a^\dagger g] - \text{Tr} [a^\dagger a a^\dagger a g] - \text{Tr} [a a^\dagger a^\dagger g] \\
&= \hbar \omega k(\bar{n}+1) \cdot \text{Tr} [a^\dagger a g] \\
&+ \hbar \omega k \bar{n} \cdot \text{Tr} [a a^\dagger (a a^\dagger - a^\dagger a) g] + \text{Tr} [a a^\dagger a a^\dagger g] - \text{Tr} [a^\dagger a a a^\dagger g] \\
&= \hbar \omega k(\bar{n}+1) 2 \text{Tr} [a^\dagger a g] + \\
&+ \hbar \omega k \bar{n} 2 \text{Tr} [a a^\dagger g] + \text{Tr} [(a a^\dagger - a^\dagger a) a a^\dagger g] \\
&= \hbar \omega k(\bar{n}+1) 2 \text{Tr} [a^\dagger a g] + \hbar \omega k \bar{n} 2 \text{Tr} [a a^\dagger g] \\
&= 2 \hbar \omega k (\bar{n} - \langle a^\dagger a \rangle)
\end{aligned}$$

Figure 9: Berechnung- $\text{Tr}[\text{HL}]$.