

# Mixed models applied to breeding

Alencar Xavier

March 13th - March 15th, 2019

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

1/171

## Instructors

### Alencar Xavier

- Quantitative Geneticist, Corteva Agrisciences
- Adjunct professor, Dpt. of Agronomy, Purdue University

### Luiz Brito

- Assistant Professor of Animal Sciences, Purdue University

### Hinayah Oliveira

- Post-doc, Dtp. of Animal Sciences, Purdue University

### Katy Rainey

- Associate Professor, Dpt. of Agronomy, Purdue University

# Schedule

- Module 1: Intro to mixed models
- Module 2: Fitting mixed models
- Module 3: Advanced topics
- Module 4: Signal detection
- Module 5: Association analysis

3/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

3/171

## Module 1 - Introduction to mixed models

# Outline

## Part 1: Concepts

- History of mixed models
- Mixed models in plant breeding
- Fixed and random terms
- Model notation
- Variance decomposition

## Part 2: Applications

- Selection models
- Practical examples
- Variance components
- Ridges and Kernels

5/171

[file:///C:/Users/rd7564/Desktop/Purdue\\_MLM/MyLecture/Purdue\\_MLM.html#8](file:///C:/Users/rd7564/Desktop/Purdue_MLM/MyLecture/Purdue_MLM.html#8)

5/171

# Part 1 - Concepts

6/171

[file:///C:/Users/rd7564/Desktop/Purdue\\_MLM/MyLecture/Purdue\\_MLM.html#8](file:///C:/Users/rd7564/Desktop/Purdue_MLM/MyLecture/Purdue_MLM.html#8)

6/171

# History of mixed models

*Francis Galton* - [1886](#): Regression and heritability

*Ronald Fisher* - [1918](#): Infinitesimal model ( $P = G + E$ )

*Sewall Wright* - [1922](#): Genetic relationship

*Charles Henderson* - [1968](#): BLUP using relationship



7/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

7/171

## Mixed models in plant breeding

- *Heart and soul* of plant breeding ([Xavier et al 2017](#))
- Variance components and heritability ([Johnson and Thompson 1994](#))
- Trait associations ([Gianola and Sorensen 2014](#))
- Estimation of genetic and breeding values ([Piepho et al 2008](#))
- Prediction of unphenotyped lines ([de los Campos et al 2013](#))
- Selection index ([Wientjes et al. 2016](#))
- Genome-wide association analysis ([Yang et al 2014](#))
- All sorts of inference ([Robinson 1991](#))

8/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

8/171

# Fixed and random terms

## Fixed effect

- Assumed to be invariable (often you cannot recollect the data)
- Inferences are made upon the parameters
- Results can not be extrapolated to other datasets
- Example: Overall mean and environmental effects

## Random effects

- You may not have all the levels available
- Inference are made on variance components
- Prior assumption: coefficients are normally distributed
- Results can not be extrapolated to other datasets
- Example: Genetic effects

9/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

9/171

# Let's unleash the beast



10/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

10/171

# Model notation

- Linear model:  $y = Xb + Zu + e$
- With variance:  $y \sim N(Xb, ZKZ\sigma_u^2 + I\sigma_e^2)$

Assuming:  $u \sim N(0, K\sigma_u^2)$  and  $e \sim N(0, I\sigma_e^2)$

Henderson equation

$$\begin{bmatrix} X'X & Z'X \\ X'Z & Z'Z + \lambda K^{-1} \end{bmatrix} \begin{bmatrix} b \\ u \end{bmatrix} = \begin{bmatrix} X'y \\ Z'y \end{bmatrix}$$

Summary:

- We know (data):  $x = \{y, X, Z, K\}$
- We want (parameters):  $\theta = \{b, u, \sigma_a^2, \sigma_e^2\}$
- Estimation based on Gaussian likelihood:  $L(x|\theta)$

11/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

11/171

# Model notation

- **y** = vector of observations ( $n$ )
- **X** = design matrix of fixed effects ( $n \times p$ )
- **Z** = design (or incidence) matrix of random effects ( $n \times q$ )
- **K** = random effect correlation matrix ( $q \times q$ )
- **u** = vector of random effect coefficients ( $q$ )
- **b** = vector of fixed effect coefficients ( $p$ )
- **e** = vector of residuals ( $n$ )
- $\sigma_a^2$  = marker effect variance (1)
- $\sigma_u^2$  = random effect variance (1)
- $\sigma_e^2$  = residual variance (1)
- $\lambda = \sigma_e^2 / \sigma_u^2$  (Regularization parameters) (1)

12/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

12/171

# Model notation

The mixed model can also be notated as follows

$$y = Wg + e$$

Solved as

$$[W'W + \Sigma]g = [W'y]$$


Where

$$W = [X, Z]$$

$$g = [b, u]$$

$$\Sigma = \begin{bmatrix} 0 & 0 \\ 0 & \lambda K^{-1} \end{bmatrix}$$

# Variance decomposition



# Part 2 - Applications

15/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

15/171

3/18/2019

Mixed models applied to breeding

## Selection

### 1 *Genetic values*

- BLUPs or BLUEs from replicated trials
- Captures additive and non-additive genetics together

### 2 *Breeding values*

- Use pedigree information to create  $K$
- Captures additive genetics (heritable)
- Trials not necessarily replicated

### 3 *Genomic Breeding values*

- Genotypic information replaces pedigree
- Any signal: additivity, dominance and epistasis

16/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

16/171



# Examples

- Example 1: Balanced data, no kinship
- Example 2: Balanced data, with kinship
- Example 3: Unbalanced data, with kinship
- Example 4: Balanced data, missing individual

17/171

17/171

# Example 1

Data:

| ##    |  | Env | Gen | Phe |
|-------|--|-----|-----|-----|
| ## 1  |  | E1  | G1  | 47  |
| ## 2  |  | E1  | G2  | 51  |
| ## 3  |  | E1  | G3  | 46  |
| ## 4  |  | E1  | G4  | 58  |
| ## 5  |  | E2  | G1  | 52  |
| ## 6  |  | E2  | G2  | 46  |
| ## 7  |  | E2  | G3  | 52  |
| ## 8  |  | E2  | G4  | 54  |
| ## 9  |  | E3  | G1  | 53  |
| ## 10 |  | E3  | G2  | 48  |
| ## 11 |  | E3  | G3  | 58  |
| ## 12 |  | E3  | G4  | 52  |

Model:  $Phenotype = Environment_{(F)} + Genotype_{(R)}$

18/171

18/171

# Example 1

Design matrix  $W$ :

| ##    | EnvE1 | EnvE2 | EnvE3 | GenG1 | GenG2 | GenG3 | GenG4 |
|-------|-------|-------|-------|-------|-------|-------|-------|
| ## 1  | 1     | 0     | 0     | 1     | 0     | 0     | 0     |
| ## 2  | 1     | 0     | 0     | 0     | 1     | 0     | 0     |
| ## 3  | 1     | 0     | 0     | 0     | 0     | 1     | 0     |
| ## 4  | 1     | 0     | 0     | 0     | 0     | 0     | 1     |
| ## 5  | 0     | 1     | 0     | 1     | 0     | 0     | 0     |
| ## 6  | 0     | 1     | 0     | 0     | 1     | 0     | 0     |
| ## 7  | 0     | 1     | 0     | 0     | 0     | 1     | 0     |
| ## 8  | 0     | 1     | 0     | 0     | 0     | 0     | 1     |
| ## 9  | 0     | 0     | 1     | 1     | 0     | 0     | 0     |
| ## 10 | 0     | 0     | 1     | 0     | 1     | 0     | 0     |
| ## 11 | 0     | 0     | 1     | 0     | 0     | 1     | 0     |
| ## 12 | 0     | 0     | 1     | 0     | 0     | 0     | 1     |

19/171

# Example 1

$W'W$ :

| ##       | EnvE1 | EnvE2 | EnvE3 | GenG1 | GenG2 | GenG3 | GenG4 |
|----------|-------|-------|-------|-------|-------|-------|-------|
| ## EnvE1 | 4     | 0     | 0     | 1     | 1     | 1     | 1     |
| ## EnvE2 | 0     | 4     | 0     | 1     | 1     | 1     | 1     |
| ## EnvE3 | 0     | 0     | 4     | 1     | 1     | 1     | 1     |
| ## GenG1 | 1     | 1     | 1     | 3     | 0     | 0     | 0     |
| ## GenG2 | 1     | 1     | 1     | 0     | 3     | 0     | 0     |
| ## GenG3 | 1     | 1     | 1     | 0     | 0     | 3     | 0     |
| ## GenG4 | 1     | 1     | 1     | 0     | 0     | 0     | 3     |

20/171

# Example 1

Left-hand side ( $W'W + \Sigma$ ):

| ##       | EnvE1 | EnvE2 | EnvE3 | GenG1 | GenG2 | GenG3 | GenG4 |
|----------|-------|-------|-------|-------|-------|-------|-------|
| ## EnvE1 | 4     | 0     | 0     | 1.00  | 1.00  | 1.00  | 1.00  |
| ## EnvE2 | 0     | 4     | 0     | 1.00  | 1.00  | 1.00  | 1.00  |
| ## EnvE3 | 0     | 0     | 4     | 1.00  | 1.00  | 1.00  | 1.00  |
| ## GenG1 | 1     | 1     | 1     | 3.17  | 0.00  | 0.00  | 0.00  |
| ## GenG2 | 1     | 1     | 1     | 0.00  | 3.17  | 0.00  | 0.00  |
| ## GenG3 | 1     | 1     | 1     | 0.00  | 0.00  | 3.17  | 0.00  |
| ## GenG4 | 1     | 1     | 1     | 0.00  | 0.00  | 0.00  | 3.17  |

Assuming independent individuals:  $K = I$

Regularization:  $\lambda = \sigma_e^2 / \sigma_u^2 = 1.64 / 9.56 = 0.17$

21/171

21/171

# Example 1

Right-hand side ( $W'y$ ):

|          |      |
|----------|------|
| ##       | [,1] |
| ## EnvE1 | 202  |
| ## EnvE2 | 204  |
| ## EnvE3 | 211  |
| ## GenG1 | 152  |
| ## GenG2 | 145  |
| ## GenG3 | 156  |
| ## GenG4 | 164  |

22/171

22/171

# Example 1

We can find coefficients through least-square solution

$$g = (LHS)^{-1}(RHS) = (W'W + \Sigma)^{-1}W'y$$

```
##      [,1]
## EnvE1 50.50
## EnvE2 51.00
## EnvE3 52.75
## GenG1 -0.71
## GenG2 -2.92
## GenG3  0.55
## GenG4  3.08
```

23/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

23/171

## Shrinkage

$$BLUE = \frac{w'y}{w'w} = \frac{sum}{n} = \text{simple average}$$

$$BLUP = \frac{w'y}{w'w + \lambda} = \frac{sum}{n + \lambda} = \text{biased average} = BLUE \times h^2$$



**Note:**

- More observations = less shrinkage
- Higher heritability = less shrinkage:  $\lambda = \frac{h^2 - 1}{h^2}$

24/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

24/171

# Example 2

If we know the relationship among individuals:

| ## |       | GenG1 | GenG2 | GenG3 | GenG4 |
|----|-------|-------|-------|-------|-------|
| ## | GenG1 | 1.00  | 0.64  | 0.23  | 0.48  |
| ## | GenG2 | 0.64  | 1.00  | 0.33  | 0.67  |
| ## | GenG3 | 0.23  | 0.33  | 1.00  | 0.31  |
| ## | GenG4 | 0.48  | 0.67  | 0.31  | 1.00  |

25/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

25/171

# Example 2

Then we estimate  $\lambda K^{-1}$

| ## |       | GenG1 | GenG2 | GenG3 | GenG4 |
|----|-------|-------|-------|-------|-------|
| ## | GenG1 | 0.15  | -0.09 | 0.00  | -0.01 |
| ## | GenG2 | -0.09 | 0.22  | -0.02 | -0.10 |
| ## | GenG3 | 0.00  | -0.02 | 0.10  | -0.02 |
| ## | GenG4 | -0.01 | -0.10 | -0.02 | 0.17  |

Regularization:  $\lambda = \sigma_e^2 / \sigma_u^2 = 1.64 / 17.70 = 0.09$

26/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

26/171

# Example 2

And the left-hand side becomes

| ## |       | EnvE1 | EnvE2 | EnvE3 | GenG1 | GenG2 | GenG3 | GenG4 |
|----|-------|-------|-------|-------|-------|-------|-------|-------|
| ## | EnvE1 | 4     | 0     | 0     | 1.00  | 1.00  | 1.00  | 1.00  |
| ## | EnvE2 | 0     | 4     | 0     | 1.00  | 1.00  | 1.00  | 1.00  |
| ## | EnvE3 | 0     | 0     | 4     | 1.00  | 1.00  | 1.00  | 1.00  |
| ## | GenG1 | 1     | 1     | 1     | 3.15  | -0.09 | 0.00  | -0.01 |
| ## | GenG2 | 1     | 1     | 1     | -0.09 | 3.22  | -0.02 | -0.10 |
| ## | GenG3 | 1     | 1     | 1     | 0.00  | -0.02 | 3.10  | -0.02 |
| ## | GenG4 | 1     | 1     | 1     | -0.01 | -0.10 | -0.02 | 3.17  |

27/171

27/171

# Example 2

We can find coefficients through least-square solution

$$g = (LHS)^{-1} (RHS) = (W'W + \Sigma)^{-1} W'y$$

|    |             |
|----|-------------|
| ## | [,1]        |
| ## | EnvE1 51.05 |
| ## | EnvE2 51.55 |
| ## | EnvE3 53.30 |
| ## | GenG1 -1.32 |
| ## | GenG2 -3.34 |
| ## | GenG3 0.03  |
| ## | GenG4 2.45  |

Genetic coefficients shrink more: Var(A) < Var(G)

28/171

28/171

# Example 3

What if we have missing data?

| ##    |  | Env | Gen | Phe |
|-------|--|-----|-----|-----|
| ## 1  |  | E1  | G1  | 47  |
| ## 2  |  | E1  | G2  | 51  |
| ## 3  |  | E1  | G3  | NA  |
| ## 4  |  | E1  | G4  | 58  |
| ## 5  |  | E2  | G1  | 52  |
| ## 6  |  | E2  | G2  | 46  |
| ## 7  |  | E2  | G3  | 52  |
| ## 8  |  | E2  | G4  | NA  |
| ## 9  |  | E3  | G1  | 53  |
| ## 10 |  | E3  | G2  | 48  |
| ## 11 |  | E3  | G3  | 58  |
| ## 12 |  | E3  | G4  | 52  |

29/171

# Example 3

Rows of missing points are removed

| ##    | EnvE1 | EnvE2 | EnvE3 | GenG1 | GenG2 | GenG3 | GenG4 |
|-------|-------|-------|-------|-------|-------|-------|-------|
| ## 1  | 1     | 0     | 0     | 1     | 0     | 0     | 0     |
| ## 2  | 1     | 0     | 0     | 0     | 1     | 0     | 0     |
| ## 4  | 1     | 0     | 0     | 0     | 0     | 0     | 1     |
| ## 5  | 0     | 1     | 0     | 1     | 0     | 0     | 0     |
| ## 6  | 0     | 1     | 0     | 0     | 1     | 0     | 0     |
| ## 7  | 0     | 1     | 0     | 0     | 0     | 1     | 0     |
| ## 9  | 0     | 0     | 1     | 1     | 0     | 0     | 0     |
| ## 10 | 0     | 0     | 1     | 0     | 1     | 0     | 0     |
| ## 11 | 0     | 0     | 1     | 0     | 0     | 1     | 0     |
| ## 12 | 0     | 0     | 1     | 0     | 0     | 0     | 1     |

30/171

# Example 3

$W'W$ :

| ##       | EnvE1 | EnvE2 | EnvE3 | GenG1 | GenG2 | GenG3 | GenG4 |
|----------|-------|-------|-------|-------|-------|-------|-------|
| ## EnvE1 | 3     | 0     | 0     | 1     | 1     | 0     | 1     |
| ## EnvE2 | 0     | 3     | 0     | 1     | 1     | 1     | 0     |
| ## EnvE3 | 0     | 0     | 4     | 1     | 1     | 1     | 1     |
| ## GenG1 | 1     | 1     | 1     | 3     | 0     | 0     | 0     |
| ## GenG2 | 1     | 1     | 1     | 0     | 3     | 0     | 0     |
| ## GenG3 | 0     | 1     | 1     | 0     | 0     | 2     | 0     |
| ## GenG4 | 1     | 0     | 1     | 0     | 0     | 0     | 2     |

31/171

31/171

# Example 3

Left-hand side ( $W'W + \Sigma$ ):

| ##       | EnvE1 | EnvE2 | EnvE3 | GenG1 | GenG2 | GenG3 | GenG4 |
|----------|-------|-------|-------|-------|-------|-------|-------|
| ## EnvE1 | 3     | 0     | 0     | 1.00  | 1.00  | 0.00  | 1.00  |
| ## EnvE2 | 0     | 3     | 0     | 1.00  | 1.00  | 1.00  | 0.00  |
| ## EnvE3 | 0     | 0     | 4     | 1.00  | 1.00  | 1.00  | 1.00  |
| ## GenG1 | 1     | 1     | 1     | 3.10  | -0.06 | 0.00  | -0.01 |
| ## GenG2 | 1     | 1     | 1     | -0.06 | 3.15  | -0.01 | -0.07 |
| ## GenG3 | 0     | 1     | 1     | 0.00  | -0.01 | 2.07  | -0.01 |
| ## GenG4 | 1     | 0     | 1     | -0.01 | -0.07 | -0.01 | 2.11  |

Regularization:  $\lambda = \sigma_c^2 / \sigma_u^2 = 1.21 / 19.61 = 0.06$

32/171

32/171



# Example 3

Right-hand side ( $W'y$ ):

```
##      [,1]
## EnvE1 156
## EnvE2 150
## EnvE3 211
## GenG1 152
## GenG2 145
## GenG3 110
## GenG4 110
```

# Example 3

Find coefficients through least-square solution

$$g = (LHS)^{-1} (RHS) = (W'W + \Sigma)^{-1} W'y$$

```
##      [,1]
## EnvE1 54.14
## EnvE2 51.70
## EnvE3 53.82
## GenG1 -2.56
## GenG2 -4.68
## GenG3 2.15
## GenG4 0.81
```

# Example 4

What if we are missing data from a individual?

| ##    |  | Env | Gen | Phe |
|-------|--|-----|-----|-----|
| ## 1  |  | E1  | G1  | NA  |
| ## 2  |  | E1  | G2  | 51  |
| ## 3  |  | E1  | G3  | 46  |
| ## 4  |  | E1  | G4  | 58  |
| ## 5  |  | E2  | G1  | NA  |
| ## 6  |  | E2  | G2  | 46  |
| ## 7  |  | E2  | G3  | 52  |
| ## 8  |  | E2  | G4  | 54  |
| ## 9  |  | E3  | G1  | NA  |
| ## 10 |  | E3  | G2  | 48  |
| ## 11 |  | E3  | G3  | 58  |
| ## 12 |  | E3  | G4  | 52  |

35/171

# Example 4

Rows of missing points are removed

| ##    | EnvE1 | EnvE2 | EnvE3 | GenG1 | GenG2 | GenG3 | GenG4 |
|-------|-------|-------|-------|-------|-------|-------|-------|
| ## 2  | 1     | 0     | 0     | 0     | 1     | 0     | 0     |
| ## 3  | 1     | 0     | 0     | 0     | 0     | 1     | 0     |
| ## 4  | 1     | 0     | 0     | 0     | 0     | 0     | 1     |
| ## 6  | 0     | 1     | 0     | 0     | 1     | 0     | 0     |
| ## 7  | 0     | 1     | 0     | 0     | 0     | 1     | 0     |
| ## 8  | 0     | 1     | 0     | 0     | 0     | 0     | 1     |
| ## 10 | 0     | 0     | 1     | 0     | 1     | 0     | 0     |
| ## 11 | 0     | 0     | 1     | 0     | 0     | 1     | 0     |
| ## 12 | 0     | 0     | 1     | 0     | 0     | 0     | 1     |

36/171

# Example 4

$W'W$ :

| ##       | EnvE1 | EnvE2 | EnvE3 | GenG1 | GenG2 | GenG3 | GenG4 |
|----------|-------|-------|-------|-------|-------|-------|-------|
| ## EnvE1 | 3     | 0     | 0     | 0     | 1     | 1     | 1     |
| ## EnvE2 | 0     | 3     | 0     | 0     | 1     | 1     | 1     |
| ## EnvE3 | 0     | 0     | 3     | 0     | 1     | 1     | 1     |
| ## GenG1 | 0     | 0     | 0     | 0     | 0     | 0     | 0     |
| ## GenG2 | 1     | 1     | 1     | 0     | 3     | 0     | 0     |
| ## GenG3 | 1     | 1     | 1     | 0     | 0     | 3     | 0     |
| ## GenG4 | 1     | 1     | 1     | 0     | 0     | 0     | 3     |

37/171

37/171

# Example 4

Left-hand side ( $W'W + \Sigma$ ):

| ##       | EnvE1 | EnvE2 | EnvE3 | GenG1 | GenG2 | GenG3 | GenG4 |
|----------|-------|-------|-------|-------|-------|-------|-------|
| ## EnvE1 | 3     | 0     | 0     | 0.00  | 1.00  | 1.00  | 1.00  |
| ## EnvE2 | 0     | 3     | 0     | 0.00  | 1.00  | 1.00  | 1.00  |
| ## EnvE3 | 0     | 0     | 3     | 0.00  | 1.00  | 1.00  | 1.00  |
| ## GenG1 | 0     | 0     | 0     | 0.14  | -0.08 | 0.00  | -0.01 |
| ## GenG2 | 1     | 1     | 1     | -0.08 | 3.19  | -0.02 | -0.09 |
| ## GenG3 | 1     | 1     | 1     | 0.00  | -0.02 | 3.09  | -0.01 |
| ## GenG4 | 1     | 1     | 1     | -0.01 | -0.09 | -0.01 | 3.15  |

Regularization:  $\lambda = \sigma_e^2 / \sigma_u^2 = 1.79 / 22.78 = 0.08$

38/171

38/171

# Example 4

Right-hand side ( $W'y$ ):

```
##      [,1]
## EnvE1 155
## EnvE2 152
## EnvE3 158
## GenG1  0
## GenG2 145
## GenG3 156
## GenG4 164
```

# Example 4

Find coefficients through least-square solution

$$g = (LHS)^{-1} (RHS) = (W'W + \Sigma)^{-1} W'y$$

```
##      [,1]
## EnvE1 52.06
## EnvE2 51.06
## EnvE3 53.06
## GenG1 -1.82
## GenG2 -3.48
## GenG3 -0.07
## GenG4  2.38
```

# Variance components

Expectation-Maximization REML ([1977](#))

$$\sigma_u^2 = \frac{u'K^{-1}u}{q - \lambda \text{tr}(K^{-1}C^{22})} \text{ and } \sigma_e^2 = \frac{e'y}{n-p}$$

Bayesian Gibbs Sampling ([1993](#))

$$\sigma_u^2 = \frac{u'K^{-1}u + S_u\nu_u}{\chi^2(q + \nu_u)} \text{ and } \sigma_e^2 = \frac{e'e + S_e\nu_e}{\chi^2(n + \nu_e)}$$

Predicted Residual Error Sum of Squares (PRESS) ([2017](#))

- $\lambda = \text{argmin}(\sum e_i^2 / (1 - h_{ii})^2)$
- Where  $H = (I + K\lambda)^{-1}$  and  $e = y - \mu - Hy$

41/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

41/171

# Ridges and Kernels

Kernel methods:

- Genetic signal is captured by the relationship matrix  $K$
- Random effect coefficients are the **breeding values** (BV)
- Efficient to compute BV when *markers*  $\gg$  *individuals*
- Easy use and combine pedigree, markers and interactions

Ridge methods:

- Genetic signal is captured by the design matrix  $M$
- Random effect coefficients are the **marker effects**
- Easy way to make predictions of unobserved individuals
- Enables to visualize where the QTLs are in the genome

42/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

42/171

# Ridges and Kernels

Kernel

$$y = Xb + Zu + e, u \sim N(0, K\sigma_u^2)$$

Ridge

$$y = Xb + Ma + e, a \sim N(0, I\sigma_a^2)$$

Where

- $M$  is the genotypic matrix,  $m_{ij} = \{0, 1, 2\}$
- $K = \alpha MM'$
- $u = Ma$
- $\sigma_a^2 = \alpha\sigma_u^2$

43/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

43/171

# Ridges and Kernels

Kernel model

$$\begin{bmatrix} X'X & Z'X \\ X'Z & Z'Z + K^{-1}(\sigma_e^2/\sigma_u^2) \end{bmatrix} \begin{bmatrix} b \\ u \end{bmatrix} = \begin{bmatrix} X'y \\ Z'y \end{bmatrix}$$

Ridge model

$$\begin{bmatrix} X'X & M'X \\ X'M & M'M + I^{-1}(\sigma_e^2/\sigma_a^2) \end{bmatrix} \begin{bmatrix} b \\ a \end{bmatrix} = \begin{bmatrix} X'y \\ M'y \end{bmatrix}$$

Both models capture same genetic signal ([de los Campos 2015](#))


44/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

44/171

# Ridges and Kernels

```
K = tcrossprod(M)/ncol(M)
GBLUP = reml(y=y,K=K); Kernel_GEBV = GBLUP$EBV
RRBLUP = reml(y=y,Z=M); Ridge_GEBV = M%*%RRBLUP$EBV
plot(Kernel_GEBV,Ridge_GEBV, main='Comparing results')
```



45/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

45/171

# Break

# Module 2 - Fitting mixed models

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

47/171

## Example 1 - Sorghum



# Example 1 - load data

Example dataset from Kevin's *agridat* package

```
data(adugna.sorghum, package = 'agridat')
dt = adugna.sorghum
head(dt)
```

```
##   gen trial env yield year   loc
## 1 G16    T2 E01   590 2001 Mieso
## 2 G17    T2 E01   554 2001 Mieso
## 3 G18    T2 E01   586 2001 Mieso
## 4 G19    T2 E01   738 2001 Mieso
## 5 G20    T2 E01   489 2001 Mieso
## 6 G21    T2 E01   684 2001 Mieso
```

49/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

49/171

# Example 1 - Getting design matrix

- Linear model:  $Phenotype = Env + Gen$
- In algebra notation:  $y = Xb + Zu + e$

```
y = dt$yield
X = model.matrix(y~env,dt)
Z = model.matrix(y~gen-1,dt) # "-1" means no intercept
```

Assuming:

- $u \sim N(0, I\sigma_g^2)$
- $e \sim N(0, I\sigma_e^2)$

50/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

50/171


## Example 1 - Visualize X and Z matrices

```
SEE=function(A,...)image(t(1-A[nrow(A):1,]),axes=F,col=gray.colors(2),...)
par(mfrow=c(1,2),mar=c(1,1,3,1))
SEE(X, main=paste("X matrix (",paste(dim(X),collapse=' x '),")" ))
SEE(Z, main=paste("Z matrix (",paste(dim(Z),collapse=' x '),")" ))
```

X matrix ( 289 x 13 )



Z matrix ( 289 x 28 )



51/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

51/171

## Example 1 - Fit the model

```
# Using the NAM package (same for rrBLUP, EMMREML, BGLR)
require(NAM, quietly = TRUE)
fit1 = reml(y=y,X=X,Z=Z)
# Alternatively, you can also use formulas with NAM
fit1b = reml(y=dt$yield,X=~dt$env,Z=~dt$gen )
# Using the lme4 package
require(lme4, quietly = TRUE)
fit2 = lmer(yield ~ env + (1|gen), data=dt)
```

52/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

52/171

# Example 1 - Variance components

```
fit1$VC[c(1:2)] # same with fit1b$VC
```

```
##           Vg           Ve
## 1 189680.4 442075.6
```

```
data.frame((summary(fit2))$varcor)$vcov
```

```
## [1] 189680.4 442075.6
```

- VC can be used to measure *broad-sense* heritability

$$H = \frac{\sigma_g^2}{\sigma_g^2 + \sigma_e^2/n} = \frac{189680.4}{189680.4 + 442075.6/10.32} = 0.82$$


53/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

53/171

# Example 1 - The coefficients

```
blue1 = fit1$Fixed[,1]; blue1 = fit1$EBV
blue2 = fit2@beta; blue2 = rowMeans(ranef(fit2)$gen)
par(mfrow=c(1,2));
plot(blue1,blue2,main="BLUE"); plot(blup1,blup2,main="BLUP")
```



54/171

# Example 1 - DIY BLUPs

```

iK = diag(ncol(Z))
Lambda = 442075.6/189680.4
W = cbind(X,Z)
Sigma = as.matrix(Matrix::bdiag(diag(0,ncol(X)),iK*Lambda))
LHS = crossprod(W) + Sigma
RHS = crossprod(W,y)
g = solve(LHS,RHS)
my_blue = g[ c(1:ncol(X))]
my_blup = g[-c(1:ncol(X))]

```

55/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8


55/171

# Example 1 - DIY BLUPs

```

par(mfrow=c(1,2))
plot(my_blue,blue1,main="BLUE")
plot(my_blup,blup1,main="BLUP")

```



56/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

56/171

# Example 1 - DIY Variance components

$$\sigma_e^2 = \frac{e'y}{n-p} \text{ and } \sigma_u^2 = \frac{u'K^{-1}u + \text{tr}(K^{-1}C^{22}\sigma_e^2)}{q}$$

```
e = y - X %%% my_blue - Z %%% my_blup
Ve = c(y%%e)/(length(y)-ncol(X))
Ve

## [1] 442075.6

trKC22 = sum(diag(ik%%(solve(LHS)[-c(1:ncol(X)), -c(1:ncol(X))])))
Vg = Vg = c(t(my_blup)%%ik%%my_blup+trKC22*Ve)/ncol(Z)
Vg

## [1] 189680.4
```

57/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

57/171

# Starting from bad variance components

```
Ve = Vg = 1
for(i in 1:25){
  Lambda = Ve/Vg;
  Sigma = as.matrix(Matrix::bdiag(diag(0,ncol(X)),ik*Lambda))
  LHS = crossprod(W) + Sigma; RHS = crossprod(W,y); g = solve(LHS,RHS)
  my_blue = g[ c(1:ncol(X))]; my_blup = g[-c(1:ncol(X))]
  e = y - X%%my_blue - Z%%my_blup; Ve = c(y%%e)/(length(y)-ncol(X))
  trKC22 = sum(diag(ik%%(solve(LHS)[(ncol(X)+1):(ncol(W)),(ncol(X)+1):(ncol(W)))]))
  Vg = c(t(my_blup)%%ik%%my_blup+trKC22*Ve)/ncol(Z)
  if(!i%5){cat('It ',i, 'VC:  Vg =',Vg, 'and Ve =',Ve, '\n')}}

## It 5 VC:  Vg = 191751.1 and Ve = 441110.7
## It 10 VC:  Vg = 189728.4 and Ve = 442053.2
## It 15 VC:  Vg = 189681.5 and Ve = 442075.1
## It 20 VC:  Vg = 189680.4 and Ve = 442075.6
## It 25 VC:  Vg = 189680.4 and Ve = 442075.6
```

58/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

58/171

# Example 2 - Barley

59/171

59/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

# Example 2 - load data

Another example dataset from Kevin's *agridat* package

```
data(stepToe.morex.pheno,package='agridat')
dt = stepToe.morex.pheno
head(dt)
```

| ##   | gen     | env   | amylase | diapow | hddate | lodging | malt | height | protein | yield  |
|------|---------|-------|---------|--------|--------|---------|------|--------|---------|--------|
| ## 1 | StepToe | MN92  | 22.7    | 46     | 149.5  | NA      | 73.6 | 84.5   | 10.5    | 5.5315 |
| ## 2 | StepToe | MTi92 | 30.1    | 72     | 178.0  | 10      | 76.5 | NA     | 11.2    | 8.6403 |
| ## 3 | StepToe | MTd92 | 26.7    | 78     | 165.0  | 15      | 74.5 | 75.5   | 13.4    | 5.8990 |
| ## 4 | StepToe | ID91  | 26.2    | 74     | 179.0  | NA      | 74.1 | 111.0  | 12.1    | 8.6290 |
| ## 5 | StepToe | OR91  | 19.6    | 62     | 191.0  | NA      | 71.5 | 90.0   | 11.7    | 5.3440 |
| ## 6 | StepToe | WA91  | 23.6    | 54     | 181.0  | NA      | 73.8 | 112.0  | 10.0    | 6.2700 |

60/171

60/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

## Example 2 - Getting design matrix

- Linear model:  $Phe = Env + Gen$
- In algebra notation:  $y = Xb + Zu + e$

```
X = model.matrix(~env,dt)
Z = model.matrix(~gen-1,dt) # "-1" means no intercept
y = dt$yield
```

61/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

61/171

## Example 2 - Fit the model

```
# Fit
fit0 = reml(y=y,X=X,Z=Z)
# BLUE and BLUP
blue0 = fit0$Fixed[,1]
blup0 = fit0$EBV
# Get VC
fit0$VC[c(1:2)]

##           Vg           Ve
## 1 0.1320092 0.6379967
```

62/171

## Example 2 - DIY BLUPs

```

iK = diag(ncol(Z))
Lambda = 0.637997/0.132009
W = cbind(X,Z)
Sigma = as.matrix(Matrix::bdiag(diag(0,ncol(X)),iK*Lambda))
LHS = crossprod(W) + Sigma
RHS = crossprod(W,y)
g = solve(LHS,RHS)
my_blue = g[ c(1:ncol(X))]
my_blup = g[-c(1:ncol(X))]

```

63/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8


63/171

## Example 2 - DIY BLUPs

```

par(mfrow=c(1,2))
plot(my_blue,blue0,main="BLUE")
plot(my_blup,blup0,main="BLUP")

```



64/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

64/171



## Example 2 - Check variance components

$$\sigma_e^2 = \frac{e'y}{n-p} \text{ and } \sigma_u^2 = \frac{u'K^{-1}u + \text{tr}(K^{-1}C^{22}\sigma_e^2)}{q}$$

```
e = y - X %>% my_blue - Z %>% my_blup
Ve = c(y%*%e)/(length(y)-ncol(X))
Ve

## [1] 0.6379967

trKC22 = sum(diag(ik%*(solve(LHS)[-c(1:ncol(X)),-c(1:ncol(X))]))))
Vg = c(t(my_blup)%*%ik%*my_blup+trKC22*Ve)/ncol(Z)
Vg

## [1] 0.1320091
```

65/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

65/171

## Starting from bad variance components

```
Ve = Vg = 1
for(i in 1:25){
  Lambda = Ve/Vg;
  Sigma = as.matrix(Matrix::bdiag(diag(0,ncol(X)),ik*Lambda))
  LHS = crossprod(W) + Sigma; RHS = crossprod(W,y); g = solve(LHS,RHS)
  my_blue = g[ c(1:ncol(X))]; my_blup = g[-c(1:ncol(X))]
  e = y - X%*%my_blue - Z%*%my_blup; Ve = c(y%*%e)/(length(y)-ncol(X))
  trKC22 = sum(diag(ik%*(solve(LHS)[(ncol(X)+1):(ncol(W)),(ncol(X)+1):(ncol(W)))]))
  Vg = c(t(my_blup)%*%ik%*my_blup+trKC22*Ve)/ncol(Z)
  if(!i%5){cat('It',i,'VC:  Vg =',Vg,'and Ve =',Ve,'\n')}}

## It 5 VC:  Vg = 0.1336139 and Ve = 0.6370751
## It 10 VC:  Vg = 0.1320386 and Ve = 0.6379797
## It 15 VC:  Vg = 0.1320097 and Ve = 0.6379964
## It 20 VC:  Vg = 0.1320092 and Ve = 0.6379967
## It 25 VC:  Vg = 0.1320092 and Ve = 0.6379967
```

66/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

66/171

## Example 3 - Barley GEBV

67/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

67/171

3/18/2019

Mixed models applied to breeding

## Using genomic information!

```
data(stepToe.morex.geno, package='agridat')
gen = do.call("cbind", lapply(stepToe.morex.geno$geno, function(x) x$data))
gen = rbind(0, 2, gen)
rownames(gen) = c('Morex', 'StepToe', as.character(stepToe.morex.geno$pheno$gen))
rownames(gen)[10] = "SM8"
gen = gen[gsub('gen', '', colnames(Z)), ]
K = GRM(IMP(gen), T)
```

68/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

68/171

## Example 3 - Fit the model

```
# Fit model
fit0 = reml(y=y,X=X,Z=Z,K=K)
# BLUE and BLUP
blue0 = fit0$Fixed[,1]
gebv0 = fit0$EBV
# Get VC
fit0$VC[c(1:2)]

##          Vg          Ve
## 1 1.027988 0.6575786
```

69/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

69/171

## Example 3 - DIY BLUPs

```
diag(K) = diag(K)+1e-08; iK = solve(K)
Lambda = 0.6575/1.0280
W = cbind(X,Z)
Sigma = as.matrix(Matrix::bdiag(diag(0,ncol(X)),iK*Lambda))
LHS = crossprod(W) + Sigma
RHS = crossprod(W,y)
g = solve(LHS,RHS)
my_blue = g[ c(1:ncol(X))]
my_gebv = g[ -c(1:ncol(X))]
```


70/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

70/171

## Example 3 - DIY BLUPs

```
par(mfrow=c(1,2))
plot(my_blue,blue0,main="BLUE")
plot(my_gebv,gebv0,main="GEBV")
```



71/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

71/171

## Example 3 - Check variance components

$$\sigma_e^2 = \frac{e'y}{n-p} \text{ and } \sigma_u^2 = \frac{u'K^{-1}u + \text{tr}(K^{-1}C^{22}\sigma_e^2)}{q}$$

```
e = y - X %*% my_blue - Z %*% my_blu
Ve = c(y%*%e)/(length(y)-ncol(X))
Ve
```

```
## [1] 0.6379967
```

```
trKC22 = sum(diag(ik%*(solve(LHS)[(ncol(X)+1):(ncol(W)):(ncol(W))]))
Vg = c(t(my_blu)%*%ik%*my_blu+trKC22*Ve)/ncol(Z)
Vg
```

```
## [1] 14.12085
```

72/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

72/171

## Example 4 - Soybeans

73/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

73/171

3/18/2019

Mixed models applied to breeding

## snp-BLUP

```
data(tpod, package='NAM')  
X = matrix(1, length(y), 1)  
Z = gen  
dim(Z)
```

```
## [1] 196 376
```


74/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

74/171

## Example 4 - Fit the model

```
# Fit using the lme4 package
fit0 = reml(y=y,X=X,Z=Z) # same as reml(y=y,Z=gen)
marker_values = fit0$EBV
gebv0 = c(gen %%% marker_values)
# Marker effects
plot(marker_values,pch=16, xlab='SNP')
```



75/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

75/171

## Example 4 - DIY BLUPs

```
iK = diag(ncol(Z))
Lambda = fit0$VC[2] / fit0$VC[1]
W = cbind(X,Z)
Sigma = diag( c(0,rep(Lambda,ncol(Z))) )
LHS = crossprod(W) + Sigma
RHS = crossprod(W,y)
g = solve(LHS,RHS)
my_mu = g[ c(1:ncol(X))]
my_marker_values = g[-c(1:ncol(X))]
my_gebv = c(gen %%% my_marker_values) # GEBVs from RR
```


76/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

76/171

## Example 4 - DIY BLUPs

```
par(mfrow=c(1,2))
plot(my_marker_values, marker_values, main="Markers")
plot(my_gebv, gebv0, main="GEBV")
```



77/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

77/171

## Example 4 - Heritability from RR

```
fit0$VC
```

```
##           Vg           Ve           h2
## 1 1.659819e-05 0.03167014 0.0005238214
```

```
Scale=sum(apply(gen,2,var)); Va=fit0$VC[1]*Scale; Ve=fit0$VC[2]
round((Va/(Va+Ve)),2)
```

```
##           Vg
## 1 0.16
```

```
K = tcrossprod(apply(gen,2,function(x) x-mean(X)))
K = K/mean(diag(K)); round(rem1(y,K=K)$VC,2)
```

```
##           Vg           Ve           h2
## 1 0.01 0.03 0.16
```

78/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

78/171

# Estimate VC from bad starters

```

W = cbind(X,Z); iK = diag(ncol(Z))
Ve = Vg = 1 # Bad starting values
for(i in 1:100){ # Check the VC convergence after few iterations
  Lambda = Ve/Vg;
  Sigma = as.matrix(Matrix::bdiag(diag(0,ncol(X)),iK*Lambda))
  LHS = crossprod(W) + Sigma; RHS = crossprod(W,y); g = solve(LHS,RHS)
  my_blue = g[ c(1:ncol(X))]; my_blup = g[-c(1:ncol(X))]
  e = y - X%*%my_blue - Z%*%my_blup; Ve = c(y%*%e)/(length(y)-ncol(X))
  trKC22 = sum(diag(iK%*(solve(LHS)[(ncol(X)+1):(ncol(W)),(ncol(X)+1):(ncol(W)))]))
  Vg = c(t(my_blup)%*%iK%*my_blup+trKC22*Ve)/ncol(Z)
  if(!i%25){cat('It',i, 'VC:  Vg = ',Vg, 'and Ve = ',Ve, '\n')}}

## It 25 VC:  Vg = 0.0002960602 and Ve = 0.01801226
## It 50 VC:  Vg = 7.428217e-05 and Ve = 0.02531553
## It 75 VC:  Vg = 4.000965e-05 and Ve = 0.02818575
## It 100 VC:  Vg = 2.887763e-05 and Ve = 0.02956763

```

79/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

79/171

# Break



# Module 3 - Advanced topics

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

81/171

3/18/2019

Mixed models applied to breeding

## Outline


- Multivariate models
- Bayesian methods
- Machine learning
- G x E interactions

82/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

82/171

## Multivariate models



file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

83/171

83/171

## Multivariate models

Mixed models also enable us to evaluate multiple traits:

- More accurate parameters: BV and variance components
- Information: Inform how traits relate to each other
- Constrains: May increase computation time considerably

It preserves the same formulation

$$y = Xb + Zu + e$$

However, we now stack the traits together:

$$y = \{y_1, y_2, \dots, y_k\}, X = \{X_1, X_2, \dots, X_k\}', b = \{b_1, b_2, \dots, b_k\}, \\ Z = \{Z_1, Z_2, \dots, Z_k\}', u = \{u_1, u_2, \dots, u_k\}, e = \{e_1, e_2, \dots, e_k\}.$$

84/171

# Multivariate models

The multivariate variance looks nice at first

$$\text{Var}(y) = \text{Var}(u) + \text{Var}(e)$$

But can get ugly with a closer look:

$$\text{Var}(u) = Z(G \otimes \Sigma_a)Z' = \begin{bmatrix} Z_1 G Z_1' \sigma_{a_1}^2 & Z_1 G Z_2' \sigma_{a_{12}} \\ Z_2 G Z_1' \sigma_{a_{21}} & Z_2 G Z_2' \sigma_{a_2}^2 \end{bmatrix}$$

and

$$\text{Var}(e) = R \otimes \Sigma_e = \begin{bmatrix} R\sigma_{e_1}^2 & R\sigma_{e_1 e_2} \\ R\sigma_{e_2 e_1} & R\sigma_{e_2}^2 \end{bmatrix}$$

85/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

85/171

# Multivariate models

You can still think the multivariate mixed model as

$$y = Wg + e$$

Where

$$y = \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}, W = \begin{bmatrix} X_1 & 0 & Z_1 & 0 \\ 0 & X_2 & 0 & Z_2 \end{bmatrix}, g = \begin{bmatrix} b_1 \\ b_2 \\ u_1 \\ u_2 \end{bmatrix}, e = \begin{bmatrix} e_1 \\ e_2 \end{bmatrix}$$

86/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

86/171

# Multivariate models

Left-hand side ( $W'R^{-1}W + \Sigma$ )

$$\begin{bmatrix} X_1'X_1\Sigma_{e11}^{-1} & X_1'X_2\Sigma_{e12}^{-1} & X_1'Z_1\Sigma_{e11}^{-1} & X_1'Z_2\Sigma_{e12}^{-1} \\ X_2'X_1\Sigma_{e12}^{-1} & X_2'X_2\Sigma_{e22}^{-1} & X_2'Z_1\Sigma_{e12}^{-1} & X_2'Z_2\Sigma_{e22}^{-1} \\ Z_1'X_1\Sigma_{e11}^{-1} & Z_1'X_2\Sigma_{e12}^{-1} & G^{-1}\Sigma_{a11}^{-1} + Z_1'Z_1\Sigma_{e11}^{-1} & G^{-1}\Sigma_{a12}^{-1} + Z_1'Z_2\Sigma_{e12}^{-1} \\ Z_2'X_1\Sigma_{e12}^{-1} & Z_2'X_2\Sigma_{e22}^{-1} & G^{-1}\Sigma_{a11}^{-1} + Z_2'Z_1\Sigma_{e12}^{-1} & G^{-1}\Sigma_{a22}^{-1} + Z_2'Z_2\Sigma_{e22}^{-1} \end{bmatrix}$$

Right-hand side ( $W'R^{-1}y$ )

$$\begin{bmatrix} X_1'(y_1\Sigma_{e1}^{-1} + y_2\Sigma_{e12}^{-1}) \\ X_2'(y_1\Sigma_{e12}^{-1} + y_2\Sigma_{e2}^{-1}) \\ Z_1'(y_1\Sigma_{e1}^{-1} + y_2\Sigma_{e12}^{-1}) \\ Z_2'(y_1\Sigma_{e12}^{-1} + y_2\Sigma_{e2}^{-1}) \end{bmatrix}$$

87/171

# Multivariate models

```
data(wheat, package = 'BGLR')
G = NAM::GRM(wheat.X);
Y = wheat.Y; colnames(Y) = c('E1', 'E2', 'E3', 'E4')
mmm = NAM::reml( y = Y, K = G )
knitr::kable( round(mmm$VC$GenCor,2) )
```

|    | E1    | E2    | E3    | E4    |
|----|-------|-------|-------|-------|
| E1 | 1.00  | -0.25 | -0.22 | -0.50 |
| E2 | -0.25 | 1.00  | 0.96  | 0.55  |
| E3 | -0.22 | 0.96  | 1.00  | 0.72  |
| E4 | -0.50 | 0.55  | 0.72  | 1.00  |

88/171

# Multivariate models

mmm\$VC\$Vg

| ## |    | E1         | E2         | E3         | E4         |
|----|----|------------|------------|------------|------------|
| ## | E1 | 0.6277835  | -0.1446924 | -0.1102175 | -0.2743640 |
| ## | E2 | -0.1446924 | 0.5440731  | 0.4419945  | 0.2822577  |
| ## | E3 | -0.1102175 | 0.4419945  | 0.3919626  | 0.3130735  |
| ## | E4 | -0.2743640 | 0.2822577  | 0.3130735  | 0.4828705  |

mmm\$VC\$Ve


| ## |    | E1          | E2         | E3         | E4         |
|----|----|-------------|------------|------------|------------|
| ## | E1 | 0.53504246  | 0.08247812 | -0.1159118 | 0.06882868 |
| ## | E2 | 0.08247812  | 0.56214755 | 0.2973841  | 0.15795801 |
| ## | E3 | -0.11591175 | 0.29738408 | 0.6714234  | 0.11086214 |
| ## | E4 | 0.06882868  | 0.15795801 | 0.1108621  | 0.59405228 |

89/171

# Multivariate models

- Selection indeces, co-heritability, indirect response to selection
- Study residual and additive genetic association among traits

biplot(princomp(mmm\$VC\$GenCor,cor=T),xlim=c(-.4,.4),ylim=c(-.11,.11))



90/171

# Multivariate models

When do additional traits contribute?

# Fit E4, predict E2

```
fit_E4=bwGR::mrr(matrix(Y[,4]),wheat.X); cor(fit_E4$hat[,1],Y[,2])
```

```
## [1] 0.3988844
```

# Fit E4 and E1, E4 predict E2

```
fit_E4E1=bwGR::mrr(Y[,c(1,4)],wheat.X); cor(fit_E4E1$hat[,2],Y[,2])
```

```
## [1] 0.3931193
```

# Fit E4 and E3, E4 predict E2

```
fit_E4E3=bwGR::mrr(Y[,3:4],wheat.X); cor(fit_E4E3$hat[,2], Y[,2])
```


```
## [1] 0.5279279
```

91/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

91/171

# Bayesian methods



92/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

92/171

# Bayesian methods

The general framework on a hierarchical Bayesian model follows:

$$p(\theta|x) \propto p(x|\theta)p(\theta)$$

Where:

- Posterior probability:  $p(\theta|x)$
- Likelihood:  $p(x|\theta)$
- Prior probability:  $p(\theta)$

93/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

93/171

# Bayesian methods

For the model:

$$y = Xb + Zu + e, \quad u \sim N(0, K\sigma_a^2), \quad e \sim N(0, I\sigma_e^2)$$

- Data ( $x = \{y, X, Z, K\}$ )
- Parameters ( $\theta = \{b, u, \sigma_a^2, \sigma_e^2\}$ )

Probabilistic model:

$$p(b, u, \sigma_a^2, \sigma_e^2 | y, X, Z, K) \propto N(y, X, Z, K | b, u, \sigma_a^2, \sigma_e^2) \times \\ N(b, u | \sigma_a^2, \sigma_e^2) \times \chi^{-2}(\sigma_a^2, \sigma_e^2 | S_a, S_e, \nu_a, \nu_e)$$

94/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

94/171

# Bayesian methods

REML: the priors  $(S_a, S_e, \nu_a, \nu_e)$  are estimated from data.

Hierarchical Bayes: You provide priors. Here is how:

$$\sigma_a^2 = \frac{u' K^{-1} u + S_a \nu_a}{\chi^2(q + \nu_a)}$$

`sigma2a=(t(u)%*%iK%*%u+Sa*dfa)/rchisq(df=ncol(Z)+dfa,n=1)`

$$\sigma_e^2 = \frac{e' e + S_e \nu_e}{\chi^2(n + \nu_e)}$$

`sigma2e=(t(e)%*%e+Se*dfe)/rchisq(df=length(y)+dfe,n=1)`

95/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

95/171

# Bayesian methods

What does it mean for **you**? If your "prior knowledge" tells you that a given trait has approximately  $h^2 = 0.5$  (nothing unreasonable). In which case, half of the phenotypic variance is due to genetics, and the other half is due to error. Your prior shape is:

$$S_a = S_e = \sigma_y^2 \times 0.5$$

We usually assign small a prior degrees of freedoms. Something like four or five prior degrees of freedom. That means that assuming  $\nu_0 = 5$ , you are yielding to your model 5 data points that support heritability 0.5

$$\nu_a = \nu_e = 5$$

Example of prior influence: In a dataset with 300 data points, 1.6% of the variance components information comes from prior (5/305), and 98.4% comes from data (300/305).

96/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

96/171



# Bayesian methods

For whole-genome regression models

$$y = \mu + Ma + e, \quad a \sim N(0, I\sigma_a^2), \quad e \sim N(0, I\sigma_e^2)$$

We scale the prior genetic variance based on allele frequencies

$$S_b = \frac{\sigma_y^2 \times 0.5}{2 \sum p_j(1 - p_j)}$$

Two common settings:

- All markers, one random effect: *Bayesian ridge regression*
- Each markers as a random effect: *BayesA*


97/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

97/171

# Machine learning methods

- Parametric methods for prediction: L1-L2
- Semi-parametric methods for prediction: Kernels
- Non-parametric methods for prediction: Trees and nets



98/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

98/171

# Machine learning methods

L1-L2 machines include all mixed and Bayesian models we have seen so far. The basic framework is driven by a single (random) term model:

$$y = Xb + e$$

The univariate solution indicates how the model is solved. A model without regularization yields the least square (LS) solution. If we regularize by deflating the nominator, we get the L1 regularization (LASSO). If we regularize by inflating the denominator, we get the L2 regularization (Ridge). For any combination of both, we get a elastic-net (EN). Thus:

$$b_{LS} = \frac{x'y}{x'x}, \quad b_{Lasso} = \frac{x'y - \lambda}{x'x}, \quad b_{Ridge} = \frac{x'y}{x'x + \lambda}, \quad b_{EN} = \frac{x'y - \lambda_1}{x'x + \lambda_2}$$

Whereas the Bayesian and mixed model framework resolves the regularization as  $\lambda = \sigma_e^2 / \sigma_b^2$ , ML methods search for  $\lambda$  through ( $k$ -fold) cross-validation.

99/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

99/171

# Machine learning methods

Common loss functions in L1-L2 machines

- LS (no prior, no shrinkage):  $\operatorname{argmin}(\sum e_i^2)$
- L1 (Laplace prior with variable selection):  $\operatorname{argmin}(\sum e_i^2 + \lambda \sum |b_j|)$
- L2 (Gaussian prior, unique solution):  $\operatorname{argmin}(\sum e_i^2 + \lambda \sum b_j^2)$

Other losses that are less popular

- Least absolute:  $\operatorname{argmin}(\sum |e_i|)$  based on  $b_{LA} = \frac{nMD(x \times y)}{x'x}$
- $\epsilon$ -loss:  $\operatorname{argmin}(\sum e_i^2, |e_i| > \epsilon)$  - used in support vector machines

100/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

100/171

# Machine learning methods

Cross-validations to search for best value of lambda

```
lasso = glmnet::cv.glmnet(x=wheat.X,y=rowMeans(Y),alpha=1);
ridge = glmnet::cv.glmnet(x=wheat.X,y=rowMeans(Y),alpha=0);
par(mfrow=c(1,2)); plot(ridge); plot(lasso)
```



101/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

101/171

# Machine learning methods

Re-fit the model using this best value

```
lasso = glmnet::glmnet(x=wheat.X,y=rowMeans(Y),lambda=lasso$lambda.min,alpha=1)
ridge = glmnet::glmnet(x=wheat.X,y=rowMeans(Y),lambda=ridge$lambda.min,alpha=0)
par(mfrow=c(1,2)); plot(lasso$beta,main='LASSO'); plot(ridge$beta,main='RIDGE');
```



102/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

102/171

# Machine learning methods

Of course, the losses presented above are not limited to the application of prediction and classification. Below, we see an example of deploying LASSO for a graphical model (Markov Random Field): How the traits of the multivariate model relate in terms of additive genetics:

```
ADJ=huge: :huge(mmm$VC$GenCor, .3, method='glasso', verbose=F)$path[[1]]
plot(igraph::graph.adjacency(adjmatrix=ADJ), vertex.label.cex=3)
```



103/171

# Machine learning methods

Reproducing kernel Hilbert Spaces (RKHS), is a generalization of a GBLUP... Most commonly instead of using the linear kernel ( $Z Z' \alpha$ ), RKHS commonly uses one or more Gaussian or exponential kernels:

$$K = \exp(-\theta D^2)$$

Where  $D^2$  is the squared Euclidean distance, and  $\theta$  is a bandwidth:

- Single kernel:  $1/mean(D^2)$
- Three kernels:  $\theta=\{5/q, 1/q, 0.2/q\}$ , where  $q=quantile(D2,0.05)$

104/171

# Machine learning methods

We can use REML, PRESS (=cross-validation) or Bayesian approach to solve RKHS

```
# Make the kernel
```

```
D2 = as.matrix(dist(wheat.X)^2)
```

$$K = \exp(-D^2/\text{mean}(D^2))$$

# Below we are going to calibrate models on Env 2 and predict Env 3

```
rkhs_press = NAM::press(y=Y[,2],K=K)$hat
```

```
rkhs_rem1 = NAM::rem1(y=Y[,2],K=K)$EBV
```

```
rkhs_bgs = NAM::gibbs(y=Y[,2],iK=solve(K))$Fit.mean
```

##

\_\_\_\_\_

| 0%

| 1%

1%

2%

105/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

105/171


# Machine learning methods

```
par(mfrow=c(1,3))
```

```
plot(rkhs_press,Y[,3],main=paste('PRESS, cor =',round(cor(rkhs_press,Y[,3]),4) ))
```

```
plot(rkhs_reml,Y[,3],main=paste('REML, cor =',round(cor(rkhs_reml,Y[,3]),4) ))
```

```
plot(rkhs_bgs,Y[,3],main=paste('Bayes, cor =',round(cor(rkhs_bgs,Y[,3]),4) ))
```




106/171

# Machine learning methods

## RKHS for epistasis and variance component analysis

```
Ks = NAM::G2A_Kernels(wheat.X) # Get all sorts of linear kernels
FIT = BGLR::BGLR(rowMeans(Y), verbose=FALSE,
  ETA=list(A=list(K=Ks$A, model='RKHS'), AA=list(K=Ks$A, model='RKHS')))
pie(c(Va=FIT$ETA$A$varU, Vaa=FIT$ETA$AA$varU, Ve=FIT$varE), main='Epistasis')
```



107/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

107/171

# Machine learning methods

For the same task (E2 predict E3), let's check members of the Bayesian alphabet

```
fit_BRR = bwGR::wgr(Y[,2],wheat.X); cor(c(fit_BRR$hat),Y[,3])

## [1] 0.5842116

fit_BayesB = bwGR::BayesB(Y[,2],wheat.X); cor(fit_BayesB$hat,Y[,3])

## [1] 0.5355413

fit_emBayesA = bwGR::emBA(Y[,2],wheat.X); cor(fit_emBayesA$hat,Y[,3])

## [1] 0.6388318
```


108/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

108/171

# Machine learning methods

## Tree regression and classifiers




109/171

109/171

# Machine learning methods

```
fit_tree = party::ctree(y~.,data.frame(y=Y[,2],wheat.X)); plot(fit_tree)
```



```
cor(c(fit_tree@predict_response()),Y[,3])
```


## [1] 0.265622

110/171

110/171

# Machine learning methods

```
fit_rf = ranger::ranger(y~,data.frame(y=Y[,2],wheat.X))
plot(fit_rf$predictions,Y[,3],xlab='RF predictions from E2',ylab='Yield E3',pch=20)
```



```
cor(fit_rf$predictions,Y[,3])
```

```
## [1] 0.4056496
```

111/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

111/171

# Genotype-Environment interactions



112/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

112/171



# Genotype-Environmnet interactions

```
y=as.vector(wheat.Y); Z=wheat.X; Zge=as.matrix(Matrix::bdiag(Z,Z,Z,Z))
#
fit_g = bwGR::BayesRR(rowMeans(wheat.Y),Z)
fit_ge = bwGR::BayesRR(y,Zge)
fit_gge = bwGR::BayesRR2(y,rbind(Z,Z,Z,Z),Zge)
#
fit_g$h2

## [1] 0.4590341

fit_ge$h2

## [1] 0.6762359

fit_gge$h2


## [1] 0.6580924
```

113/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8113/171


# Genotype-Environmnet interactions

GxE design matrix: Example of 4 environments, 30 individuals, 70 SNPs



# Genotype-Environment interactions

```
GE1=matrix(fit_ge$hat,ncol=4); GE2=matrix(fit_ge$hat,ncol=4)
plot(data.frame(G=fit_g$hat,GE=rowMeans(GE1),G_and_GE=rowMeans(GE2)),main='GEBV across E')
```




115/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

115/171

# Genotype-Environment interactions

```
par(mfrow=c(1,3))
plot(fit_g$hat,rowMeans(Y),main=round(cor(fit_g$hat,rowMeans(Y)),4),xlab='G',ylab='Obs',pch=20)
plot(c(GE1),y,rowMeans(Y),main=round(cor(c(GE1),y),4),xlab='GE model',ylab='Observed',pch=20)
plot(c(GE2),y,rowMeans(Y),main=round(cor(c(GE2),y),4),xlab='G+GE model',ylab='Observed',pch=20)
```



116/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

116/171

# Break

# Module 4 - Signal detection

# Outline

- Test statistics
- Allele coding
- Power & resolution
- Linkage mapping
- LD mapping
- Structure
- Imputation
- GLM
- MLM
- WGR
- Rare-variants
- Validation studies

119/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8


119/171

3/18/2019

Mixed models applied to breeding

## Test statistics

- Testing associations are as simple as t-test and ANOVA



120/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

120/171

# Test statistics

- A more generalized framework: Likelihood test

$$LRT = L_0/L_1 = -2(\log L_1 - \log L_0)$$

For the model:

$$\begin{aligned} y &= Xb + Zu + e \\ y &\sim N(Xb, V) \end{aligned}$$

REML function is given by:

$$L(\sigma_u^2, \sigma_e^2) = -0.5(\ln|V| + \ln|X'V^{-1}X| + y'Py)$$

Where  $V = ZKZ'\sigma_u^2 + I\sigma_e^2$  and  $y'Py = y'e$

121/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

121/171

# Allele coding

Types of allele coding

1. Add. (1 df):  $\{-1,0,1\}$  or  $\{0,1,2\}$  - **Very popular** (Lines, GCA)
2. Dom. (1 df):  $\{0,1,0\}$  - **Popular** (Trees, clonals and Hybrids)
3. Jointly A+D (2 df): Popular on QTL mapping in F2s
4. Complete dominance (1 df):  $\{0,0,1\}$  or  $\{0,1,1\}$  - **Very unusual**
5. Interactions (X df): *Marker x Factor* (epistasis and GxE)

122/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

122/171

# Power and resolution

## Power

- Key: Number of individuals & allele frequency
- More DF = lower power
- Multiple testing: Bonferroni and FDR
- Tradeoff: Power vs false positives

## Resolution

- Genotyping density
- LD blocks
- Recombination

123/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

123/171

3/18/2019

Mixed models applied to breeding

# Power: Variance of X



124/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

124/171

# Beavis effect: 1000 is just OK

Xu S. Theoretical basis of the Beavis effect. Genetics. 2003 1;165(4):2259-68.

TABLE 4  
Comparisons of predicted and observed (estimated) biases in estimated QTL effects and variances from Beavis F<sub>2</sub> simulation experiments

| Simulated conditions <sup>a</sup> | Variance explained |          |                        | Additive effect |          |                        | Average estimated location |
|-----------------------------------|--------------------|----------|------------------------|-----------------|----------|------------------------|----------------------------|
|                                   | Simulated          | Observed | Predicted <sup>b</sup> | Simulated       | Observed | Predicted <sup>c</sup> |                            |
| 10-30-100                         | 3.00               | 16.76    | 16.0537                | 2.45            | 4.96     | 5.6410                 | 11.3                       |
| 10-30-500                         | 3.00               | 4.33     | 4.1890                 | 2.45            | 2.89     | 2.8617                 | 10.53                      |
| 10-30-1000                        | 3.00               | 3.02     | 3.1846                 | 2.45            | 2.56     | 2.4868                 | 10.8                       |
| 10-63-100                         | 6.25               | 12.65    | 16.5984                | 3.55            | 4.68     | 5.7328                 | 10.51                      |
| 10-63-500                         | 6.25               | 7.08     | 6.5381                 | 3.55            | 3.73     | 3.5829                 | 10.96                      |
| 10-63-1000                        | 6.25               | 6.34     | 6.3566                 | 3.55            | 3.60     | 3.5500                 | 11.04                      |
| 10-95-100                         | 9.50               | 18.68    | 17.3883                | 4.36            | 5.85     | 5.8466                 | 10.58                      |
| 10-95-500                         | 9.50               | 10.1     | 9.7082                 | 4.36            | 4.49     | 4.3607                 | 11.08                      |
| 10-95-1000                        | 9.50               | 9.67     | 9.6028                 | 4.36            | 4.44     | 4.3600                 | 11.19                      |
| 40-30-100                         | 0.75               | 15.78    | 15.6270                | 1.22            | 4.40     | 5.5436                 | 10.83                      |
| 40-30-500                         | 0.75               | 3.17     | 3.3332                 | 1.22            | 2.35     | 2.5671                 | 10.17                      |
| 40-30-1000                        | 0.75               | 1.46     | 1.7961                 | 1.22            | 1.85     | 1.8790                 | 10.17                      |
| 40-63-100                         | 1.56               | 16.31    | 15.7983                | 1.77            | 4.71     | 5.5999                 | 10.45                      |
| 40-63-500                         | 1.56               | 3.54     | 3.5783                 | 1.77            | 2.59     | 2.6582                 | 10.13                      |
| 40-63-1000                        | 1.56               | 1.96     | 2.1435                 | 1.77            | 2.09     | 2.0494                 | 10.37                      |
| 40-95-100                         | 2.40               | 16.55    | 15.9694                | 2.18            | 5.02     | 5.6236                 | 10.45                      |
| 40-95-500                         | 2.40               | 3.97     | 3.9190                 | 2.18            | 2.79     | 2.7641                 | 10.12                      |
| 40-95-1000                        | 2.40               | 2.58     | 2.6970                 | 2.18            | 2.36     | 2.2784                 | 10.29                      |

<sup>a</sup> Numerical values denote the number of QTL-heritability-number of progeny.  
<sup>b</sup> Using Equation 17.  
<sup>c</sup> Using Equation 8.

125/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

125/171

# Multiple testing:

GWAS tests *m* hypothesis:

- No correction:  $\alpha = 0.05$
- Bonferroni:  $\alpha = 0.05/m$
- FDR (25%):  $\alpha = 0.05/(m \times 0.75)$




126/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

126/171

# Linkage mapping

- Generally on experimental pops (F2, DH, RIL, BC)
- Based on single-marker analysis or interval mapping
- Recombination rates would increase **power**




127/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

127/171

# LD mapping (or association mapping)

- Use of historical LD between marker and QTL
- AM allowed studies on random panels
- Dense SNP panels would increase **resolution**



128/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

128/171




# Structure

- 1. Confounding associations with sub-populations
- 2. Major limitation of association mapping
- 3. Structure: *PCs, clusters, subpopulation* (eg. race)



129/171

# Structure



**Genes mirror geography within Europe**


nature Vol 456|6 November 2008|doi:10.1038/nature07331

130/171

# Imputation

Less missing values = more obs. = more detection power

- Markov models: Based on flanking markers
- Random forest: Multiple decision trees capture LD
- kNN & Projections: Fill with similar haplotypes



131/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

131/171

# GLM (generalized linear models)

- Full model ( $L_1$ ):

$$y = Xb + m_j a + e$$

- Null model ( $L_0$ ):

$$y = Xb + e$$

1. **Advantage:** Fast, not restricted to Gaussian traits
2. Popular methodology on human genetic studies
3.  $Xb$  includes (1) environment, (2) structure and (3) covariates

132/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

132/171

# MLM (mixed linear models)

- Full model ( $L_1$ ):

$$y = Xb + Zu + m_j a + e$$

- Null model ( $L_0$ ):

$$y = Xb + Zu + e$$

1. The famous "Q+K model"
2. **Advantage:** Better control of false positives, no need for PCs
3. Polygenic effect ( $u$ ) assumes  $u \sim N(0, K\sigma_u^2)$
4. Faster if we don't reestimate  $\lambda = \sigma_e^2 / \sigma_u^2$  for each SNP

133/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

133/171

# cMLM (compressed MLM)

1. Uses the same base model as MLM
2. **Advantage:** Faster than MLM
3. Based on clustered individuals:
  - $Z$  indicates the subgroup
  - $K$  is the relationship among subgroup
  - Often needs PCs to complement  $K$

134/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

134/171

# WGR (whole-genome regression)

- 1. Tests all markers at once
- 2. **Advantage:** No double-fitting, no PCs, no Bonferroni
- Model (BayesB, BayesC, SSVS):

$$y = Xb + Ma + e$$

- Marker effects are from a mixture of distributions


$a_j \sim \text{Binomial}$  with  $p(\pi) = 0$  and  $p(1 - \pi) = a_j$

135/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

136/171

# WGR (whole-genome regression)



136/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

136/171

# Rare variants

1. Screen a set ( $s$ ) of low MAF markers on NGS data
2. **Advantage:** Detect signals from low power SNPs
3. Applied to uncommon diseases (not seen in plant breeding)
4. Two possible model

- Full model 1 ( $L_1$ ):  $y = Xb + M_s a + e$
- Full model 2 ( $L_2$ ):  $y = Xb + PC_1(M_s) + e$
- Null model ( $L_0$ ):  $y = Xb + e$

Test either  $LR(L_1, L_0)$  or  $LR(L_2, L_0)$

137/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

137/171

# Validation studies

- QTLs detected with 3 methods, across 3 mapping pops
- Validations made on 3 unrelated populations

<https://doi.org/10.1534/g3.118.200636>



**Figure 2** Stacked plots of GWAS and QTL results. From upper to lower panels are results from the Bayesian-based multi-variant (A) stepwise regression (B) and single variant (C) models for GWAS and the joint QTL mapping result (D). The red dashed line in the QTL plot indicates the 1,000 permutation threshold and black lines show the QTL confidence intervals. Red squares in panel (A), triangles in panel (B) and circles in panel (C) indicate the kernel row number associated variants selected for further genetic validation.

138/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

138/171

# Break

# Module 5 - Association analysis

# Prelude: Data & Structure

141/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

141/171

## Getting some data

Example dataset from the *SoyNAM* package. We are querying two of the forty biparental families with a shared parental IA3023, grown in 18 environment.

```
Data = SoyNAM::BLUP(trait = 'yield', family = 2:3)

## solving BLUE of checks
## solving BLUP of phenotypes
## No redundant SNPs found
## There are 312 markers with MAF below the threshold
## Removing markers with more than 50% missing values
## Imputing with expectation (based on transition prob)
## removing repeated genotypes
## solving identity matrix
## individual 1 had 37 duplicate(s)
## individual 169 had 1 duplicate(s)
## individual 182 had 1 duplicate(s)
```

142/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

142/171

# Genomic relationship matrix

```

y = Data$Phen
M = Data$Gen
#
Z = apply(M, 2, function(snp) snp-mean(snp))
ZZ = tcrossprod(Z)
Sum2pq = sum(apply(M, 2, function(snp){p=mean(snp)/2; return(2*p*(1-p))}))
G = ZZ/Sum2pq

```

Kernel commonly deployed, referred in VanRaden (2008)

$$G = \frac{(M - P)(M - P)'}{2 \sum_{j=1}^J p_j(1 - p_j)}$$

143/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

143/171

# Genomic relationship matrix

```
image(G[, 241:1], main='GRM heatmap', xaxt='n', yaxt='n')
```

**GRM heatmap**



144/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

144/171



## Structure parameters (1) PCs

```
Spectral = eigen(G,symmetric = TRUE)
PCs = Spectral$eigenvectors[,1:5]
plot(PCs,xlab='PC 1',ylab='PC 2',main='Population on eigen spaces',col=Data$Fam,pch=20)
```



145/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

145/171

## Structure parameters (2) Clusters

```
GeneticDistance = Gdist(M,method=6)

## Modified Rogers' distance

Tree = hclust(GeneticDistance,method = 'ward.D2')
Clst = factor(cutree(Tree,k=2))
plot(Tree,labels = FALSE)
```

146/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

146/171

# Single marker analysis

147/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

147/171

3/18/2019

Mixed models applied to breeding

## GLM (1) - No structure

```

Marker = M[,117]
#
fit = lm( y ~ Marker )
anova( fit )

## Analysis of Variance Table
##
## Response: y
##           Df  Sum Sq Mean Sq F value    Pr(>F)
## Marker      1  476321  476321   20.172 1.102e-05 ***
## Residuals 239 5643504    23613
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

-log(anova(fit)$`Pr(>F)`[1],base = 10)

## [1] 4.957736

```

148/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

148/171

## GLM (2) - Principal Components

```

reduced_model = lm( y ~ PCs )
full_model = lm( y ~ PCs + Marker )
anova( reduced_model, full_model )

## Analysis of Variance Table
##
## Model 1: y ~ PCs
## Model 2: y ~ PCs + Marker
##   Res.Df    RSS Df Sum of Sq   F    Pr(>F)
## 1      235 4060362
## 2      234 3562067  1    498295 32.734 3.215e-08 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

-log((anova( reduced_model, full_model ))$`Pr(>F)`[2],base = 10)

## [1] 7.492813

```

149/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

149/171

## GLM (3) - Population Clusters

```

reduced_model = lm( y ~ Clst )
full_model = lm( y ~ Clst + Marker )
anova( reduced_model, full_model )

## Analysis of Variance Table
##
## Model 1: y ~ Clst
## Model 2: y ~ Clst + Marker
##   Res.Df    RSS Df Sum of Sq   F    Pr(>F)
## 1      239 4275698
## 2      238 3652041  1    623657 40.643 9.4e-10 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

-log( anova(reduced_model,full_model)$`Pr(>F)`[2],base = 10)

## [1] 9.026884

```

150/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

150/171

# MLM - K+Q model

```
Q = model.matrix(~C1st)
reduced_model = reml( y=y, X=Q, K=G)
full_model = reml( y=y, X=cbind(Q, Marker), K=G)
LRT = -2*(full_model$loglik - reduced_model$loglik)
-log(pchisq(LRT,1,lower.tail=FALSE),base=10)

## [1] 10.80903
```

151/171

[file:///C:/Users/rd7564/Desktop/Purdue\\_MLM/MyLecture/Purdue\\_MLM.html#8](file:///C:/Users/rd7564/Desktop/Purdue_MLM/MyLecture/Purdue_MLM.html#8)

151/171

# Whole genome screening


152/171

[file:///C:/Users/rd7564/Desktop/Purdue\\_MLM/MyLecture/Purdue\\_MLM.html#8](file:///C:/Users/rd7564/Desktop/Purdue_MLM/MyLecture/Purdue_MLM.html#8)

152/171

## DIY (example with GLM)

```
reduced_model = lm( y ~ C1st )
glm_pval = apply(M,2,function(Marker){
  pval = anova(reduced_model, lm(y~C1st+Marker) )$`Pr(>F)`[2]
  return(-log(pval,base = 10)))
plot(glm_pval,pch=20,xlab="SNP",main="My first GLM GWAS")
```



153/171

file:///C:/Users/rd7564/Desktop/Purdue MLM/MyLecture/Purdue MLM.html#8

153/171

## Using CRAN implementations

NAM random model:  $y = \mu + Marker \times Pop + Zu + e$

```
fit_gwa = gwas3(y=y, gen=M, fam=c(C1st), chr=Data$Chrom)
```

```
## Calculating G matrix
## Solving polygenic model
## Starting Eigendecomposition
## Starting Marker Analysis
##
```

|    |  |    |
|----|--|----|
|    |  | 0% |
|    |  |    |
|    |  | 1% |
|    |  |    |
| =  |  | 1% |
|    |  |    |
| =  |  | 2% |
|    |  |    |
| == |  | 2% |


154/171

file:///C:/Users/rd7564/Desktop/Purdue MLM/MyLecture/Purdue MLM.html#8

154/171

# Manhattan plot

```
plot(fit_gwa, pch=20, main = "My first MLM GWAS")
```



155/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

155/171

# Yet another R implementations

```
require(rrBLUP,quietly = TRUE); COL = fit_gwa$MAP[,1]%2+1 # Color chromosomes
geno=data.frame(colnames(M),fit_gwa$MAP[,1:2],t(M-1),row.names=NULL)
pheno=data.frame(line=colnames(geno)[-c(1:3)],Pheno=y,C1st,row.names=NULL)
fit_another_gwa=GWAS(pheno,geno,fixed='C1st',plot=FALSE)
```

```
## [1] "GWAS for trait: Pheno"
## [1] "Variance components estimated. Testing markers."
```




156/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

156/171

# Comparing results


```
m1m_pval=fit_another_gwa$Pheno; m1m_pval[m1m_pval==0]=NA
plot(glm_pval,m1m_pval,xlab='GLM',ylab='MLM',main='Compare')
```



157/171

# Power analysis - QQ plot

```
nam_pval = fit_gwa$PolyTest$pval
par(mfrow=c(1,3))
qqman::qq(glm_pval,main='GLM')
qqman::qq(m1m_pval,main='MLM')
qqman::qq(nam_pval,main='RLM')
```



158/171

# Multiple testing

159/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

159/171

# Multiple testing

*Wikipedia:* In statistics, the **multiple comparisons, multiplicity or multiple testing** problem occurs when one considers a set of statistical inferences simultaneously or infers a subset of parameters selected based on the observed values. In certain fields it is known as the look-elsewhere effect: *The more inferences are made, the more likely erroneous inferences are to occur.* Several statistical techniques have been developed to prevent this from happening, allowing significance levels for single and multiple comparisons to be directly compared. These techniques generally require a **stricter significance threshold** for individual comparisons, so as to compensate for the number of inferences being made.

160/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8


160/171



# Baseline - No correction

Base significance threshold:  $\alpha = 0.05$

```
plot(fit_gwa, alpha=0.05, main = "No correction")
```



161/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

161/171

# Multiple testing correction

Bonferroni:  $\alpha = 0.05/m$

```
plot(fit_gwa, alpha=0.05/ncol(M), main = "Bonferroni correction")
```



162/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

162/171

# False-Discovery Rate

$$\text{Benjamini-Hochberg FDR: } \alpha = \frac{0.05}{m \times (1 - \text{FDR})}$$

```
plot(fit_gwa, alpha=0.05/(ncol(M)*.75), main = "FDR 25%")
```



163/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

163/171

# False-Discovery Rate

Unique segments based on Eigenvalues:  $m^* = D > 1$

```
m_star = sum(Spectral$values>1)
plot(fit_gwa, alpha=0.05/m_star, main="Bonferroni on unique segments")
```



164/171


file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

164/171

# Multi-loci analysis


# Whole genome regression

```
fit_wgr = bwGR::BayesDpi(y=y,X=M,it=3000); par(mfrow=c(1,2));  
plot(fit_wgr$PVAL,col=COL,pch=20,ylab=' -log(p) ',main='GWA')  
plot(fit_wgr$b,col=COL,pch=20,ylab='Marker effect',main='GWP')
```



# WGR - No need for multiple testing

```
thr_none = -log(pchisq(qchisq(1-0.05/ncol(M),1),1,lower.tail=FALSE),base=10)
thr_bonf = -log(pchisq(qchisq(1-0.05,1),1,lower.tail=FALSE),base=10)
par(mfrow=c(1,2)); plot(fit_gwa,alpha=NULL,main="MLM",pch=20); abline(h=thr_none,col=3)
plot(fit_wgr$PVAL,col=COL,ylab=' -log(p)',main="WGR",pch=20); abline(h=thr_bonf,col=3)
```




167/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

167/171

# Approaches are complementary




168/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

168/171

# Random forest

```
fit_rf = ranger::ranger(y~,data= data.frame(y=y,M),importance='impurity')
plot(fit_rf$variable.importance,ylab='Importance',main='Random Forest',col=COL+7,pch=20)
```



169/171

file:///C:/Users/rd7564/Desktop/Purdue\_MLM/MyLecture/Purdue\_MLM.html#8

169/171

# Thanks!

Thanks!

- e-mail: [xaviera@purdue.edu](mailto:xaviera@purdue.edu) or [alencar.xavier@corteva.com](mailto:alencar.xavier@corteva.com)
- other resources: <https://alensex.wixsite.com/home>
- material: [https://github.com/alensex/Lectures/tree/master/Purdue\\_MLM](https://github.com/alensex/Lectures/tree/master/Purdue_MLM)

170/171

# Break