

HW1

Sandra Villamar and Shobhit Dronamraju

2023-01-19

```
library(ggplot2)
```


1. Write a function `confBand(x, y, conf=0.95)` taking in a predictor vector x and a response vector y and return a plot with (x, y) points, the least squares line, and the confidence band at level $conf$. Apply the function to `hp` and `mpg` from `04cars` dataset.

```
confBand <- function(x, y, conf=0.95){  
  
  # compute confidence band lines  
  fit = lm(y ~ x)  
  pred_interval = predict(fit, data.frame(x), interval='confidence')  
  lower = pred_interval[,2]  
  upper = pred_interval[,3]  
  
  # plot  
  plot(x, y, pch=16) # points  
  abline(fit$coefficients[1], fit$coefficients[2], col='blue') # reg line  
  points(x, lower, col='red', type='l') # lower band  
  points(x, upper, col='red', type='l') # upper band  
}
```

```
load("HW1/04cars.rda") # load cars dataset to "dat"  
dat = dat[,c(13,15,16,18,19)] # extract selected variables  
dat = dat[complete.cases(dat),] # extract complete cases  
names(dat) = c("hp", "mpg", "wt", "len", "wd") # abbreviate names  
str(dat)
```

```
## 'data.frame': 387 obs. of 5 variables:  
## $ hp : num 103 103 140 140 140 132 132 130 110 130 ...  
## $ mpg: num 34 34 37 37 37 36 36 33 36 33 ...  
## $ wt : num 2370 2348 2617 2676 2617 ...  
## $ len: num 167 153 183 183 183 174 174 168 168 168 ...  
## $ wd : num 66 66 69 68 69 67 67 67 67 67 ...
```

```
confBand(dat$hp, dat$mpg) # apply function
```



2. Let $n = 100$. Draw n i.i.d samples from $Unif(0, 1)$ which stay fixed in what follows. Repeat the following experiment $N=1000$ times.

- Generate $y_i = 1 + x_i + \epsilon_i$ with ϵ_i i.i.d. $N(0, 0.2)$
- Compute the 99% confidence band and record whether or not it contains the true line or not

Summarize the result of this numerical experiment by returning the proportion of times (out of N) that the confidence band contained the true line.

```
n = 100
N = 1000
x = runif(n, 0, 1)
y_true = 1 + x
p = 1
count = 0

for(i in 1:N){

  # setting up formula
  e = rnorm(100, 0, sqrt(0.2))
  y = 1 + x + e

  # computing variables for confidence band
  fit = lm(y~x)
  pred = predict(fit, data.frame(x), se.fit=TRUE)
```

```

y_hat = pred$fit
se_hat = pred$se.fit

# compute confidence band
interval = sqrt((p + 1) * qf(0.99, p + 1, n - p - 1)) * se_hat
lower = y_hat - interval
upper = y_hat + interval

# check if true line within band
if(all(lower <= y_true) && all(y_true <= upper)){
  count = count + 1
}
}

proportion = count/N

```

The experiment produced the proportion: 0.993

Contributions:

We worked on all parts of this assignment together.