
Algorithm 1 Sampling of the locations of the points given the set I of active Bernoulli random variables

Input: R, I

Output: $(X_i)_{i \in I}$

Let $\varphi_I^R(x) = (\varphi_i^R(x), i \in I)$

Draw X_1 from the distribution with density $\|\varphi_I(x)\|_{C^{|I|}}^2/|I|$

$e_1 \leftarrow \frac{\varphi_I^R(X_1)}{\|\varphi_I^R(X_1)\|_{C^{|I|}}}$

for $i \leftarrow 2$ to $|I|$ **do**

Draw X_i from the distribution with density

$$p_i(x) = \frac{1}{|I| - i + 1} \left(\|\varphi_I^R(x)\|_{C^{|I|}}^2 - \sum_{k=1}^{i-1} |\langle e_k, \varphi_I^R(x) \rangle|^2 \right)$$

$$u_i \leftarrow \varphi_I^R(X_i) - \sum_{k=1}^{i-1} \langle e_k, \varphi_I^R(X_i) \rangle e_k$$

$$e_i \leftarrow \frac{u_i}{\|u_i\|_{C^{|I|}}}$$

end for

where the eigenvalues are

$$\lambda_k^R = R^{2k+2},$$

and the eigenfunctions

$$\phi_k^R : x \mapsto \sqrt{\frac{k+1}{\pi}} \frac{1}{R^{k+1}} x^k.$$

Proof.

We have, for the (original) Bergman kernel

$$k(x, y) = \frac{1}{\pi} \frac{1}{(1 - x\bar{y})^2},$$

using $\frac{1}{(1-u)^2} = \sum_{k \geq 0} (k+1)u^k$ for all $u \in \mathcal{B}(0, R)$, we have

$$\begin{aligned} \sum_{k \geq 0} \frac{k+1}{\pi} x^k \bar{y}^k &= \sum_{k \geq 0} R^{2k+2} \frac{k+1}{\pi} \frac{x^k}{R^{k+1}} \frac{\bar{y}^k}{R^{k+1}} \\ &= \sum_{k \geq 0} \lambda_n^R \phi_n^R(x) \overline{\phi_n^R(y)}, \end{aligned}$$

taking ϕ_n^R and λ_n^R as in the theorem statement. Because we are restricting onto $L^2(\mathcal{B}(0, R), \lambda)$, we ought to consider the inner product on this space and not $L^2(\mathcal{B}(0, 1), \lambda)$. We have

$$\langle \phi_n^R, \phi_m^R \rangle_{L^2(\mathcal{B}(0, R))} = \frac{\sqrt{(n+1)(m+1)}}{\pi} \frac{1}{R^{n+m+2}} \int_{\mathcal{B}(0, R)} z^n \bar{z}^m dz.$$

However,

$$\int_{\mathcal{B}(0, R)} z^n \bar{z}^m dz = \int_0^R \int_0^{2\pi} r^{n+m} e^{i\theta(n-m)} d\theta dr = 2\pi \delta_{n,m} \int_0^R r^{n+m+1} dr = 2\pi \delta_{n,m} \frac{R^{2n+2}}{2n+2}.$$