Typhoid Fever

Tuesday — الثلاثاء

6 July — ٦ يوليه

* Typhoid fever endemic in Egypt

* Transmission
poor sanitation
lack of clean water = fecal oral route

* Risk factors :-
Disposal of sewage in water
الكلام دة قل دلوقتي الا في الناس اللي بتشرب مية آبار عشان ممكن تكون مختلطة بالصرف الصحي

* Pathogenesis
4) Invade reticuloendothelial system

* Week 2 :- Relative bradycardia
↑HR المفروض لما بيحصل fever الجسم
but this doesn't occur in typhoid fever

* Week 3 :- neurological complication (confusion)



* Stepladder fever in week 1

* Bradycardia = Faget's sign

* RLQ pain due to affecting payer's patch of illium.

Complications

Usually in 3weeks if untreated

DIC = Dissiminated intravascular coagulation

بيحصل ايه ؟

coagulation → thrombosis → تفك

صغيرة

متفكش → كبيرة

مشكلة الموضوع لما يحصل مرة واحدة في كذا مكان العيان يبقى عنده مشكليتين :-

1) Intravascular coagulation (Thrombosis)

2) Bleeding why?

لان لما بتحصل مرة واحدة الجسم مبيلحقش يصنع coagulation factors بدل اللي استخدموها في عمل الـ thrombus

JULY

SAT	SUN	MON	TUE	WED	THUR	FRI
					1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24/31	25	26	27	28	29	30

CamScanner

Sunday الأحد

4 July ١٤ محرم ٢٧ يونيه ٤ يوليه

اعالج الـ bleeding يحصل و thrombosis العكس .. عشان كدة بيتسمى Death is comming (DIC)

TTT :- treatment of the cause

INR يعلى , Hb ↓↓

Question :-

Melena occur in which week?

Investigations

1) Clinically :-

step ladder fever, RLQ abdominal pain, with alternating constipation and diarrhea

" Most common presentation"

*In Egypt we rarely see pts in week 3 due to antibiotic abuse

Antibiotic بياخد و نفسه العيان مبيبقش

ممكن يظبط و يخف

2) Blood culture done in week 1 during bacteremia.



3) Widal test الى بيتعمل فحو ال
Low specifity as if pt is having old infection, Ab is still present.


O → recent infection
H → old infection

Titre of O in non-endemic countries :-
1/160 → Typhoid fever
حد مذبش قبل كدة infection
" " in endemic countries as Egypt :-
1/320 → Typhoid fever
or having Raising Titre
* حد جالى typhoid fever مبيشخبش 1/160

4) CBC ; leucopenia
elevated liver enzymes are non specific
not present in all pts

* Cefexime 3rd generation cephalosporin

* Severe typhoid as typhoid state



CamScanner

* MDR = Multiple drug Resistance
Resistant to at least one antibiotic in 3 different groups.
(one or more drug)

* XDR = Extended drug resistance

Resistant to all classes except one drug or more in one or two groups.

* PDR = Pan drug resistance

Resistant to all classes of Antibiotics.

* There are 6 Antibiotic groups.

Examples :-
pt resistant to penicillins only → Not MDR.
" " penicillin, cephalosporin → not MDR

يوليه

السبت	الأحد	الاثنين	الثلاثاء	الأربعاء	الخميس	الجمعة
					١	٢
٣	٤	٥	٦	٧	٨	٩
١٠	١١	١٢	١٣	١٤	١٥	١٦
١٧	١٨	١٩	٢٠	٢١	٢٢	٢٣
٢٤	٢٥	٢٦	٢٧	٢٨	٢٩	٣٠

* مبيحصلش ان نعمل culture قبل ما ادي antibiotic

III of Carriers
(having the infection but without symptoms)

1) Quinolone oral for 2weeks - 1 Month
or
2) Vaccine not commonly given but given to pts travelling from endemic areas to non-endemic ones

. Ty2 la not given in immunocompromised
. Conjugate given from age 1 year

JULY

SAT	SUN	MON	TUE	WED	THUR	FR
					1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24/31	25	26	27	28	29	30

CamScanner

Notes

Brucellosis

مذكرات

* Zoonotic : from animal to man .

* V.Imp Rare mode of transmission.

* LPS : week endotoxin cause continous low stimmulation of immunity يحصلها ارهاق

* Virulence factors التلاتة مع بعض اسمهم
Stealth mechanism

* Undulant fever كام يوم كويس وكام يوم تعبان

* Night sweats
(Drenching sweats) more than excessive من غير مجهود

* Sacroilitis cause lowerback pain

Diagnosis

1) Clinically :- fever, night sweats, direct contact with animals, drinking unpasterized milk

2) Culture is a biohazard for lab workers (due to risk of inhaling aerosols) so we don't start with it.

3) We start with serology :-

- Rose Bengal test = Malta test
- ELISA most accurate

4) PCR expensive

TTT :- Any intracellular organism treated by combination

- Rifampin can be given also as 300 mg BID

- Cotrimoxazole cause Jaundice to both child and mother if given in last 3mo of pregnancy.

CamScanner

Wednesday الأربعاء

30 June ٣٠ يونيه

١٠ محرم ٢٣ يونيه

- pregnant woman :-

Rifampin + Aminoglycoside (streptomycin / Gentamycin)

For 2 weeks

- Endocarditis , ttt given for 3-6 months

* Diagnosis

Serology then → PCR then → Culture

يونيه

السبت	الأحد	الاثنين	الثلاثاء	الأربعاء	الخميس	الجمعة
			١	٢	٣	٤
٥	٦	٧	٨	٩	١٠	١١
١٢	١٣	١٤	١٥	١٦	١٧	١٨
١٩	٢٠	٢١	٢٢	٢٣	٢٤	٢٥
٢٦	٢٧	٢٨	٢٩	٣٠		