Department of pharmacognosy

phytotherapy exam

Name :..... Date : .....

Q1- answer the following questions by select (A) for true and select (B) for false one.

1. Ginkgo biloba is used mainly to prevent or reduce memory deterioration during old age.
2. St. John's Wort is used mainly to treat low mood, depression, seasonal affective disorder, anxiety and insomnia.
3. St. John's Wort contains naphthodianthrones as the main active constituent.
4. Kava kava can be taken with substances acting on the central nervous system, such as alcohol, barbiturates and psychopharmacological agents.
5. Valerian is used as a mild sedative to treat symptoms of anxiety, stress and insomnia, including (especially) during menopause.
6. Valerian may inhibit the effects of anesthetics that act on GABA receptors.
7. Squill is used for the relief of chesty coughs, as an expectorant.
8. Squill should be avoided by patients with Impaired hepatic or renal function.
9. Garlic supplements may interact with antiplatelet drugs, warfarin and related drugs, increasing the risk of bleeding.

10. Garlic is also used to relieve the symptoms of catarrh, rhinitis and nasal congestion.
11. Gentian is used to treat loss of appetite and gastrointestinal complaints such as dyspepsia, bloating and flatulence.
12. Centaury has been used to treat temporary loss of appetite and as a liver and kidney tonic.
13. The major constituent of peppermint leaves is the essential oil, which contains the monoterpenes menthol and menthone.
14. Chamomile is widely used both internally and externally to treat inflammatory conditions.
15. Liquorice has long been used for the relief of digestive symptoms such as dyspepsia and to aid the healing of gastric ulcers.
16. The antitussive and expectorant properties of liquorice have been attributed to the increase in tracheal mucus secretion caused by glycyrrhizin.
17. Milk Thistle used to relieve the symptoms associated with over indulgence of food and drink, including indigestion and upset stomach.
18. dandelion is used as a diuretic and adjuvant in major urinary complaints.
19. uva ursi is used for urinary tract infections (UTIs), inflammatory conditions of the efferent urinary tract, benign prostatic hyperplasia (BPH), and bronchitis.
20. Saw palmetto extracts are used to relieve lower urinary tract symptoms such as dysuria, polyuria and urine retention, in men with a confirmed diagnosis of benign prostate hyperplasia (BPH).
21. Pumpkin seed extract is used for the symptomatic treatment of difficulties with urination in men due to benign prostatic hyperplasia (BPH).

22. Elderberry and elderflower extracts have a long history of use for the relief of the symptoms of the common cold and flu, chills and sore throats.

23. The main active compounds of Ipecacuanha are the volatile oils emetine, cephaeline, and emetamine.

24. Anethole is the major component of the fennel oil, with more than 10% estragole and more than 7.5% fenchone.

25. Eucalyptus oil has a good empirical evidence base for use as a decongestant in the symptomatic relief of cold and coughs, and as an insect repellant.

26. Evening primrose oil is taken to provide essential fatty acids, and for the symptomatic relief of itching in acute and chronic dry skin conditions.

27. Calendula extract-containing creams and gels are used to treat skin irritation, inflammation and burns, especially after radiotherapy.

28. Arnica is often taken orally as a homeopathic preparation to treat or prevent bruising, including after surgery.

29. Devil's claw extracts are used to relieve rheumatic or muscular pain, general aches and pains in the muscles and joints and backache.

30. Ginseng is used as a tonic and 'adaptogen' (i.e. to increase the body's ability to adapt to stress), to overcome fatigue, improve cognitive function and combat ageing.

31. Ginseng should be taking with MAOIs, nifedipine and in cancer chemotherapy.

32. Siberian ginseng contains triterpene saponins, which are chemically distinct from American and Asian ginseng, and known as eleutherosides.

33. The main active constituents of the ashwagandha root are the steroidal lactones, the withanolides, which include withaferin A, and sitoindosides IX and X (glycowithanolides).

34. Ashwagandha is considered to be an adaptogen in herbal medicine and a Rasayana in Ayurveda.

35. Astragalus is used widely in traditional Chinese medicine (TCM) for similar indications, as a tonic to strengthen the immune system, and for viral infections, fatigue and loss of blood.

36. Echinacea is used to treat common cold and influenza type infections and other upper respiratory tract infections.

37. Echinacea is used to treat Inflammation of the mouth and throat and major skin conditions

38. Arnica may cause contact dermatitis. It is an irritant to mucous membranes and may be fatal if taken internally.

39. Evening primrose oil may potentially interact with antiplatelet drugs such as warfarin.

40. Agents that improve the resistance to physical and psychological stress called Adaptogens.

1	2	3	4	5	6	7	8	9	10
11	12	13	14	15	16	17	18	19	20
21	22	23	24	25	26	27	28	29	30
31	32	33	34	35	36	37	38	39	40

	A	A	A	B	A	A	A	A	B
A	A	A	A	A	A	B	A	A	A
21	22	23	24	25	26	27	28	29	30
A	A	B	B	A	A	A	A	A	A
31	32	33	34	35	36	37	38	39	40
A	A	A	A	A	A	B	A	A	A
41	42	43	44	45	46	47	48	49	50
1	2	3	4	5	6	7	8	9	10
A	A	A	B	A	A	A	A	A	A
11	12	13	14	15	16	17	18	19	20