

Published on 10.04.15 in Vol 17, No 4 (2015): April

This paper is in the following e-collection/theme issue:

[Web-based and Mobile Health Interventions](#)

[Article](#) Cited By (0) Tweetations (147) Metrics

Original Paper

Long-Term Outcomes of a Web-Based Diabetes Prevention Program: 2-Year Results of a Single-Arm Longitudinal Study

S Cameron Sepah^{1,2}, PhD ; Luohua Jiang³, PhD ; Anne L Peters⁴, MD

¹University of California, San Francisco, Department of Psychiatry, San Francisco, CA, United States

²Omada Health, San Francisco, CA, United States

³University of California, Irvine, Department of Epidemiology, Irvine, CA, United States

⁴University of Southern California, Keck School of Medicine, Los Angeles, CA, United States

Corresponding Author

ABSTRACT

Background: Digital therapeutics are evidence-based behavioral treatments delivered online that can increase accessibility and effectiveness of health care. However, few studies have examined long-term clinical outcomes of digital therapeutics.

Objective: The objective of this study was to conduct a 2-year follow-up on participants in the Internet-based *Prevent* diabetes prevention program pilot study, specifically examining the effects on body weight and A1c, which are risk factors for diabetes development.

Methods: A quasi-experimental research design was used, including a single-arm pre- and post-intervention assessment of outcomes. Participants underwent a 16-week weight loss intervention and an ongoing weight maintenance intervention. As part of the program, participants received a wireless scale, which was used to collect body weight data on an ongoing basis. Participants also received A1c test kits at baseline, 0.5 year, 1 year, and 2-year time points.

Results: Participants previously diagnosed with prediabetes (n=220) were originally enrolled in the pilot study. A subset of participants (n=187) met Centers for Disease Control and Prevention (CDC) criteria for starting the program (starters), and a further subset (n=155) met CDC criteria for completing the program (completers) and were both included in analyses. Program starters lost an average of 4.7% (SD 0.4) of baseline body weight after 1 year and 4.2% (SD 0.8) after 2 years, and reduced A1c by mean 0.38% (SD 0.07) after 1 year and 0.43% (SD 0.08) after 2 years. Program completers lost mean 4.9% (SD 0.5) of baseline body weight after 1 year and 4.3% (SD 0.8) after 2 years, and reduced A1c by 0.40% (SD 0.07) after 1 year and 0.46% (SD 0.08) after 2 years. For both groups, neither 2-year weight loss nor A1c results were significantly different from 1-year results.

Conclusions: Users of the *Prevent* program experienced significant reductions in body weight and A1c that are maintained after 2 years. Contrary to the expected progression from prediabetes

**J Med Internet Res 2015;17(4):e92)**

doi:10.2196/jmir.4052

KEYWORDS**prediabetes; type 2 diabetes; obesity; diabetes prevention program; internet; online; mobile apps; mhealth; digital health; intervention****We also recommend**

Effects of Internet behavioral counseling on weight loss in adults at risk for type 2 diabetes: a randomized trial.

Sponsored. Deborah F Tate et al., J Am Med, 2003

Nutritional Education Through Internet-Delivered Menu Plans Among Adults With Type 2 Diabetes Mellitus: Pilot Study

Abeer Bader et al., JMIR Res Protoc, 2013

Diabetes in midlife and cognitive change over 20 years: a cohort study.

Sponsored. Andreea M Rawlings et al., Ann Intern Med, 2014

A Fully Automated Diabetes Prevention Program, Alive-PD: Program Design and Randomized Controlled Trial Protocol

Clifford H Block et al., JMIR Res Protoc, 2015

Association of an intensive lifestyle intervention with remission of type 2 diabetes.

Sponsored. Edward W Gregg et al., J Am Med, 2012

Effects of a Web-Based Intervention on Physical Activity and Metabolism in Older Adults: Randomized Controlled Trial

Carolien A Wijsman et al., J Med Internet Res, 2013

Feasibility, acceptability and outcomes at a 12-month follow-up of a novel community-based intervention to prevent type 2 diabetes in adults at high risk: mixed methods pilot study.

Sponsored. Linda Penn et al., BMJ Open, 2013

A Smart Web Aid for Preventing Diabetes in Rural China: Preliminary Findings and Lessons

Penglai Chen et al., J Med Internet Res, 2014

Aerobic neuromuscular electrical stimulation--an emerging technology to improve haemoglobin A1c in type 2 diabetes mellitus: results of a pilot study.

Sponsored. Louis Crowe et al., BMJ Open, 2012

Women Veterans' Experience With a Web-Based Diabetes Prevention Program: A Qualitative Study to Inform Future Practice

Tannaz Moin et al., J Med Internet Res, 2015

Powered by

Introduction

Prediabetes, the clinical precursor to type 2 diabetes, continues to a

Sections



interventions such as the Diabetes Prevention Program (DPP) [3]. As a result, there is an urgent need for effective chronic disease prevention interventions that can be disseminated to millions of people. To address such needs, a new field of “digital therapeutics” has emerged —evidence-based treatments from the field of behavioral medicine that are delivered online [4].

One of the most established digital therapeutics is *Prevent*, an Internet-based version of the DPP clinical trial. Though the DPP was primarily an individual treatment, most DPP translations including *Prevent* have successfully used group-based approaches to minimize cost, and thus group-based interventions have been recommended in evidence-based guidelines for diabetes prevention [5].

Previous published research showed that *Prevent* program starters were able to achieve clinically significant reductions in body weight and A1c after 1 year [6]. Of note, this was the first study to show that a scalable Internet-based intervention met all efficacy benchmarks for DPP programs set by CDC’s Diabetes Prevention and Recognition Program (DPRP) standards [7]. This addressed a key concern raised by Yudkin & Montori that “rolling out intensive lifestyle interventions like these to populations with pre-diabetes...would be challenging” [8].

Furthermore, a critique of DPP studies by Kahn and Davidson noted that weight loss results of DPP translations have often been low (a meta-analysis of 22 studies showing 2.4% weight loss at 1 year) [9] and often show significant weight regain after the intervention ends. However, they concede that real-world DPPs “may have promise if the weight loss achieved at year 2 can be sustained and replicated” [10].

Thus, the current study seeks to make an original contribution to the scientific literature by addressing such points through investigating the long-term outcomes and sustainability of an Internet-based DPP.

Methods

Research Design

A quasi-experimental research design was used, including a single-arm pre- and post-intervention assessment of body weight, A1c, and program engagement outcomes [6]. Participants were not compensated for participation but were enrolled in the program at no cost. Institutional Review Board (IRB) exemption was granted by Western IRB for secondary analyses of previously collected and de-identified data.

Participants

Patients were recruited via craigslist advertisements seeking participants for an Internet-based diabetes prevention program. Patients were screened for a self-reported clinical diagnosis of prediabetes within the past year and meeting CDC DPRP eligibility criteria: 18 years of age or older, have a body mass index (BMI) of ≥ 24 kg/m² (≥ 22 kg/m² if Asian).

Sections





Program

A full description of the program components has been previously published [6]. Briefly, *Prevent* is an Internet-based translation of the DPP lifestyle intervention, which includes small group support, personalized health coaching, a weekly DPP-based curriculum, and digital tracking tools.

Participants were demographically matched into groups of 10-15 participants and placed into a private online social network resembling Facebook where they could discuss goal progress and provide social support to one another. At any convenient time or place using Internet-enabled devices, they could asynchronously complete weekly DPP-based health education lessons, privately message and call a health coach for individual counseling, track weight loss, and physical activity using a wireless weight scale and pedometer, and monitor their engagement and weight loss progress.

Prevent starts with a 16-week “Core” curriculum focusing on weight loss and continues with a 36-week post-core “Sustain” curriculum focusing on weight maintenance, with this “active” intervention totaling 12 months. During this time, participants also engage with health coaching, small group discussion, and tracking of body weight/food/physical activity. After 12 months, participants continue with an ongoing intervention that is more proactive, in which they continue to have access to past curriculum, a larger *Prevent* “alumni” group discussion forum, and tracking capabilities.

Measures

Baseline demographic and health information were collected prior to program start using an Internet-based questionnaire self-completed by the subject. Body weight, the study’s primary outcome, was measured serially in pounds using a validated, wireless-enabled scale that was mailed to participants [12]. Weights were transmitted over a 2G cellular network to a central study server. Participants were encouraged to weigh themselves daily and reminded via email/telephone calls to obtain weights at the baseline, 6, 12, and 24-month assessment timepoints. Weight measurements were highly stable and the scale’s coefficient of variation was ± 0.2 lb.

A1c was measured in percent (NGSP/DCCT units) using self-administered AccuBase A1c test kits by DTI laboratories, an FDA-cleared whole blood test that uses a capillary tube blood collection method. This allows for reliable home-based data collection and valid lab testing using high-performance liquid chromatography (HPLC-IE/HPLC-BA), including abnormal hemoglobin screening [13]. A1c test kits were mailed to participants’ homes around the baseline, 6, 12, and 24-month assessment timepoints.

Analyses

Results were analyzed for two subgroups based on CDC DPRP standards: “program starters” were those who completed at least 4 core lessons, and “program completers” were those who completed at least 9 core lessons.





change point after the last core lesson day included in the model. For A1c, an additional change point at 12 months was added to the mixed model due to significant change in slope at that timepoint and better model fit statistics.

Models used an autoregressive-moving-average covariance structure to statistically account for the correlation of frequently measured weight data. Repeated A1c measures were also correlated but not measured on a daily basis. Therefore a spatial power covariance structure (with time as the distance measure) was used to account for the correlation among repeated measures of A1c from the same participant.

Results

Demographics and Participation

As shown in [Figure 1](#), 254 participants from across the United States responded to online advertisements and met CDC DPRP eligibility criteria. Of these, 220 participants completed the initial assessment and online set-up process and enrolled in the intervention on April 29, 2012. Demographic characteristics of study participants are reported in a previous study [5]. Briefly, as outlined in [Table 1](#), the participants were socioeconomically diverse. The baseline BMI of these 220 participants was 36.6 kg/m². A subset of these participants (n=187) completed at least 4 core lessons and were thus designated program starters, and a further subset of these participants (n=155) completed at least 9 core lessons and were thus designated program completers.






Figure 1. Participant recruitment and retention flow chart.

[View this figure](#)

Characteristic	N (%)	P
Gender		
Male	108 (50)	
Female	107 (50)	
Race		
White	108 (50)	
Black	107 (50)	
Hispanic	108 (50)	
Asian	107 (50)	
Other	108 (50)	
Ethnicity		
Hispanic	108 (50)	
Non-Hispanic	107 (50)	
Education		
High school or less	108 (50)	
Some college	107 (50)	
College graduate	108 (50)	
Postgraduate	107 (50)	
Income		
< \$10,000	108 (50)	
\$10,000 - \$19,999	107 (50)	
\$20,000 - \$29,999	108 (50)	
\$30,000 - \$39,999	107 (50)	
\$40,000 - \$49,999	108 (50)	
\$50,000 - \$59,999	107 (50)	
\$60,000 - \$69,999	108 (50)	
\$70,000 - \$79,999	107 (50)	
\$80,000 - \$89,999	108 (50)	
\$90,000 - \$99,999	107 (50)	
\$100,000 or more	108 (50)	

Table 1. Baseline demographic characteristics of participants.

[View this table](#)

Engagement

As reported in a previous publication, during the first year of the program, participants were engaged in the program through various activities.

Sections





Because lessons are limited to the first year, no engagement benchmarks exist beyond 1 year. However, participants continued to weigh in an average of 6.2 (SD 0.3) of 12 months and logged in an average of 3.5 (SD 0.3) of 12 months between years 1 and 2. In order to evenly compare engagement rates between years 1 and 2, the last 8 months of the year 1 (ie, the Sustain post-core period) was compared to the first 8 months of year 2. *Prevent* program starters weighed in an average of 4.6 (SD 0.2) of the last 8 months of year 1 versus 4.5 (SD 0.2) of the first 8 months of year 2, a non-significant difference ($P=.708$). *Prevent* program starters logged in on an average of 3.5 of 8 (SD 0.3) last months of the year 1 versus 2.7 of 8 (SD 0.3) first months of year 2, which was a significant difference ($P=.008$).

Changes in Body Weight and A1c

Among program starters, 100% (187/187) had an initial baseline weight measurement, 78.6% (147/187) had at least one weight measurement between 15 and 17 weeks, 79.1% (148/187) had at least one weight measurement between 11 and 13 months, and 70.1% (131/187) had at least one weight measurement between 22 and 26 months.

Changes in body weight and A1c after 16 weeks and 1 year were reported in a previous publication and reported here alongside 2-year data in [Tables 2](#) and [3](#) [6]. From baseline to year 2, program starters and completers achieved mean weight loss of 4.2% (SD 0.8%) and 4.3% (SD 0.8%), respectively. Program starters' and completers' weight loss was also maintained from year 1 to year 2, with no significant change ($P=.25$ and $.20$, respectively). Intention-to-treat analyses were also conducted on all enrollees: from baseline to year 2, enrollees achieved an average weight loss of 4.5% (SD 0.7%), though the majority of these additional participants did not weigh in after 1 year, which contributed to a higher possibility of model error. Among the program starters who reported weight between 22 and 26 months ($n=131$), 52 (40.0%) continued to meet or exceeded the CDC DPRP 5% weight loss benchmark at 24 months.

Because A1c measurement was optional, compliance was lower. Among program starters, 75.4% (141/187) had an initial baseline A1c measurement, 36.4% (68/187) had an A1c measurement between months 6-8 (due to A1c being a lagging measurement), 55.6% (104/187) had an A1c measurement between months 12-14, and 52.9% (99/187) had an A1c measurement between months 24-28.

As shown in [Tables 2](#) and [3](#), from baseline to year 2, program starters and completers reduced A1c by 0.43% (SD 0.08) and 0.46% (SD 0.08), respectively. Program starters' and completers' A1c reduction was also maintained from year 1 to year 2, with no significant change ($P=.39$ and $.38$, respectively).

Measure	Baseline	Year 1	Year 2
Program starters	187	187	187
Completers	147	147	147
Enrollees	187	187	187
Program starters	187	187	187
Completers	147	147	147
Enrollees	187	187	187
Program starters	187	187	187
Completers	147	147	147
Enrollees	187	187	187
Program starters	187	187	187
Completers	147	147	147
Enrollees	187	187	187

Sections



[illegible]

Table 3. Changes in body weight and A1c of participants over time.

[View this table](#)

Discussion

Principal Findings

Results indicate that the *Prevent* Internet-based diabetes prevention program helped participants who started or completed the program achieve significant reductions in body weight and A1c after 2 years. Of note, weight loss was largely maintained between years 1 and 2; this is one of the few studies to exhibit this weight maintenance effect after the active intervention ended.

Furthermore, A1c continued to show an average reduction from within the prediabetes range (5.7%-6.4%) to the normal range (<5.7%), in contrast to an expected annual rate of progression of approximately 4% from prediabetes to type 2 diabetes [14]. Thus, these results address critiques regarding the ability of DPP translations to show long-term effects on weight loss and A1c reduction [7,9].

The ability of *Prevent* to show sustained effects may be due to unique aspects of the program. Although the active intervention was 12 months, participants have continued access to *Prevent* and can proactively use the intervention. Thus, by allowing ongoing health tracking and social support, digital therapeutics can promote long-term maintenance of lifestyle changes with little ongoing intervention cost or maintenance.

Although *Prevent* exceeded CDC DPRP benchmarks for program completion during the 16-week core phase, 30% of participants who initially enrolled still did not complete the program. We hypothesize that one of the key advantages of digital therapeutics—accessibility—may also be a disadvantage when it comes to program completion. The lower barrier to Internet-based program entry inevitably increases the number of participants with lower motivation, who may have never made the effort to show up to an in-person program.

Strengths and Limitations

Study strengths include longitudinal collection of body weight and A1c data and statistical analysis using linear mixed models, which allow for more robust estimation over time and missing data. Furthermore, recruitment, intervention, and assessment were done exclusively remotely, in contrast to most digital health studies that require in-person orientation or follow-up assessment. This enhances the generalizability of the findings to “real-world” com

Sections





70% of program starters had weight data at year 2 and 53% had A1c data at year 2, which limits generalizability regarding outcomes on all participants. Thus, conclusions regarding weight loss and A1c reductions are limited to program starters and completers. In addition, while adherence to program behaviors (eg, lesson completion, tracking of weight and physical activity) were assessed according to CDC DPP standards, adherence to health behaviors (eg, diet and exercise goals) were not, which limits causal inference.

Conclusions

Results of this study suggest that the *Prevent* Internet-based diabetes prevention program was able to produce significant reductions in body weight and A1c that are sustained over the course of 2 years, even after active intervention ended after 1 year. Further investigation is warranted to test digital therapeutics as a scalable solution to address national diabetes and cardiovascular disease prevention efforts.

Acknowledgments

The authors wish to acknowledge Dr Ann Albright and Dr Ed Gregg at the CDC for their support of this work, and the Diabetes Prevention Program Research Group for performing the original DPP trial, on which the *Prevent* curriculum is based.

Authors' Contributions

Data were collected by SCS and provided unmodified to LJ, who independently conducted data analysis and wrote the results section. ALP provided consultation and editing of the paper for accuracy.

Conflicts of Interest

This work was funded by Omada Health, a company that makes and owns online behavior change programs, including the *Prevent* program, which is subject of this study. SCS is medical director for Omada Health and receives a salary and stock options. LJ is a paid statistical consultant, and ALP is a scientific advisor for Omada Health and receives stock options.

References

1. Bullard KM, Saydah SH, Imperatore G, Cowie CC, Gregg EW, Geiss LS, et al. Secular changes in U.S. Prediabetes prevalence defined by hemoglobin A1c and fasting plasma glucose: National Health and Nutrition Examination Surveys, 1999-2010. *Diabetes Care* 2012;35(9):2286-2292 [FREE Full text] [CrossRef] [Medline]

Sections





healthful diet and physical activity for cardiovascular disease prevention in adults with cardiovascular risk factors: U.S. Preventive Services Task Force Recommendation Statement. *Ann Intern Med* 2014;161(8):587-593. [[CrossRef](#)] [[Medline](#)]

4. Webb TL, Joseph J, Yardley L, Michie S. Using the internet to promote health behavior change: a systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy. *J Med Internet Res* 2010;12(1):e4 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
5. Paulweber B, Valensi P, Lindström J, Lalic NM, Greaves CJ, McKee M, et al. A European evidence-based guideline for the prevention of type 2 diabetes. *Horm Metab Res* 2010;42 Suppl 1:S3-36 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
6. Sepah SC, Jiang L, Peters AL. Translating the Diabetes Prevention Program into an Online Social Network: Validation against CDC Standards. *Diabetes Educ* 2014;40(4):435-443 [[FREE Full text](#)] [[CrossRef](#)] [[Medline](#)]
7. Centers for Disease Control and Prevention. 2011. Diabetes Prevention Recognition Program Standards and Operating Procedures URL: http://www.cdc.gov/diabetes/prevention/pdf/DPRP_Standards_09-02-2011.pdf [accessed 2014-11-18] [[WebCite Cache](#)]
8. Yudkin JS, Montori VM. The epidemic of pre-diabetes: the medicine and the politics. *BMJ* 2014;349:g4485. [[CrossRef](#)] [[Medline](#)]
9. Dunkley AJ, Bodicoat DH, Greaves CJ, Russell C, Yates T, Davies MJ, et al. Diabetes prevention in the real world: effectiveness of pragmatic lifestyle interventions for the prevention of type 2 diabetes and of the impact of adherence to guideline recommendations: a systematic review and meta-analysis. *Diabetes Care* 2014 Apr;37(4):922-933. [[CrossRef](#)] [[Medline](#)]
10. Kahn R, Davidson MB. The reality of type 2 diabetes prevention. *Diabetes Care* 2014;37(4):943-949. [[CrossRef](#)] [[Medline](#)]
11. Prevent by Omada Health. URL: <http://preventnow.com/> [accessed 2014-12-21] [[WebCite Cache](#)]
12. Steinberg DM, Tate DF, Bennett GG, Ennett S, Samuel-Hodge C, Ward DS. Daily self-weighing and adverse psychological outcomes: a randomized controlled trial. *Am J Prev Med* 2014;46(1):24-29. [[CrossRef](#)] [[Medline](#)]
13. DTI Laboratories, Inc. Technical Data. URL: http://dtilaboratories.com/new_page_9.htm [accessed 2015-01-02] [[WebCite Cache](#)]
14. Morris DH, Khunti K, Achana F, Srinivasan B, Gray LJ, Davies MJ, et al. Progression rates from HbA1c 6.0-6.4% and other prediabetes definitions to type 2 diabetes: a meta-analysis. *Diabetologia* 2013 Jul;56(7):1489-1493. [[CrossRef](#)] [[Medline](#)]





DPP: Diabetes Prevention Program

DPRP: Diabetes Prevention Recognition Program

HPLC: High-Performance Liquid Chromatography

IRB: Institutional Review Board

USPSTF: US Preventive Services Task Force

Edited by G Eysenbach; submitted 17.11.14; peer-reviewed by L Saslow, K Schneider; comments to author 11.12.14; revised version received 02.01.15; accepted 16.03.15; published 10.04.15

Copyright

©S Cameron Sepah, Luohua Jiang, Anne L Peters. Originally published in the Journal of Medical Internet Research (<http://www.jmir.org>), 10.04.2015.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in the Journal of Medical Internet Research, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.jmir.org/>, as well as this copyright and license information must be included.





Journals

JMIR Research Protocols

JMIR mHealth and uHealth

interactive Journal of Medical Research

JMIR Medical Informatics

JMIR Mental Health

Medicine 2.0

JMIR Serious Games

JMIR Human Factors

JMIR Rehabilitation and Assistive Technologies

JMIR Public Health and Surveillance

JMIR Cancer

JMIR Medical Education

Editorial Policies

Focus and Scope

Section Policies

Peer Review Process

Publication Frequency

Memberships

Author Self-Archiving

Why you should choose JMIR to publish your research - advantages of the journal

Indexing and Impact Factor

Editorial Board Policy

Record Keeping and Research Policy

Subscriptions (Membership)

Theme Issues and Guest Editors

Trademarks and Service Marks Policy

Sections





Submissions

[Online Submissions](#)

[Author Guidelines](#)

[Copyright Notice](#)

[Privacy Statement](#)

Other

[Announcements](#)

[Editorial Team](#)

[Feedback](#)

[Contact Us](#)

Copyright © 2015 JMIR Publications Inc.

Sections

