

**SAMPLE QUESTION PAPER**


**MARKING SCHEME**


**CLASS XII**

**MATHEMATICS (CODE-041)**

**SECTION: A (Solution of MCQs of 1 Mark each)**

Q no.	ANS	HINTS/SOLUTION
1	(d)	$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, A^2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$
2	(d)	$(A+B)^{-1} = B^{-1} + A^{-1}.$
3	(b)	<p>Area = <math>\frac{1}{2} \begin{vmatrix} -3 &amp; 0 &amp; 1 \\ 3 &amp; 0 &amp; 1 \\ 0 &amp; k &amp; 1 \end{vmatrix}</math>, given that the area = 9 sq unit.</p> <p><math>\Rightarrow \pm 9 = \frac{1}{2} \begin{vmatrix} -3 &amp; 0 &amp; 1 \\ 3 &amp; 0 &amp; 1 \\ 0 &amp; k &amp; 1 \end{vmatrix}</math>; expanding along <math>C_2</math>, we get <math>\Rightarrow k = \pm 3</math>.</p>
4	(a)	<p>Since, <math>f</math> is continuous at <math>x = 0</math>,</p> <p>therefore, <math>L.H.L = R.H.L = f(0) = a \text{ finite quantity}.</math></p> <p><math>\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)</math></p> <p><math>\Rightarrow \lim_{x \rightarrow 0^-} \frac{-kx}{x} = \lim_{x \rightarrow 0^+} 3 = 3 \Rightarrow k = -3.</math></p>
5	(d)	<p>Vectors <math>2\hat{i} + 3\hat{j} - 6\hat{k}</math> &amp; <math>6\hat{i} + 9\hat{j} - 18\hat{k}</math> are parallel and the fixed point <math>\hat{i} + \hat{j} - \hat{k}</math> on the line <math>\vec{r} = \hat{i} + \hat{j} - \hat{k} + \lambda(2\hat{i} + 3\hat{j} - 6\hat{k})</math> does not satisfy the other line <math>\vec{r} = 2\hat{i} - \hat{j} - \hat{k} + \mu(6\hat{i} + 9\hat{j} - 18\hat{k})</math>; where <math>\lambda</math> &amp; <math>\mu</math> are scalars.</p>
6	(c)	The degree of the differential equation $\left[1 + \left(\frac{dy}{dx}\right)^2\right]^3 = \left(\frac{d^2y}{dx^2}\right)^2$ is 2
7	(b)	<p><math>Z = px + qy \dots (i)</math></p> <p>At <math>(3,0)</math>, <math>Z = 3p \dots (ii)</math> and at <math>(1,1)</math>, <math>Z = p + q \dots (iii)</math></p> <p>From <math>(ii)</math> &amp; <math>(iii)</math>, <math>3p = p + q \Rightarrow 2p = q</math>.</p>

8	<b>(a)</b>	<p>Given, <math>ABCD</math> is a rhombus whose diagonals bisect each other. <math> \vec{EA}  =  \vec{EC} </math> and <math> \vec{EB}  =  \vec{ED} </math> but since they are opposite to each other so they are of opposite signs  <math>\Rightarrow \vec{EA} = -\vec{EC}</math> and <math>\vec{EB} = -\vec{ED}</math>.</p>  <p><math>\Rightarrow \vec{EA} + \vec{EC} = \vec{0} \dots\dots (i)</math> and <math>\vec{EB} + \vec{ED} = \vec{0} \dots\dots (ii)</math>  Adding (i) and (ii), we get <math>\vec{EA} + \vec{EB} + \vec{EC} + \vec{ED} = \vec{0}</math>.</p>
9	<b>(b)</b>	$f(x) = e^{\cos^2 x} \sin^3(2n+1)x$ $f(-x) = e^{\cos^2(-x)} \sin^3(2n+1)(-x)$ $f(-x) = -e^{\cos^2 x} \sin^3(2n+1)x$ $\therefore f(-x) = -f(x)$ $\text{So, } \int_{-\pi}^{\pi} e^{\cos^2 x} \sin^3(2n+1)x \, dx = 0$
10	<b>(b)</b>	<p>Matrix <math>A</math> is a skew symmetric matrix of odd order. <math>\therefore  A  = 0</math>.</p>
11	<b>(c)</b>	<p>We observe, <math>(0,0)</math> does not satisfy the inequality <math>x - y \geq 1</math>  So, the half plane represented by the above inequality will not contain origin therefore, it will not contain the shaded feasible region.</p>
12	<b>(b)</b>	<p>Vector component of <math>\vec{a}</math> along <math>\vec{b} = \left( \frac{\vec{a} \cdot \vec{b}}{ \vec{b} ^2} \right) \vec{b} = \frac{18}{25} (3\hat{j} + 4\hat{k})</math>.</p>
13	<b>(d)</b>	$ \text{adj}(2A)  =  (2A) ^2 = (2^3  A )^2 = 2^6  A ^2 = 2^6 \times (-2)^2 = 2^8$ .
14	<b>(d)</b>	<p><b>Method 1:</b>  Let <math>A, B, C</math> be the respective events of solving the problem. Then, <math>P(A) = \frac{1}{2}, P(B) = \frac{1}{3}</math> and <math>P(C) = \frac{1}{4}</math>. Here, <math>A, B, C</math> are independent events.  Problem is solved if at least one of them solves the problem.  Required probability is <math>P(A \cup B \cup C) = 1 - P(\bar{A})P(\bar{B})P(\bar{C})</math></p>

		$= 1 - \left(1 - \frac{1}{2}\right) \left(1 - \frac{1}{3}\right) \left(1 - \frac{1}{4}\right) = 1 - \frac{1}{4} = \frac{3}{4}.$ <p><b>Method 2:</b></p> <p>The problem will be solved if one or more of them can solve the problem. The probability is</p> $P(\overline{ABC}) + P(\overline{AB}\overline{C}) + P(\overline{A}\overline{BC}) + P(\overline{AB}\overline{C}) + P(\overline{A}\overline{B}\overline{C}) + P(\overline{ABC}) + P(ABC)$ $= \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{3}{4} + \frac{1}{2} \cdot \frac{1}{3} \cdot \frac{3}{4} + \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{1}{4} + \frac{1}{2} \cdot \frac{1}{3} \cdot \frac{1}{4} + \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{1}{4} + \frac{1}{2} \cdot \frac{1}{3} \cdot \frac{1}{4} + \frac{1}{2} \cdot \frac{1}{3} \cdot \frac{1}{4} = \frac{3}{4}.$ <p><b>Method 3:</b></p> <p>Let us think quantitatively. Let us assume that there are 100 questions given to <b>A</b>. <b>A</b> solves <math>\frac{1}{2} \times 100 = 50</math> questions then remaining <b>50</b> questions is given to <b>B</b> and <b>B</b> solves <math>50 \times \frac{1}{3} = 16.67</math> questions. Remaining <math>50 \times \frac{2}{3}</math> questions is given to <b>C</b> and <b>C</b> solves <math>50 \times \frac{2}{3} \times \frac{1}{4} = 8.33</math> questions.</p> <p>Therefore, number of questions solved is <math>50 + 16.67 + 8.33 = 75</math>.</p> <p>So, required probability is <math>\frac{75}{100} = \frac{3}{4}</math>.</p>
15	(c)	<p><b>Method 1:</b></p> $ydx - xdy = 0 \Rightarrow \frac{ydx - xdy}{y^2} = 0 \Rightarrow d\left(\frac{x}{y}\right) = 0 \Rightarrow x = \frac{1}{c}y \Rightarrow y = cx.$ <p><b>Method 2:</b></p> $ydx - xdy = 0 \Rightarrow ydx = xdy \Rightarrow \frac{dy}{y} = \frac{dx}{x}; \text{ on integrating } \int \frac{dy}{y} = \int \frac{dx}{x}$ $\log_e  y  = \log_e  x  + \log_e  c $ <p>since <math>x, y, c &gt; 0</math>, we write <math>\log_e y = \log_e x + \log_e c \Rightarrow y = cx</math>.</p>
16	(d)	Dot product of two mutually perpendicular vectors is zero. $\Rightarrow 2 \times 3 + (-1)\lambda + 2 \times 1 = 0 \Rightarrow \lambda = 8$ .
17	(c)	<p><b>Method 1:</b></p> $f(x) = x +  x  = \begin{cases} 2x, & x \geq 0 \\ 0, & x < 0 \end{cases}$  <p>There is a sharp corner at <math>x = 0</math>, so <math>f(x)</math> is not differentiable at <math>x = 0</math>.</p> <p><b>Method 2:</b></p>

		<p><math>Lf'(0) = 0</math> &amp; <math>Rf'(0) = 2</math>; so, the function is not differentiable at <math>x = 0</math></p> <p>For <math>x \geq 0, f(x) = 2x</math> (linear function) &amp; when <math>x &lt; 0, f(x) = 0</math> (constant function)</p> <p>Hence <math>f(x)</math> is differentiable when <math>x \in (-\infty, 0) \cup (0, \infty)</math>.</p>
18	(d)	We know, $l^2 + m^2 + n^2 = 1 \Rightarrow \left(\frac{1}{c}\right)^2 + \left(\frac{1}{c}\right)^2 + \left(\frac{1}{c}\right)^2 = 1 \Rightarrow 3\left(\frac{1}{c}\right)^2 = 1 \Rightarrow c = \pm\sqrt{3}$ .
19	(a)	$\frac{d}{dx}(f(x)) = (x-1)^3(x-3)^2$ Assertion : $f(x)$ has a minimum at $x = 1$ is true as $\frac{d}{dx}(f(x)) < 0, \forall x \in (1-h, 1)$ and $\frac{d}{dx}(f(x)) > 0, \forall x \in (1, 1+h)$ ; where, 'h' is an infinitesimally small positive quantity, which is in accordance with the Reason statement.
20	(d)	<b>Assertion is false.</b> As element 4 has no image under $f$ , so relation $f$ is not a function. <b>Reason is true.</b> The given function $f : \{1, 2, 3\} \rightarrow \{x, y, z, p\}$ is one – one, as for each $a \in \{1, 2, 3\}$ , there is different image in $\{x, y, z, p\}$ under $f$ .

### Section –B

[This section comprises of solution of very short answer type questions (VSA) of 2 marks each]

21	$\begin{aligned} \sin^{-1}\left(\cos\left(\frac{33\pi}{5}\right)\right) &= \sin^{-1}\cos\left(6\pi + \frac{3\pi}{5}\right) = \sin^{-1}\cos\left(\frac{3\pi}{5}\right) = \sin^{-1}\sin\left(\frac{\pi}{2} - \frac{3\pi}{5}\right) \\ &= \frac{\pi}{2} - \frac{3\pi}{5} = -\frac{\pi}{10}. \end{aligned}$	1 1
21 OR	$-1 \leq (x^2 - 4) \leq 1 \Rightarrow 3 \leq x^2 \leq 5 \Rightarrow \sqrt{3} \leq  x  \leq \sqrt{5}$ $\Rightarrow x \in [-\sqrt{5}, -\sqrt{3}] \cup [\sqrt{3}, \sqrt{5}]$ . So, required domain is $[-\sqrt{5}, -\sqrt{3}] \cup [\sqrt{3}, \sqrt{5}]$ .	1 1
22	$f(x) = xe^x \Rightarrow f'(x) = e^x(x+1)$ When $x \in [-1, \infty), (x+1) \geq 0$ & $e^x > 0 \Rightarrow f'(x) \geq 0 \therefore f(x)$ increases in this interval. <i>or, we can write</i> $f(x) = xe^x \Rightarrow f'(x) = e^x(x+1)$ For $f(x)$ to be increasing, we have $f'(x) = e^x(x+1) \geq 0 \Rightarrow x \geq -1$ as $e^x > 0, \forall x \in \mathbb{R}$ Hence, the required interval where $f(x)$ increases is $[-1, \infty)$ .	1 1 $\frac{1}{2}$ 1 $\frac{1}{2}$
23	<b>Method 1 :</b> $f(x) = \frac{1}{4x^2 + 2x + 1}$ ,	



	<p>and when <math>x \in \left(-\frac{1}{4}, -\frac{1}{4} + h\right)</math>, <math>4x &gt; -1 \Rightarrow 8x &gt; -2 \Rightarrow 8x + 2 &gt; 0 \Rightarrow -(8x + 2) &lt; 0</math>          and <math>(4x^2 + 2x + 1)^2 &gt; 0 \Rightarrow f'(x) &lt; 0</math>. This shows that <math>x = -\frac{1}{4}</math> is the point of local maxima.  <math>\therefore</math> maximum value of <math>f(x)</math> is <math>f\left(-\frac{1}{4}\right) = \frac{1}{4\left(-\frac{1}{4}\right)^2 + 2\left(-\frac{1}{4}\right) + 1} = \frac{4}{3}</math>.</p>	$\frac{1}{2}$
23 OR	<p>For maxima and minima, <math>P'(x) = 0 \Rightarrow 42 - 2x = 0</math>  <math>\Rightarrow x = 21</math> and <math>P''(x) = -2 &lt; 0</math>          So, <math>P(x)</math> is maximum at <math>x = 21</math>.          The maximum value of <math>P(x) = 72 + (42 \times 21) - (21)^2 = 513</math> i.e., the maximum profit is ₹ 513.</p>	$\frac{1}{2}$
24	<p>Let <math>f(x) = \log\left(\frac{2-x}{2+x}\right)</math>          We have, <math>f(-x) = \log\left(\frac{2+x}{2-x}\right) = -\log\left(\frac{2-x}{2+x}\right) = -f(x)</math>          So, <math>f(x)</math> is an odd function. <math>\therefore \int_{-1}^1 \log\left(\frac{2-x}{2+x}\right) dx = 0</math>.</p>	1 1
25	<p><math>f(x) = x^3 + x</math>, for all <math>x \in \mathbb{R}</math>.  <math>\frac{d}{dx}(f(x)) = f'(x) = 3x^2 + 1</math>; for all <math>x \in \mathbb{R}</math>, <math>x^2 \geq 0 \Rightarrow f'(x) &gt; 0</math>          Hence, no critical point exists.</p>	$\frac{1}{2}$ $\frac{1}{2}$


### Section -C

**[This section comprises of solution short answer type questions (SA) of 3 marks each]**

26	<p>We have, <math>\frac{2x^2+3}{x^2(x^2+9)}</math>. Now, let <math>x^2 = t</math>          So, <math>\frac{2t+3}{t(t+9)} = \frac{A}{t} + \frac{B}{t+9}</math>, we get <math>A = \frac{1}{3}</math> &amp; <math>B = \frac{5}{3}</math>  <math>\int \frac{2x^2+3}{x^2(x^2+9)} dx = \frac{1}{3} \int \frac{dx}{x^2} + \frac{5}{3} \int \frac{dx}{x^2+9}</math>  <math>= -\frac{1}{3x} + \frac{5}{9} \tan^{-1}\left(\frac{x}{3}\right) + c</math>, where 'c' is an arbitrary constant of integration.</p>	$\frac{1}{2}$ 1 $\frac{1}{2}$ 1
27	<p>We have, (i) <math>\sum P(X_i) = 1 \Rightarrow k + 2k + 3k = 1 \Rightarrow k = \frac{1}{6}</math>.</p>	1 1



	<p>So equation (i) becomes <math>v + y \frac{dv}{dy} = v + \frac{y}{e^v}</math></p> $\Rightarrow y \frac{dv}{dy} = \frac{y}{e^v}$ $\Rightarrow e^v dv = dy$ <p>On integrating we get, <math>\int e^v dv = \int dy \Rightarrow e^v = y + c \Rightarrow e^{x/y} = y + c</math>  <i>where 'c' is an arbitrary constant of integration.</i></p>	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
29 OR	<p>The given Differential equation is</p> $(\cos^2 x) \frac{dy}{dx} + y = \tan x$ <p>Dividing both the sides by <math>\cos^2 x</math>, we get</p> $\frac{dy}{dx} + \frac{y}{\cos^2 x} = \frac{\tan x}{\cos^2 x}$ $\frac{dy}{dx} + y(\sec^2 x) = \tan x(\sec^2 x) \dots\dots\dots(i)$ <p>Comparing with <math>\frac{dy}{dx} + Py = Q</math></p> $P = \sec^2 x, Q = \tan x \cdot \sec^2 x$ <p>The Integrating factor is, <math>IF = e^{\int P dx} = e^{\int \sec^2 x dx} = e^{\tan x}</math></p> <p>On multiplying the equation (i) by <math>e^{\tan x}</math>, we get</p> $\frac{d}{dx}(y \cdot e^{\tan x}) = e^{\tan x} \tan x (\sec^2 x) \Rightarrow d(y \cdot e^{\tan x}) = e^{\tan x} \tan x (\sec^2 x) dx$ <p>On integrating we get, <math>y \cdot e^{\tan x} = \int t \cdot e^t dt + c_1</math>; where, <math>t = \tan x</math> so that <math>dt = \sec^2 x dx</math></p> $= te^t - e^t + c = (\tan x) e^{\tan x} - e^{\tan x} + c$ <p><math>\therefore y = \tan x - 1 + c_1 (e^{-\tan x})</math>, where '<math>c_1</math>' &amp; '<math>c</math>' are arbitrary constants of integration.</p>	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
30	<p>The feasible region determined by the constraints, <math>x + 2y \geq 100</math>, <math>2x - y \leq 0</math>, <math>2x + y \leq 200</math>, <math>x, y \geq 0</math>, is given below.</p>	



**A (0, 50), B (20, 40), C (50, 100) and D (0, 200)** are the corner points of the feasible region.

The values of  $Z$  at these corner points are given below.

Corner point	Corresponding value of $Z = x + 2y$	
<b>A (0, 50)</b>	<b>100</b>	Minimum
<b>B (20, 40)</b>	<b>100</b>	Minimum
<b>C (50, 100)</b>	<b>250</b>	
<b>D (0, 200)</b>	<b>400</b>	

**30 OR**

The minimum value of  $Z$  is **100** at all the points on the line segment joining the points **(0,50)** and **(20,40)**.


The feasible region determined by the constraints,  $x \geq 3, x + y \geq 5, x + 2y \geq 6, y \geq 0$ . is given below.

**1**  
**2**

**1**  
**2**

**1**  
**2**

Here, it  
be seen  
the



feasible region is unbounded.

The values of  $Z$  at corner points  $A(3, 2)$ ,  $B(4, 1)$  and  $C(6, 0)$  are given below.

Corner point	Corresponding value of $Z = -x + 2y$
$A(3, 2)$	1 (may or may not be the maximum value)
$B(4, 1)$	-2
$C(6, 0)$	-6

Since the feasible region is unbounded,  $Z = 1$  may or may not be the maximum value.

Now, we draw the graph of the inequality,  $-x + 2y > 1$ , and we check whether the resulting open half-plane has any point/s, in common with the feasible region or not.

Here, the resulting open half plane has points in common with the feasible region.

Hence,  $Z = 1$  is not the maximum value. We conclude,  $Z$  has no maximum value.

31

$$\frac{y}{x} = \log_e \left( \frac{x}{a+bx} \right) = \log_e x - \log_e (a+bx)$$

On differentiating with respect to  $x$ , we get

$$\Rightarrow \frac{x \frac{dy}{dx} - y}{x^2} = \frac{1}{x} - \frac{1}{a+bx} \frac{d}{dx}(a+bx) = \frac{1}{x} - \frac{b}{a+bx}$$

$$\Rightarrow x \frac{dy}{dx} - y = x^2 \left( \frac{1}{x} - \frac{b}{a+bx} \right) = \frac{ax}{a+bx}$$

On differentiating again with respect to  $x$ , we get

$$\Rightarrow x \frac{d^2y}{dx^2} + \frac{dy}{dx} - \frac{dy}{dx} = \frac{(a+bx)a - ax(b)}{(a+bx)^2}$$

can  
that

1

$\frac{1}{2}$

$\frac{1}{2}$

1

$\frac{1}{2}$

	$\Rightarrow x \frac{d^2y}{dx^2} = \left( \frac{a}{a+bx} \right)^2.$	$\frac{1}{2}$
--	--	---------------


### Section -D

[This section comprises of solution of long answer type questions (LA) of 5 marks each]

32		1
	<p>To find the point of intersections of the curve <math>y = x^2 + 1</math> and the line <math>y = x + 1</math>, we write <math>x^2 + 1 = x + 1 \Rightarrow x(x - 1) = 0 \Rightarrow x = 0, 1</math>.</p> <p>So, the point of intersections <math>P(0, 1)</math> and <math>Q(1, 2)</math>.</p> <p>Area of the shaded region <math>OPQRTSO = (\text{Area of the region } OSQPO + \text{Area of the region } STRQS)</math></p> $  \begin{aligned}  &= \int_0^1 (x^2 + 1) dx + \int_1^2 (x + 1) dx \\  &= \left[ \frac{x^3}{3} + x \right]_0^1 + \left[ \frac{x^2}{2} + x \right]_1^2 \\  &= \left[ \left( \frac{1}{3} + 1 \right) - 0 \right] + \left[ (2 + 2) - \left( \frac{1}{2} + 1 \right) \right] \\  &= \frac{23}{6} \quad \text{Hence the required area is } \frac{23}{6} \text{ sq units.}  \end{aligned}  $	1 1 $\frac{1}{2}$ $\frac{1}{2}$ 1
33	<p>Let <math>(a, b)</math> be an arbitrary element of <math>\mathbb{N} \times \mathbb{N}</math>. Then, <math>(a, b) \in \mathbb{N} \times \mathbb{N}</math> and <math>a, b \in \mathbb{N}</math></p> <p>We have, <math>ab = ba</math>; (As <math>a, b \in \mathbb{N}</math> and multiplication is commutative on <math>\mathbb{N}</math>)</p> <p><math>\Rightarrow (a, b)R(a, b)</math>, according to the definition of the relation <math>R</math> on <math>\mathbb{N} \times \mathbb{N}</math></p> <p>Thus <math>(a, b)R(a, b)</math>, <math>\forall (a, b) \in \mathbb{N} \times \mathbb{N}</math>.</p> <p>So, <math>R</math> is reflexive relation on <math>\mathbb{N} \times \mathbb{N}</math>.</p> <p>Let <math>(a, b), (c, d)</math> be arbitrary elements of <math>\mathbb{N} \times \mathbb{N}</math> such that <math>(a, b)R(c, d)</math>.</p>	1

	<p>Then, <math>(a,b)R(c,d) \Rightarrow ad = bc \Rightarrow bc = ad</math>; (changing LHS and RHS)</p> <p><math>\Rightarrow cb = da</math>; (As <math>a, b, c, d \in \mathbb{N}</math> and multiplication is commutative on <math>\mathbb{N}</math>)</p> <p><math>\Rightarrow (c,d)R(a,b)</math>; according to the definition of the relation <math>R</math> on <math>\mathbb{N} \times \mathbb{N}</math></p> <p>Thus <math>(a,b)R(c,d) \Rightarrow (c,d)R(a,b)</math></p> <p>So, <math>R</math> is symmetric relation on <math>\mathbb{N} \times \mathbb{N}</math>.</p> <p>Let <math>(a,b), (c,d), (e,f)</math> be arbitrary elements of <math>\mathbb{N} \times \mathbb{N}</math> such that</p> <p><math>(a,b)R(c,d)</math> and <math>(c,d)R(e,f)</math>.</p> <p>Then <math>\begin{cases} (a,b)R(c,d) \Rightarrow ad = bc \\ (c,d)R(e,f) \Rightarrow cf = de \end{cases} \Rightarrow (ad)(cf) = (bc)(de) \Rightarrow af = be</math></p> <p><math>\Rightarrow (a,b)R(e,f)</math>; (according to the definition of the relation <math>R</math> on <math>\mathbb{N} \times \mathbb{N}</math>)</p> <p>Thus <math>(a,b)R(c,d)</math> and <math>(c,d)R(e,f) \Rightarrow (a,b)R(e,f)</math></p> <p>So, <math>R</math> is transitive relation on <math>\mathbb{N} \times \mathbb{N}</math>.</p> <p>As the relation <math>R</math> is reflexive, symmetric and transitive so, it is equivalence relation on <math>\mathbb{N} \times \mathbb{N}</math>.</p> <p><math>\begin{aligned} [(2,6)] &amp;= \{(x,y) \in \mathbb{N} \times \mathbb{N} : (x,y)R(2,6)\} \\ &amp;= \{(x,y) \in \mathbb{N} \times \mathbb{N} : 3x = y\} \\ &amp;= \{(x,3x) : x \in \mathbb{N}\} = \{(1,3), (2,6), (3,9), \dots\} \end{aligned}</math></p>	1 1 $\frac{1}{2}$ $\frac{1}{2}$ 1
33 OR	<p>We have, <math>f(x) = \begin{cases} \frac{x}{1+x}, &amp; \text{if } x \geq 0 \\ \frac{x}{1-x}, &amp; \text{if } x &lt; 0 \end{cases}</math></p> <p>Now, we consider the following cases</p> <p><b>Case 1:</b> when <math>x \geq 0</math>, we have <math>f(x) = \frac{x}{1+x}</math></p> <p>Injectivity: let <math>x, y \in \mathbb{R}^+ \cup \{0\}</math> such that <math>f(x) = f(y)</math>, then</p> $\Rightarrow \frac{x}{1+x} = \frac{y}{1+y} \Rightarrow x + xy = y + xy \Rightarrow x = y$ <p>So, <math>f</math> is injective function.</p> <p>Surjectivity : when <math>x \geq 0</math>, we have <math>f(x) = \frac{x}{1+x} \geq 0</math> and <math>f(x) = 1 - \frac{1}{1+x} &lt; 1</math>, as <math>x \geq 0</math></p> <p>Let <math>y \in [0,1)</math>, thus for each <math>y \in [0,1)</math> there exists <math>x = \frac{y}{1-y} \geq 0</math> such that <math>f(x) = \frac{\frac{y}{1-y}}{1+\frac{y}{1-y}} = y</math>.</p>	1 1 1

	<p>So, <math>f</math> is onto function on <math>[0, \infty)</math> to <math>[0, 1)</math>.</p> <p><b>Case 2:</b> when <math>x &lt; 0</math>, we have <math>f(x) = \frac{x}{1-x}</math></p> <p>Injectivity: Let <math>x, y \in \mathbb{R}^-</math> i.e., <math>x, y &lt; 0</math>, such that <math>f(x) = f(y)</math>, then</p> $\Rightarrow \frac{x}{1-x} = \frac{y}{1-y} \Rightarrow x - xy = y - xy \Rightarrow x = y$ <p>So, <math>f</math> is injective function.</p> <p>Surjectivity : <math>x &lt; 0</math>, we have <math>f(x) = \frac{x}{1-x} &lt; 0</math> also, <math>f(x) = \frac{x}{1-x} = -1 + \frac{1}{1-x} &gt; -1</math></p> $-1 < f(x) < 0.$ <p>Let <math>y \in (-1, 0)</math> be an arbitrary real number and there exists <math>x = \frac{y}{1+y} &lt; 0</math> such that,</p> $f(x) = f\left(\frac{y}{1+y}\right) = \frac{\frac{y}{1+y}}{1 - \frac{y}{1+y}} = y.$ <p>So, for <math>y \in (-1, 0)</math>, there exists <math>x = \frac{y}{1+y} &lt; 0</math> such that <math>f(x) = y</math>.</p> <p>Hence, <math>f</math> is onto function on <math>(-\infty, 0)</math> to <math>(-1, 0)</math>.</p> <p><b>Case 3:</b></p> <p>(Injectivity): Let <math>x &gt; 0</math> &amp; <math>y &lt; 0</math> such that <math>f(x) = f(y) \Rightarrow \frac{x}{1+x} = \frac{y}{1-y}</math></p> $\Rightarrow x - xy = y + xy \Rightarrow x - y = 2xy, \text{ here } LHS > 0 \text{ but } RHS < 0, \text{ which is inadmissible.}$ <p>Hence, <math>f(x) \neq f(y)</math> when <math>x \neq y</math>.</p> <p>Hence <math>f</math> is one-one and onto function.</p>	1
34	<p>The given system of equations can be written in the form <math>AX = B</math>,</p> <p>Where, <math>A = \begin{bmatrix} 2 &amp; 3 &amp; 10 \\ 4 &amp; -6 &amp; 5 \\ 6 &amp; 9 &amp; -20 \end{bmatrix}, X = \begin{bmatrix} 1/x \\ 1/y \\ 1/z \end{bmatrix}</math> and <math>B = \begin{bmatrix} 4 \\ 1 \\ 2 \end{bmatrix}</math></p> <p>Now, <math> A  = \begin{vmatrix} 2 &amp; 3 &amp; 10 \\ 4 &amp; -6 &amp; 5 \\ 6 &amp; 9 &amp; -20 \end{vmatrix} = 2(120 - 45) - 3(-80 - 30) + 10(36 + 36)</math></p> $= 2(75) - 3(-110) + 10(72) = 150 + 330 + 720 = 1200 \neq 0 \therefore A^{-1}$ exists. <p><math>\therefore adj A = \begin{bmatrix} 75 &amp; 110 &amp; 72 \\ 150 &amp; -100 &amp; 0 \\ 75 &amp; 30 &amp; -24 \end{bmatrix}^T = \begin{bmatrix} 75 &amp; 150 &amp; 75 \\ 110 &amp; -100 &amp; 30 \\ 72 &amp; 0 &amp; -24 \end{bmatrix}</math></p>	<p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p>

	<p>Hence, <math>A^{-1} = \frac{1}{ A }(\text{adj}A) = \frac{1}{1200} \begin{bmatrix} 75 &amp; 150 &amp; 75 \\ 110 &amp; -100 &amp; 30 \\ 72 &amp; 0 &amp; -24 \end{bmatrix}</math></p> <p>As, <math>AX = B \Rightarrow X = A^{-1}B \Rightarrow \begin{bmatrix} \frac{1}{x} \\ \frac{1}{y} \\ \frac{1}{z} \end{bmatrix} = \frac{1}{1200} \begin{bmatrix} 75 &amp; 150 &amp; 75 \\ 110 &amp; -100 &amp; 30 \\ 72 &amp; 0 &amp; -24 \end{bmatrix} \begin{bmatrix} 4 \\ 1 \\ 2 \end{bmatrix}</math></p> $= \frac{1}{1200} \begin{bmatrix} 300+150+150 \\ 440-100+60 \\ 288+0-48 \end{bmatrix} \Rightarrow \begin{bmatrix} \frac{1}{x} \\ \frac{1}{y} \\ \frac{1}{z} \end{bmatrix} = \frac{1}{1200} \begin{bmatrix} 600 \\ 400 \\ 240 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ \frac{1}{3} \\ \frac{1}{5} \end{bmatrix}$ <p>Thus, <math>\frac{1}{x} = \frac{1}{2}, \frac{1}{y} = \frac{1}{3}, \frac{1}{z} = \frac{1}{5}</math> Hence, <math>x = 2, y = 3, z = 5</math>.</p>	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $1$
35	<p>Let <math>P(1,6,3)</math> be the given point, and let '<math>L</math>' be the foot of the perpendicular from '<math>P</math>' to the given line <math>AB</math> (as shown in the figure below). The coordinates of a general point on the given line are given by</p>  $\frac{x-0}{1} = \frac{y-1}{2} = \frac{z-2}{3} = \lambda; \lambda \text{ is a scalar, i.e., } x = \lambda, y = 2\lambda + 1 \text{ and } z = 3\lambda + 2$ <p>Let the coordinates of <math>L</math> be <math>(\lambda, 2\lambda + 1, 3\lambda + 2)</math>.</p> <p>So, direction ratios of <math>PL</math> are <math>\lambda - 1, 2\lambda + 1 - 6</math> and <math>3\lambda + 2 - 3</math>, i.e. <math>\lambda - 1, 2\lambda - 5</math> and <math>3\lambda - 1</math>.</p> <p>Direction ratios of the given line are <math>1, 2</math> and <math>3</math>, which is perpendicular to <math>PL</math>.</p> <p>Therefore, <math>(\lambda - 1)1 + (2\lambda - 5)2 + (3\lambda - 1)3 = 0 \Rightarrow 14\lambda - 14 = 0 \Rightarrow \lambda = 1</math></p> <p>So, coordinates of <math>L</math> are <math>(1, 3, 5)</math>.</p>	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $1$




On solving the above equations , we get  $\lambda = \frac{2}{3}$  and  $\mu = 0$

So, the position vector of the points, at which they should be so that the distance between them is the shortest, are  $\frac{2}{3}(\hat{i} - \hat{j} + \hat{k})$  and  $\hat{i} - \hat{j}$ .

$$\overrightarrow{PQ} = \overrightarrow{OQ} - \overrightarrow{OP} = \frac{1}{3}\hat{i} - \frac{1}{3}\hat{j} - \frac{2}{3}\hat{k} \text{ and } |\overrightarrow{PQ}| = \sqrt{\left(\frac{1}{3}\right)^2 + \left(-\frac{1}{3}\right)^2 + \left(-\frac{2}{3}\right)^2} = \sqrt{\frac{2}{3}}$$

The shortest distance =  $\sqrt{\frac{2}{3}}$  units.

**Method 2:**



The equation of two given straight lines in the Cartesian form are  $\frac{x}{1} = \frac{y}{-1} = \frac{z}{1} \dots\dots\dots(i)$  and

$$\frac{x-1}{0} = \frac{y+1}{-2} = \frac{z}{1} \dots\dots\dots(ii)$$

The lines are not parallel as direction ratios are not proportional. Let  $P$  be a point on straight line (i) and  $Q$  be a point on straight line (ii) such that line  $PQ$  is perpendicular to both of the lines.

Let the coordinates of  $P$  be  $(\lambda, -\lambda, \lambda)$  and that of  $Q$  be  $(1, -2\mu - 1, \mu)$ ; where ' $\lambda$ ' and ' $\mu$ ' are scalars.

Then the direction ratios of the line  $PQ$  are  $(\lambda - 1, -\lambda + 2\mu + 1, \lambda - \mu)$

Since  $PQ$  is perpendicular to straight line (i), we have,

$$(\lambda - 1).1 + (-\lambda + 2\mu + 1).(-1) + (\lambda - \mu).1 = 0$$

$$\Rightarrow 3\lambda - 3\mu = 2 \dots\dots\dots(iii)$$

Since ,  $PQ$  is perpendicular to straight line (ii), we have

$$0.(\lambda - 1) + (-\lambda + 2\mu + 1).(-2) + (\lambda - \mu).1 = 0 \Rightarrow 3\lambda - 5\mu = 2 \dots\dots\dots(iv)$$

Solving (iii) and (iv), we get  $\mu = 0, \lambda = \frac{2}{3}$

**Therefore**, the **Coordinates** of  $P$  are  $\left(\frac{2}{3}, -\frac{2}{3}, \frac{2}{3}\right)$  and that of  $Q$  are  $(1, -1, 0)$

So, the required shortest distance is  $\sqrt{\left(1 - \frac{2}{3}\right)^2 + \left(-1 + \frac{2}{3}\right)^2 + \left(0 - \frac{2}{3}\right)^2} = \sqrt{\frac{2}{3}} \text{ units.}$

### Section -E

[This section comprises solution of 3 case- study/passage based questions of 4 marks each with two sub parts. Solution of the first two case study questions have three sub parts (i),(ii),(iii) of marks 1,1,2 respectively. Solution of the third case study question has two sub parts of 2 marks each.)

- 36 Let  $E_1, E_2, E_3$  be the events that Jayant, Sonia and Oliver processed the form, which are clearly pairwise mutually exclusive and exhaustive set of events.

$$\text{Then } P(E_1) = \frac{50}{100} = \frac{5}{10}, P(E_2) = \frac{20}{100} = \frac{1}{5} \text{ and } P(E_3) = \frac{30}{100} = \frac{3}{10}.$$

Also, let  $E$  be the event of committing an error.

We have,  $P(E | E_1) = 0.06, P(E | E_2) = 0.04, P(E | E_3) = 0.03$ .

- (i) The probability that Sonia processed the form and committed an error is given by

$$P(E \cap E_2) = P(E_2) \cdot P(E | E_2) = \frac{1}{5} \times 0.04 = 0.008.$$

1

- (ii) The total probability of committing an error in processing the form is given by

$$P(E) = P(E_1) \cdot P(E | E_1) + P(E_2) \cdot P(E | E_2) + P(E_3) \cdot P(E | E_3)$$

1

$$P(E) = \frac{50}{100} \times 0.06 + \frac{20}{100} \times 0.04 + \frac{30}{100} \times 0.03 = 0.047.$$

- (iii) The probability that the form is processed by Jayant given that form has an error is given by

$$P(E_1 | E) = \frac{P(E | E_1) \times P(E_1)}{P(E | E_1) \cdot P(E_1) + P(E | E_2) \cdot P(E_2) + P(E | E_3) \cdot P(E_3)}$$

1

$$= \frac{0.06 \times \frac{50}{100}}{0.06 \times \frac{50}{100} + 0.04 \times \frac{20}{100} + 0.03 \times \frac{30}{100}} = \frac{30}{47}$$

1

$$\text{Therefore, the required probability that the form is not processed by Jayant given that form has an error} = P(\overline{E_1} | E) = 1 - P(E_1 | E) = 1 - \frac{30}{47} = \frac{17}{47}.$$

1

$$\text{(iii) OR } \sum_{i=1}^3 P(E_i | E) = P(E_1 | E) + P(E_2 | E) + P(E_3 | E) = 1$$

1

Since, sum of the posterior probabilities is 1.

	$  \begin{aligned}  & (\text{We have, } \sum_{i=1}^3 P(E_i   E) = P(E_1   E) + P(E_2   E) + P(E_3   E)) \\  & = \frac{P(E \cap E_1) + P(E \cap E_2) + P(E \cap E_3)}{P(E)} \\  & = \frac{P((E \cap E_1) \cup (E \cap E_2) \cup (E \cap E_3))}{P(E)} \text{ as } E_i \& E_j; i \neq j, \text{ are mutually exclusive events} \\  & = \frac{P(E \cap (E_1 \cup E_2 \cup E_3))}{P(E)} = \frac{P(E \cap S)}{P(E)} = \frac{P(E)}{P(E)} = 1; 'S' \text{ being the sample space)  \end{aligned}  $	
37	<p>We have ,</p> $ \vec{F}_1  = \sqrt{6^2 + 0^2} = 6kN,  \vec{F}_2  = \sqrt{(-4)^2 + 4^2} = \sqrt{32} = 4\sqrt{2} kN,  \vec{F}_3  = \sqrt{(-3)^2 + (-3)^2} = \sqrt{18} = 3\sqrt{2} kN.$ <p>(i) Magnitude of force of Team <math>A = 6kN</math>. <span style="float: right;">1</span></p> <p>(ii) Since <math>\vec{a} + \vec{c} = 3(\hat{i} - \hat{j})</math> and <math>\vec{b} = -4(\hat{i} - \hat{j})</math> <span style="float: right;">1</span></p> <p>So, <math>\vec{b}</math> and <math>\vec{a} + \vec{c}</math> are unlike vectors having same intial point <span style="float: right;">1</span></p> <p>and <math> \vec{b}  = 4\sqrt{2}</math> &amp; <math> \vec{a} + \vec{c}  = 3\sqrt{2}</math> <span style="float: right;">1</span></p> <p>Thus <math> \vec{F}_2  &gt;  \vec{F}_1 + \vec{F}_3 </math> also <math>\vec{F}_2</math> and <math>\vec{F}_1 + \vec{F}_3</math> are unlike <span style="float: right;">1</span></p> <p>Hence B will win the game</p> <p>(iii) <math>\vec{F} = \vec{F}_1 + \vec{F}_2 + \vec{F}_3 = 6\hat{i} + 0\hat{j} - 4\hat{i} + 4\hat{j} - 3\hat{i} - 3\hat{j} = -\hat{i} + \hat{j}</math> <span style="float: right;">1</span></p> $\therefore  \vec{F}  = \sqrt{(-1)^2 + (1)^2} = \sqrt{2} kN. 1$ <p><b>OR</b> <span style="float: right;">1</span></p> $\vec{F} = -\hat{i} + \hat{j}$ $\therefore \theta = \pi - \tan^{-1}\left(\frac{1}{1}\right) = \pi - \frac{\pi}{4} = \frac{3\pi}{4}; \text{ where '}\theta\text{' is the angle made by the resultant force with the } +ve \text{ direction of the } x-\text{axis.} 1$	
38	$y = 4x - \frac{1}{2}x^2$ <p>(i) The rate of growth of the plant with respect to the number of days exposed to sunlight is given by <math>\frac{dy}{dx} = 4 - x</math>. <span style="float: right;">2</span></p> <p>(ii) Let rate of growth be represented by the function <math>g(x) = \frac{dy}{dx}</math>.</p>	

Now,  $g'(x) = \frac{d}{dx} \left( \frac{dy}{dx} \right) = -1 < 0$

$\Rightarrow g(x)$  decreases.

So the rate of growth of the plant decreases for the first three days.

Height of the plant after 2 days is  $y = 4 \times 2 - \frac{1}{2}(2)^2 = 6\text{cm.}$

1

1


# **Lorem ipsum**

**Lorem ipsum dolor sit amet, consectetur adipiscing elit. Nunc ac faucibus odio.**

Vestibulum neque massa, scelerisque sit amet ligula eu, congue molestie mi. Praesent ut varius sem. Nullam at porttitor arcu, nec lacinia nisi. Ut ac dolor vitae odio interdum condimentum. **Vivamus dapibus sodales ex, vitae malesuada ipsum cursus convallis. Maecenas sed egestas nulla, ac condimentum orci.** Mauris diam felis, vulputate ac suscipit et, iaculis non est. Curabitur semper arcu ac ligula semper, nec luctus nisl blandit. Integer lacinia ante ac libero lobortis imperdiet. *Nullam mollis convallis ipsum, ac accumsan nunc vehicula vitae.* Nulla eget justo in felis tristique fringilla. Morbi sit amet tortor quis risus auctor condimentum. Morbi in ullamcorper elit. Nulla iaculis tellus sit amet mauris tempus fringilla.

Maecenas mauris lectus, lobortis et purus mattis, blandit dictum tellus.

- **Maecenas non lorem quis tellus placerat varius.**
- *Nulla facilisi.*
- Aenean congue fringilla justo ut aliquam.
- Mauris id ex erat. Nunc vulputate neque vitae justo facilisis, non condimentum ante sagittis.
- Morbi viverra semper lorem nec molestie.
- Maecenas tincidunt est efficitur ligula euismod, sit amet ornare est vulputate.



In non mauris justo. Duis vehicula mi vel mi pretium, a viverra erat efficitur. Cras aliquam est ac eros varius, id iaculis dui auctor. Duis pretium neque ligula, et pulvinar mi placerat et. Nulla nec nunc sit amet nunc posuere vestibulum. Ut id neque eget tortor mattis tristique. Donec ante est, blandit sit amet tristique vel, lacinia pulvinar arcu. Pellentesque scelerisque fermentum erat, id posuere justo pulvinar ut. Cras id eros sed enim aliquam lobortis. Sed lobortis nisl ut eros efficitur tincidunt. Cras justo mi, porttitor quis mattis vel, ultricies ut purus. Ut facilisis et lacus eu cursus.

In eleifend velit vitae libero sollicitudin euismod. Fusce vitae vestibulum velit. Pellentesque vulputate lectus quis pellentesque commodo. Aliquam erat volutpat. Vestibulum in egestas velit. Pellentesque fermentum nisl vitae fringilla venenatis. Etiam id mauris vitae orci maximus ultricies.

## **Cras fringilla ipsum magna, in fringilla dui commodo**

**a.**

1	In eleifend velit vitae libero sollicitudin euismod.		
2	Cras fringilla ipsum magna, in fringilla dui commodo a.		
3	Aliquam erat volutpat.		
4	Fusce vitae vestibulum velit.		
5	Etiam vehicula luctus fermentum.		

Etiam vehicula luctus fermentum. In vel metus congue, pulvinar lectus vel, fermentum dui. Maecenas ante orci, egestas ut aliquet sit amet, sagittis a magna. Aliquam ante quam, pellentesque ut dignissim quis, laoreet eget est. Aliquam erat volutpat. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos himenaeos. Ut ullamcorper justo sapien, in cursus libero viverra eget. Vivamus auctor imperdiet urna, at pulvinar leo posuere laoreet. Suspendisse neque nisl, fringilla at iaculis scelerisque, ornare vel dolor. Ut et pulvinar nunc. Pellentesque fringilla mollis efficitur. Nullam venenatis commodo imperdiet. Morbi velit neque, semper quis lorem quis, efficitur dignissim ipsum. Ut ac lorem sed turpis imperdiet eleifend sit amet id sapien.

# **Lorem ipsum dolor sit amet, consectetur adipiscing elit.**

Nunc ac faucibus odio. Vestibulum neque massa, scelerisque sit amet ligula eu, congue molestie mi. Praesent ut varius sem. Nullam at porttitor arcu, nec lacinia nisi. Ut ac dolor vitae odio interdum condimentum. Vivamus dapibus sodales ex, vitae malesuada ipsum cursus convallis. Maecenas sed egestas nulla, ac condimentum orci. Mauris diam felis, vulputate ac suscipit et, iaculis non est. Curabitur semper arcu ac ligula semper, nec luctus nisl blandit. Integer lacinia ante ac libero lobortis imperdiet. Nullam mollis convallis ipsum, ac accumsan nunc vehicula vitae. Nulla eget justo in felis tristique fringilla. Morbi sit amet tortor quis risus auctor condimentum. Morbi in ullamcorper elit. Nulla iaculis tellus sit amet mauris tempus fringilla.

## **Maecenas mauris lectus, lobortis et purus mattis, blandit dictum tellus.**

Maecenas non lorem quis tellus placerat varius. Nulla facilisi. Aenean congue fringilla justo ut aliquam. Mauris id ex erat. Nunc vulputate neque vitae justo facilisis, non condimentum ante sagittis. Morbi viverra semper lorem nec molestie. Maecenas tincidunt est efficitur ligula euismod, sit amet ornare est vulputate.

In non mauris justo. Duis vehicula mi vel mi pretium, a viverra erat efficitur. Cras aliquam est ac eros varius, id iaculis dui auctor. Duis pretium neque ligula, et pulvinar mi placerat et. Nulla nec nunc sit amet nunc posuere vestibulum. Ut id neque eget tortor mattis tristique. Donec ante est, blandit sit amet tristique vel, lacinia pulvinar arcu. Pellentesque scelerisque fermentum erat, id posuere justo pulvinar ut. Cras id eros sed enim aliquam lobortis. Sed lobortis nisl ut eros efficitur tincidunt. Cras justo mi, porttitor quis mattis vel, ultricies ut purus. Ut facilisis et lacus eu cursus.

## **In eleifend velit vitae libero sollicitudin euismod.**

Fusce vitae vestibulum velit. Pellentesque vulputate lectus quis pellentesque commodo. Aliquam erat volutpat. Vestibulum in egestas velit. Pellentesque fermentum nisl vitae fringilla venenatis. Etiam id mauris vitae orci maximus ultricies. Cras fringilla ipsum magna, in fringilla dui commodo a.

Etiam vehicula luctus fermentum. In vel metus congue, pulvinar lectus vel, fermentum dui. Maecenas ante orci, egestas ut aliquet sit amet, sagittis a magna. Aliquam ante quam, pellentesque ut dignissim quis, laoreet eget est. Aliquam erat volutpat. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos himenaeos. Ut ullamcorper justo sapien, in cursus libero viverra eget. Vivamus auctor imperdiet urna, at pulvinar leo posuere laoreet. Suspendisse neque nisl, fringilla at iaculis scelerisque, ornare vel dolor. Ut et pulvinar nunc. Pellentesque fringilla mollis efficitur. Nullam venenatis commodo imperdiet. Morbi velit neque, semper quis lorem quis, efficitur dignissim ipsum. Ut ac lorem sed turpis imperdiet eleifend sit amet id sapien.

