

---

# Phys 434

## QUANTUM PHYSICS 3

University of Waterloo

Course notes by: TC Fraser  
Instructor: Anton Burkov

---

[tcfraser@tcfraser.com](mailto:tcfraser@tcfraser.com)

Version: 1.0

## Table Of Contents

	Page
<b>1 Review</b>	<b>4</b>
1.1 Discrete Spectrum . . . . .	4
1.2 Continuous Spectrum . . . . .	6
1.3 Infinitesimal Translations . . . . .	7
1.4 Transformations Between Position and Momentum Representations . . . . .	8
1.5 Time Dependence of Kets . . . . .	10

## Disclaimer

These notes are intended to be a reference for my future self (TC Fraser). If you the reader find these notes useful in any capacity, please feel free to use these notes as you wish, free of charge. However, I do not guarantee their complete accuracy and mistakes are likely present. If you notice any errors please email me at **[tcfraser@tcfraser.com](mailto:tcfraser@tcfraser.com)**, or contribute directly at **<https://github.com/tcfraser/course-notes>**. If you are the professor of this course (Anton Burkov) and you've managed to stumble upon these notes and would like to make large changes or additions, email me please.

Latest versions of all my course notes are available at **[www.tcfraser.com/coursenotes](http://www.tcfraser.com/coursenotes)**.

# 1 Review

## 1.1 Discrete Spectrum

States in quantum mechanics are vectors in Hilbert space  $\mathcal{H}$ . In Dirac notation, states are denoted as *kets*  $|\psi\rangle$ . Observables in quantum mechanics are operators  $A : \mathcal{H} \rightarrow \mathcal{H}$  such that  $|\psi\rangle \mapsto A|\psi\rangle$ . Every operator  $A$  has a set of eigenkets  $\{|a'\rangle\}$ ,

$$A|a'\rangle = a'|a'\rangle$$

The eigenvalue corresponding to the eigenket  $|a'\rangle$  is denoted  $a' \in \mathbb{R}$ . The dual Hilbert space will be called the bra space and elements of the bra space will be denoted with a ket space  $\langle\varphi|$ .

We will denote the *inner product* (scalar product) to be  $\langle\varphi|\psi\rangle$ . By definition,

$$\langle\varphi|\psi\rangle = \langle\psi|\varphi\rangle^*$$

$$\langle\psi|\psi\rangle = \|\psi\|^2 \geq 0$$

Every state in the Hilbert space can be normalized,

$$|\tilde{\psi}\rangle = \frac{1}{\sqrt{\langle\psi|\psi\rangle}}|\psi\rangle$$

In doing so, we have,

$$\langle\tilde{\psi}|\tilde{\psi}\rangle = \frac{\langle\psi|\psi\rangle}{\langle\psi|\psi\rangle} = 1$$

Evidently, if we have that  $\langle\varphi|\psi\rangle = \langle\psi|\varphi\rangle$ , then  $\langle\varphi|\psi\rangle$  must be real. A bra  $\langle\varphi|$  and ket  $|\psi\rangle$  are said to be *orthogonal* if  $\langle\varphi|\psi\rangle = 0$ .

The dual of  $A|\psi\rangle$  is  $\langle\psi|A^\dagger$ . Where  $A^\dagger$  is the Hermitian conjugate (adjoint) of  $A$ . We can act on the ket  $A|\psi\rangle$  with the bra  $\langle\varphi|$  and obtain,

$$\langle\varphi|A|\psi\rangle = \langle\psi|A^\dagger|\varphi\rangle^*$$

The operator  $A$  is *Hermitian* if and only if  $A = A^\dagger$ .

If  $A$  is a Hermitian operator, then  $A$ 's eigenvalues and eigenkets have particularly nice properties. Let  $(a', |a'\rangle)$  and  $(a'', |a''\rangle)$  be two eigen-pairs.

$$A|a'\rangle = a'|a'\rangle \tag{1.1}$$

$$A|a''\rangle = a''|a''\rangle \tag{1.2}$$

Let  $\langle\varphi|$  be an arbitrary bra. By eq. (1.2) we have that,

$$\langle\varphi|A|a''\rangle = a''\langle\varphi|a''\rangle$$

The adjoint to this equation yields,

$$\langle a''|A|\varphi\rangle^* = a''\langle a''|\varphi\rangle^*$$

Conjugating each term,

$$\langle a''|A|\varphi\rangle = a''^*\langle a''|\varphi\rangle \tag{1.3}$$

Since eq. (1.3) is true for an arbitrary  $\langle\varphi|$ , it must be that

$$\langle a''|A = a''^*\langle a''| \tag{1.4}$$

Combining eqs. (1.4) and (1.1), and recognizing that  $A$  is Hermitian,

$$\underbrace{\langle a''|A|a'\rangle - \langle a''|A^\dagger|a'\rangle}_0 = a'\langle a''|a'\rangle = a''^*\langle a''|a'\rangle$$

Therefore,

$$(a' - a''^*)\langle a''|a'\rangle = 0 \quad (1.5)$$

As an example, we can chose  $|a''\rangle = |a'\rangle$  to see that

$$(a' - a'^*)\langle a'|a'\rangle = 0 \implies a' = a'^*$$

Therefore all eigenvalues of Hermitian operators are always real. Since the spectrum of an operator represents all physical observables, this observation is in agreement with the fact that all physical quantities are real-valued.

Moreover returning to eq. (1.5) we can consider  $|a'\rangle$  and  $|a''\rangle$  to be different eigenkets that are non-degenerate (their eigenvalues differ). Then be eq. (1.5),

$$\langle a''|a'\rangle = 0$$

Therefore eigenkets of Hermitian operators are orthogonal (or can at least be orthogonalized). Since the norm of an eigenket is arbitrary, we will hence forth assert that all eigenkets are normalized. Each of these properties can be summarized with a Kronecker delta.

$$\langle a|a'\rangle = \delta_{a,a'} \quad (1.6)$$

In summary, the set of eigenkets of any Hermitian operator forms a complete orthonormal set of states. Effectively, the set of eigenkets form a basis for the Hilbert space. Consequently, we can write any ket  $|\psi\rangle$  in terms of the eigenkets for any Hermitian operator  $A$

$$|\psi\rangle = \sum_{a'} C_{a'} |a'\rangle \quad (1.7)$$

Where  $C_{a'} \in \mathbb{C}$  are uniquely defined through acting with the dual eigenket  $\langle a''|$ ,

$$\langle a''|\psi\rangle = \sum_{a'} C_{a'} \langle a''|a'\rangle = \sum_{a'} C_{a'} \delta_{a'',a'} = C_{a''} \implies C_{a'} = \langle a'|\psi\rangle \quad (1.8)$$

Physically, the coefficient  $C_{a'}$  is called a *probability amplitude*. When a given system is in state  $|\psi\rangle$ , the probability of measuring the value  $a'$  when making the observation or measurement  $A$  is given by the square modulus of  $C_{a'}$ ,

$$P_A(a') = |\langle a'|\psi\rangle|^2$$

We now have the luxury of re-writing eq. (1.7) as a spectral decomposition,

$$|\psi\rangle = \sum_{a'} |a'\rangle \langle a'|\psi\rangle \quad (1.9)$$

Since  $|\psi\rangle$  is *arbitrary*, we obtain a closure relation (otherwise known as the resolution of identity).

$$\sum_{a'} |a'\rangle \langle a'| = \mathbb{I} \quad (1.10)$$

We define the projection operator  $\Lambda_{a'} = |a'\rangle \langle a'|$ .

$$\Lambda_{a''} |\psi\rangle = |a''\rangle \langle a''|\psi\rangle = \sum_{a'} |a'\rangle \underbrace{\langle a'|a''\rangle}_{\delta_{a',a''}} \langle a''|\psi\rangle = \langle a''|\psi\rangle |a''\rangle$$

As such,  $\Lambda_a$  *projects*  $|\psi\rangle$  into the direction of  $|a\rangle$ . Using the closure operation eq. (1.10) and the spectral decomposition of a ket eq. (1.9) one can recover the spectral decomposition of an operator  $A$ . For each eigenket  $|a'\rangle$ , multiply eq. (1.1) by  $\langle a'|$ ,

$$A|a'\rangle \langle a'| = a'|a'\rangle \langle a'|$$

And summing over all eigenkets,

$$A = \sum_{a'} a' |a'\rangle \langle a'|$$

Additionally consider another operator  $B$ ,

$$B = \mathbb{I} \cdot B \cdot \mathbb{I} = \sum_{a', a''} |a''\rangle \langle a''| B |a'\rangle \langle a'|$$

Where  $\langle a''|B|a'\rangle$  can be interpreted as a matrix indexed by  $|a''\rangle$  and  $|a'\rangle$ ,

$$\langle a''|B|a'\rangle = B_{a'', a'}$$

Where refer to  $B_{a'', a'}$  as the matrix elements of an operator  $B$  with respect to the a complete orthonormal set of eigenstates of a Hermitian operator  $A$ . The entries in  $B_{a'', a'}$  have the following property,

$$\langle a''|B|a'\rangle = \langle a'|B^\dagger|a''\rangle^*$$

Therefore the matrix that corresponds to  $B^\dagger$  is the complex conjugate transposed of the matrix corresponding to  $B$ .

## 1.2 Continuous Spectrum

Of course, there exists operators with non-discrete spectrum. We will now generalize to operators with continuous spectrum. The two most important of such operators are position and momentum. Let  $|\vec{x}'\rangle$  a position eigenket corresponding to the state of a particle at position  $\vec{x}'$  in space. Let  $\vec{x}$  be the position operator defined as,

$$\vec{x}|\vec{x}'\rangle = \vec{x}'|\vec{x}'\rangle$$

It is important not to get confused about notation:

- $\vec{x}$  – Position operator
- $\vec{x}'$  – Position eigenket

The wave function  $\psi(\vec{x}')$  is the probability amplitude to find a particle in a state  $|\psi\rangle$  at position  $\vec{x}'$  and is defined as,

$$\langle \vec{x}'|\psi\rangle = \psi(\vec{x}')$$

We also have the ability to generalize eq. (1.6) to a continuous spectrum. The continuous generalization of the Kronecker delta is the Dirac delta function.

$$\langle \vec{x}'|\vec{x}''\rangle = \delta(\vec{x}' - \vec{x}'')$$

Where  $\delta\vec{x}'$  is defined as,

$$\int_{\mathbb{R}^3} d^3x' f(\vec{x}') \delta(\vec{x}') = f(\vec{0})$$

Where  $f(\vec{x}') : \mathbb{R}^3 \rightarrow \mathbb{R}$  is a function on  $\mathbb{R}^3$ .

The closure relation becomes,

$$\mathbb{I} = \int_{\mathbb{R}^3} d^3x' |\vec{x}'\rangle \langle \vec{x}'|$$

Therefore we have that,

$$|\psi\rangle = \mathbb{I} \cdot |\psi\rangle = \int_{\mathbb{R}^3} d^3x' |\vec{x}'\rangle \langle \vec{x}'|\psi\rangle$$

Now let  $|\phi\rangle$  be another space in the same Hilbert space as  $|\psi\rangle$ ,

$$\begin{aligned}\langle\phi|\psi\rangle &= \int_{\mathbb{R}^3} d^3x' \langle\phi|\vec{x}'\rangle \langle\vec{x}'|\psi\rangle \\ &= \int_{\mathbb{R}^3} d^3x' \langle\vec{x}'|\phi\rangle^* \langle\vec{x}'|\psi\rangle \\ &= \int_{\mathbb{R}^3} d^3x' \phi(\vec{x}')^* \psi(\vec{x}')\end{aligned}$$

### 1.3 Infinitesimal Translations

The operator of infinitesimal translations  $T$  is defined as,

$$T(d\vec{x}')|\vec{x}'\rangle = |\vec{x}' + d\vec{x}'\rangle$$

Where  $d\vec{x}'$  is an infinitesimally small vector. Acting on an arbitrary state  $|\psi\rangle$ ,

$$\begin{aligned}T(d\vec{x}')|\psi\rangle &= T(d\vec{x}') \left\{ \int_{\mathbb{R}^3} d^3x' |\vec{x}'\rangle \langle\vec{x}'|\psi\rangle \right\} \\ &= \int_{\mathbb{R}^3} d^3x' |\vec{x}' + d\vec{x}'\rangle \langle\vec{x}'|\psi\rangle \\ &= \int_{\mathbb{R}^3} d^3x' |\vec{x}'\rangle \langle\vec{x}' - d\vec{x}'|\psi\rangle \quad \vec{x}' \mapsto \vec{x}' - d\vec{x}'\end{aligned}$$

Next without loss of generality, let  $|\psi\rangle$  be normalized  $\langle\psi|\psi\rangle = 1$ . Moreover, we may let  $T(d\vec{x}')|\psi\rangle$  be normalized as well.

$$\langle\psi|T^\dagger(d\vec{x}')T(d\vec{x}')|\psi\rangle \quad (1.11)$$

If we wish for eq. (1.11) to hold for all states  $|\psi\rangle$ , it must be that  $T(d\vec{x}')$  is *unitary*.

$$T^\dagger(d\vec{x}')T(d\vec{x}') = \mathbb{I} \implies T^\dagger(d\vec{x}') = T^{-1}(d\vec{x}') \quad (1.12)$$

Another desired property of translations  $T(d\vec{x}')$  is that they are additive,

$$T(d\vec{x}')T(d\vec{x}'') = T(d\vec{x}' + d\vec{x}'') \quad (1.13)$$

Consequently,

$$T^{-1}(d\vec{x}') = T(-d\vec{x}') \quad T(\vec{0}) = \mathbb{I}$$

All of the above properties are satisfied if,

$$T(d\vec{x}') = \mathbb{I} - i\vec{K} \cdot d\vec{x}'$$

Where  $\vec{K} = (K_x, K_y, K_z)$  is a vector operator that is Hermitian ( $\vec{K}^\dagger = \vec{K}$ ) to be determined. First we demonstrate that such a  $T(d\vec{x}')$  is unitary (eq. (1.12)),

$$\begin{aligned}T^\dagger(d\vec{x}')T(d\vec{x}') &= \left(\mathbb{I} + i\vec{K}^\dagger \cdot d\vec{x}'\right) \left(\mathbb{I} - i\vec{K} \cdot d\vec{x}'\right) \\ &= \mathbb{I} + \underbrace{i\vec{K}^\dagger \cdot d\vec{x}' - i\vec{K} \cdot d\vec{x}'}_0 + \mathcal{O}(|d\vec{x}'|^2) \xrightarrow{0} \\ &= \mathbb{I}\end{aligned}$$

Next we demonstrate additivity (eq. (1.13)),

$$T(d\vec{x}'')T(d\vec{x}') = \left(\mathbb{I} - i\vec{K} \cdot d\vec{x}''\right) \left(\mathbb{I} - i\vec{K} \cdot d\vec{x}'\right)$$

$$\begin{aligned}
&= \mathbb{I} - i\vec{K} \cdot d\vec{x}'' - i\vec{K} \cdot d\vec{x}' + \mathcal{O}(|d\vec{x}'|^2) \rightarrow 0 \\
&= \mathbb{I} - i\vec{K} \cdot (d\vec{x}'' + d\vec{x}') \\
&= T(d\vec{x}'' + d\vec{x}')
\end{aligned}$$

In order to illuminate the specific form of  $\vec{K}$ , we calculate the commutator  $[\vec{x}, T(d\vec{x}')] ,$

$$\begin{aligned}
[\vec{x}, T(d\vec{x}')] &= [\vec{x}, \mathbb{I} - i\vec{K} \cdot d\vec{x}'] \\
&= -i\vec{x}\vec{K}d\vec{x}' + i\vec{K}d\vec{x}'\vec{x}
\end{aligned}$$

Choose  $d\vec{x}' = dx'\hat{x}_j$  and  $\vec{K} \cdot \hat{x}_j = K_j$  where  $\hat{x}_j$  is the unit vector in the direction of one of the basis vectors.

$$[\vec{x}, T(d\vec{x}')] = -i\vec{x}\vec{K}d\vec{x}' + i\vec{K}d\vec{x}'\vec{x}$$

Moreover  $[X_i, K_j] = 0$  whenever  $i \neq j$ .

$$-ix_jK_j + iK_jx_j = 1 \implies -i[x_j, K_j] = 1 \implies [x_j, K_j] = i$$

Therefore,

$$[x_i, K_j] = i\delta_{ij} \implies \vec{K} = \frac{1}{\hbar}\vec{p}$$

Where  $\vec{p}$  is the generator of infinitesimal translations,

$$[x_i, p_j] = i\hbar\delta_{ij}$$

Such that,

$$T(d\vec{x}') = 1 - \frac{i}{\hbar}\vec{p} \cdot d\vec{x}'$$

## 1.4 Transformations Between Position and Momentum Representations

Calculate for a 1D system,

$$\begin{aligned}
T(\Delta x')|\psi\rangle &= \left(1 - \frac{i}{\hbar}p\Delta x'\right)|\psi\rangle \\
&= \int_{\mathbb{R}} dx' \left(1 - \frac{i}{\hbar}p\Delta x'\right)|x'\rangle\langle x'|\psi\rangle \\
&= \int_{\mathbb{R}} dx' T(\Delta x')|x'\rangle\langle x'|\psi\rangle \\
&= \int_{\mathbb{R}} dx' |x' + \Delta x'\rangle\langle x'|\psi\rangle \\
&= \int_{\mathbb{R}} dx' |x'\rangle\langle x' - \Delta x'|\psi\rangle
\end{aligned}$$

Examine  $\langle x' - \Delta x'|\psi\rangle$ ,

$$\langle x' - \Delta x'|\psi\rangle \approx \langle x'|\psi\rangle - \Delta x' \frac{\partial}{\partial x'} \langle x'|\psi\rangle$$

Therefore,

$$\begin{aligned}
T(\Delta x')|\psi\rangle &= \int_{\mathbb{R}} dx' |x'\rangle \left[ \langle x'|\psi\rangle - \Delta x' \frac{\partial}{\partial x'} \langle x'|\psi\rangle \right] \\
&= |\psi\rangle - \Delta x' \int_{\mathbb{R}} dx' |x'\rangle \left[ \frac{\partial}{\partial x'} \langle x'|\psi\rangle \right]
\end{aligned}$$



Which in turn implies that,

$$p|\psi\rangle = \int_{\mathbb{R}} dx' |x'\rangle \left( -i\hbar \frac{\partial}{\partial x'} \right) \langle x'|\psi\rangle$$

Since a given ket  $|\psi\rangle$  can be written in *any* basis or representation, we can transform  $|\psi\rangle$  in the momentum basis. Recall the momentum eigenkets form a complete orthonormal set of states,

$$\vec{p}|\vec{p}'\rangle = \vec{p}'|\vec{p}'\rangle \quad \langle \vec{p}'|\vec{p}''\rangle = \delta(\vec{p}' - \vec{p}'')$$

Moreover we have the resolution of identity,

$$\mathbb{I} = \int d^3p' |\vec{p}'\rangle \langle \vec{p}'|$$

Therefore we have that,

$$|\psi\rangle = \mathbb{I} \cdot |\psi\rangle = \int d^3p' |\vec{p}'\rangle \langle \vec{p}'|\psi\rangle$$

So we define the wavefunction in momentum representation  $\langle \vec{p}'|\psi\rangle = \psi(\vec{p}')$ . We will now discover how to transform from  $\psi(\vec{p}')$  to  $\psi(\vec{x}')$  in 1D. By definition we have that,

$$\langle x'|p|\psi\rangle = -i\hbar \frac{\partial}{\partial x'} \langle x'|\psi\rangle$$

We now choose  $|\psi\rangle = |p'\rangle$ ,

$$\langle x'|p|p'\rangle = -i\hbar \frac{\partial}{\partial x'} \langle x'|p'\rangle$$

But  $|p'\rangle$  is an eigenket of  $p$ ,

$$p'\langle x'|p'\rangle = -i\hbar \frac{\partial}{\partial x'} \langle x'|p'\rangle$$

Therefore we have a differential equation for  $f(x') = \langle x'|p'\rangle$ ,

$$p'f(x') = -i\hbar \frac{\partial f}{\partial x'} \tag{1.14}$$

Which has the well known solution,

$$f(x') = \langle x'|p'\rangle = Ne^{\frac{i}{\hbar}p'x'} \tag{1.15}$$

Where  $N$  is an arbitrary constant. To confirm eq. (1.14) check  $\frac{\partial f}{\partial x'}$

$$\frac{\partial f}{\partial x'} = N \frac{i}{\hbar} p' e^{\frac{i}{\hbar}p'x'} = \frac{i}{\hbar} p' f(x')$$

As a quick trick notice that,

$$\langle x'|x''\rangle = \delta(x' - x'') = \langle x'|\mathbb{I}|x''\rangle = \int dp' \langle x'|p'\rangle \langle p'|x''\rangle$$

Substitute in eq. (1.15),

$$\delta(x' - x'') = N^2 \int dp' e^{\frac{i}{\hbar}p'x'} e^{-\frac{i}{\hbar}p'x''} = N^2 \int dp' e^{\frac{i}{\hbar}p'(x' - x'')} \tag{1.16}$$

Recall a integral representation of the Dirac-delta function,

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ikx} dk = \delta(x) \tag{1.17}$$

Comparing eqs. (1.16) and (1.17) (and using  $\mu_0 \pi a^2 n \dot{I}_s$ ) we have that,

$$N^2 = \frac{1}{2\pi\hbar} \implies N = \frac{1}{\sqrt{2\pi\hbar}}$$

This we have that,

$$\langle x' | p' \rangle = \frac{1}{\sqrt{2\pi\hbar}} e^{\frac{i}{\hbar} p' x'} \quad (1.18)$$

Which refers to the usual plane wave wavefunction. Alternatively, one can obtain this result from the Schrodinger Equation  $H\psi = E\psi$  and using a free Hamiltonian  $H = \frac{p^2}{2m} = -\frac{\hbar^2 d^2}{2mdx^2}$ . Generalizing eq. (1.18) to more than one dimension gives (say 3 dimensions),

$$\langle \vec{x}' | \vec{p}' \rangle = \frac{1}{(2\pi\hbar)^{3/2}} e^{\frac{i}{\hbar} \vec{p}' \cdot \vec{x}'}$$

This result allows us to convert from the momentum representation  $\psi(\vec{p}')$  to the position representation  $\psi(\vec{x}')$  and backward,

$$\begin{aligned} \langle x' | \psi \rangle &= \int dp' \langle x' | p' \rangle \langle p' | \psi \rangle \\ &= \frac{1}{\sqrt{2\pi\hbar}} \int dp' e^{\frac{i}{\hbar} p' x'} \langle p' | \psi \rangle \end{aligned}$$

Analogously we can rotate state space to give,

$$\langle p' | \psi \rangle = \frac{1}{\sqrt{2\pi\hbar}} \int dx' e^{-\frac{i}{\hbar} p' x'} \langle x' | \psi \rangle$$

Which is nothing more than the **(Inverse) Fourier Transform**.

## 1.5 Time Dependence of Kets

Let  $|\psi, t_0; t\rangle$  be the state which is  $|\psi, t_0\rangle \equiv |\psi\rangle$  at time  $t_0$  which becomes a different state  $|\psi, t_0; t\rangle$  at a later time  $t > t_0$ . Of course, there must exist an operator that transforms initial states  $|\psi, t_0\rangle$  into final state  $|\psi, t_0; t\rangle$ . Let  $U(t, t_0)$  be this unknown operator,

$$|\psi, t_0; t\rangle = U(t, t_0) |\psi, t_0\rangle$$

Which we will now discover. To do so, we will demand some properties of  $U(t, t_0)$ . Consider some physical quantity with corresponding operator  $A$  with eigenkets  $|a'\rangle$  and eigenvalues  $a'$ . Then we can write  $|\psi, t_0\rangle$  in terms of the orthonormal set of states defined by  $\{|a'\rangle\}$ <sup>1</sup>,

$$|\psi, t_0\rangle = \sum_{a'} C_{a'}(t_0) |a'\rangle$$

Analogously at time  $t$ ,

$$|\psi, t_0; t\rangle = \sum_{a'} C_{a'}(t) |a'\rangle$$

The coefficients  $C_{a'}$  are determined by eq. (1.8),

$$\langle \psi, t_0 | \psi, t_0 \rangle = \sum_{a', a''} C_{a'}^*(t_0) C_{a''}(t_0) \underbrace{\langle a' | a'' \rangle}_{\delta_{a', a''}}$$

<sup>1</sup>We will also assign all of the time evolution to the coefficients  $C_{a'} = C_{a'}(t_0)$

Notice that normalization dictates that  $\sum_{a'} |C_{a'}(t_0)|^2 = 1$ . Therefore we can interpret  $|C_{a'}(t_0)|$  as the probability that a measurement of a physical system  $A$  gives  $a'$ . Analogously at later times  $t$ ,

$$\sum_{a'} |C_{a'}(t)|^2 = 1$$

Therefore it must be that  $U(t_0; t)$  is unitary.

$$\langle \psi, t_0; t | \psi, t_0; t \rangle = \langle \psi, t_0 | U^\dagger(t, t_0) U(t, t_0) | \psi, t_0 \rangle = \langle \psi, t_0 | \psi, t_0 \rangle$$

Which holds for all  $|\psi\rangle$ , thus,

$$U^\dagger(t, t_0) U(t, t_0) = \mathbb{I} \implies U^\dagger(t, t_0) = U^{-1}(t, t_0) \quad (1.19)$$

Another desired property of the time evolution operator  $U(t; t_0)$  is *composition*. For  $t_2 > t_1 > t_0$ ,

$$U(t_2, t_0) = U(t_2, t_1) U(t_1, t_0) \quad (1.20)$$

It turns out that eqs. (1.19) and (1.20) uniquely characterize  $U(t, t_0)$ . Consider an infinitesimal time evolution operator  $U(t_0 + dt, t_0)$ . We now prove that,

$$U(t_0 + dt, t_0) = \mathbb{I} - i\Omega dt$$

Where  $\Omega = \Omega^\dagger$  is an unknown Hermitian operator. Consider,

$$\begin{aligned} U^\dagger(t_0 + dt, t_0) U(t_0 + dt, t_0) &= (\mathbb{I} + i\Omega^\dagger dt)(\mathbb{I} - i\Omega dt) \\ &= \mathbb{I} + i \left( \underbrace{\Omega^\dagger - \Omega}_0 \right) dt + \mathcal{O}(dt^2) \xrightarrow{0} \mathbb{I} \\ &= \mathbb{I} \end{aligned}$$

Thus satisfying unitary properties. Next examine composition,

$$\begin{aligned} U(t_0 + dt_1 + dt_2, t_0 + dt_1) U(t_0 + dt_1, t_0) &= (\mathbb{I} - i\Omega dt_2)(\mathbb{I} - i\Omega dt_1) \\ &= \mathbb{I} - i\Omega(dt_1 + dt_2) + \mathcal{O}(dt_1 dt_2) \\ &= U(t_0 + dt_1 + dt_2, t_0) \end{aligned}$$

By dimensional analysis,  $\Omega$  needs to have dimensions of inverse time or *frequency*. Of course the energy and frequency of a system are related by  $E = \hbar\omega$ . We conclude that,

$$\Omega = \frac{1}{\hbar} H$$

Where  $H$  is the usual Hamiltonian operator. This result is analogous to  $\vec{K} = \frac{1}{\hbar} \vec{p}$ . We have that,

$$U(t_0 + dt, t_0) = \mathbb{I} - \frac{i}{\hbar} H dt$$

Using this result, we will recover the Schrodinger equation. Consider the difference of two time evolution operators,

$$U(t + dt, t_0) - U(t, t_0) = -\frac{i}{\hbar} H dt U(t, t_0)$$

Dividing both sides by  $dt$  and taking a limit,

$$\lim_{dt \rightarrow 0} \frac{U(t + dt, t_0) - U(t, t_0)}{dt} = -\frac{i}{\hbar} H U(t, t_0)$$

One obtains,

$$\frac{\partial}{\partial t} U(t, t_0) = -\frac{i}{\hbar} H U(t, t_0)$$

Which is identical to the Schrodinger equation for operators,

$$i\hbar \frac{\partial}{\partial t} U(t, t_0) = H U(t, t_0)$$

We can also recover the more familiar Schrodinger equation for states through the following process,

$$i\hbar \frac{\partial}{\partial t} U(t, t_0) |\psi, t_0\rangle = H U(t, t_0) |\psi, t_0\rangle$$

$$i\hbar \frac{\partial}{\partial t} |\psi, t_0; t\rangle = H |\psi, t_0; t\rangle$$

Now multiply by  $\langle \vec{x}' |$ ,

$$i\hbar \frac{\partial}{\partial t} \langle \vec{x}' | \psi, t_0; t \rangle = \langle \vec{x}' | H | \psi, t_0; t \rangle$$

Inserting resolution of identity,

$$\langle \vec{x}' | H | \psi, t_0; t \rangle = \int d^3x'' \langle \vec{x}' | H | \vec{x}'' \rangle \langle \vec{x}'' | \psi, t_0; t \rangle$$

Where  $\langle \vec{x}' | H | \vec{x}'' \rangle$  is the Hamiltonian in terms of the position basis,

$$\langle \vec{x}' | H | \vec{x}'' \rangle = -\frac{\hbar^2}{2m} \vec{\nabla} \delta(\vec{x}' - \vec{x}'')$$