

---

# Phys 442

## ELECTRICITY & MAGNETISM 3

University of Waterloo

Course notes by: TC Fraser  
Instructor: Chris O'Donovan

---

[tcfrazer@tcfrazer.com](mailto:tcfrazer@tcfrazer.com)

Version: 1.0

## Table Of Contents

	Page
<a href="#">1 Coordinates and Symmetry</a>	<a href="#">4</a>
<a href="#">2 First Assignment?</a>	<a href="#">4</a>

## Disclaimer

These notes are intended to be a reference for my future self (TC Fraser). If you the reader find these notes useful in any capacity, please feel free to use these notes as you wish, free of charge. However, I do not guarantee their complete accuracy and mistakes are likely present. If you notice any errors please email me at **[tcfraser@tcfraser.com](mailto:tcfraser@tcfraser.com)**, or contribute directly at **<https://github.com/tcfraser/course-notes>**. If you are the professor of this course (Chris O'Donovan) and you've managed to stumble upon these notes and would like to make large changes or additions, email me please.

Latest versions of all my course notes are available at **[www.tcfraser.com/coursenotes](http://www.tcfraser.com/coursenotes)**.

## 1 Coordinates and Symmetry

A clever choice of coordinates systems typically makes solving a problem considerably easier. Mathematically, this is due to *Noether's Theorem*. A typical three dimensional Lagrangian will have three dependent generalized coordinates  $L = L(x, y, z) = L(s, \theta, \zeta) = \dots$ . However, if one can identify generalized coordinates  $q$  that make the Lagrangian invariant  $\frac{\partial L}{\partial q} = 0$ , then the *Euler-Lagrange* equations are considerably similar,

$$\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{q}} \right) - \frac{\partial L}{\partial q} = 0 \implies \frac{\partial L}{\partial \dot{q}} = \text{const.} \implies L \propto \dot{q}$$

As such, the number of equations that remain to be solved has been reduced.

## 2 First Assignment?

**A1.1:** Use cylindrical coordinates with  $\zeta$  along the axis of the cable,

$$V(\zeta) = \frac{1}{4\pi\epsilon_0} \int_C \frac{d\rho}{z}$$

Where  $\vec{z} = \vec{r} - \vec{r}'$ ,  $\vec{r}'$  is the source point and  $\vec{r}$  is the field point. The entire cylinder is the set of all source points  $\vec{r}'$  that are contained inside  $|\vec{r}'| \leq R$ .

$$\begin{aligned} \vec{r} &= \zeta \hat{\zeta} \\ \vec{r}' &= s' \hat{s}' + \zeta' \hat{\zeta} \end{aligned}$$

$$V(\zeta) = \frac{\rho}{4\pi\epsilon_0} \int_C \frac{dV}{|\vec{r} - \vec{r}'|}$$

Where  $dV = s ds d\theta d\zeta$ . One can then find the electric *field* by doing  $\vec{E} = -\vec{\nabla} V = E_{\zeta} \hat{\zeta} = -\frac{\partial V}{\partial \zeta} \hat{\zeta}$

**A1.2:**

Between the two conductors, there will be a radial electric field  $\vec{E} = E(s)\hat{s}$  and parallel magnetism field  $\vec{B} = B(s)\hat{\zeta}$ . Outside the two conductors, there will be no electric or magnetic field.

$$\begin{aligned} E_{\text{vac}}^{\parallel} &= 0 \\ E_{\text{vac}}^{\perp} &= \frac{\sigma}{\epsilon_0} \\ \vec{\nabla} \cdot \vec{E} &= \frac{\rho}{\epsilon_0} \end{aligned}$$

Todo (TC Fraser): Figure out O'Donovan