To: The Editor, *The Open Journal of Astrophysics*
From: Sanskar Sontakke (Independent Researcher)
Date: October 2026
Subject: New submission, with the record of a pre-submission review and the revisions it required: *Time-Domain Imaging of Rotating, Cloud-Covered Exoplanets with the Solar Gravitational Lens: Reconstruction Algorithms, Observing-Cadence Requirements, and Mission Targets*

Dear Editor,

Please find enclosed the manuscript, submitted for consideration as a new submission. No referee report on it has been issued by *OJA* or any other journal, and no editorial decision is quoted or implied anywhere in this letter. What follows instead is the record of a structured *pre-submission review* that I commissioned and ran over the manuscript in the *OJA* referee-report format before submitting. I include it because that review, rather than the item's wording, changed the paper's content: it identified claim-attribution inconsistencies, unfulfilled forward references, omitted analytical diagnostics, and statistical reporting gaps, and acting on it re-ran the pipeline end to end, which is the reason the enclosed numbers can be checked rather than trusted.

Rather than treating those findings as cosmetic or rhetorical issues, I addressed every item in the manuscript and the repository before submitting. Every reported number now matches the archived calculations exactly, the causal attributions reflect the true experimental contrasts, and negative and structural failure modes (such as surface–cloud phase locking and the collapse of single-pixel scalar photometry at Earth-like cloud coverage) are given their rightful prominence in the Abstract and Conclusions.

Below, I summarize the primary structural improvements and then respond item by item to all **ten major** and **nine minor** comments of that internal review, quoting each comment as it was written down.

Executive Summary of Substantive Revisions Made Before Submission

1. **Causal Attribution of Algorithmic Gain (Major items 1 and 4):** I have corrected the causal attribution of the matched ablation gain. The factor-by-factor comparison in Table 6 clearly demonstrates that essentially the entire algorithmic gain ($\Delta r = +0.054 \pm 0.006$ over white-noise GLS; $+0.0568 \pm 0.0046$ in Row D vs. Row A, $p_t = 5.9 \times 10^{-7}$) arises from per-dwell-slot nuisance profiling, not from the Ornstein–Uhlenbeck (OU) temporal covariance. Adding the OU covariance at this cadence slightly reduces correlation ($\Delta r = -0.0032 \pm 0.0019$, Row F vs. Row D). I explicitly defined baseline B3 in §4.6 and linked it to Table 6 Row A. Furthermore, I replaced the heuristic slot deflation and the "6.25% dimension reduction preserves 93.75% of information" claim with the exact anchored profile likelihood (Eqs. 10–12), derived in closed form as a rank-one correction to the block-diagonal normal equations ($A_{\text{anchored}} = A_{\text{free}} + \mathbf{q}\mathbf{q}^{\mathsf{T}}/S$, $\mathbf{b}_{\text{anchored}} = \mathbf{b}_{\text{free}} + \mathbf{q}m/S$).

2. **Resolution of the Spin Identifiability Degeneracy (Major item 2):** I withdrew the claim that residual spin misalignments can be refined in flight by minimizing photometric residuals. I proved analytically (Eqs. 10–11 in the profile-likelihood derivation) and numerically (§5.9) that the whitened profile likelihood χ^2 is invariant to the assumed initial spin phase ($\leq 9.2 \times 10^{-12}$ relative spread across $\pm 10^{\circ}$), because an initial phase shift is absorbed into the map's longitude origin. I regenerated Figure 6b with an explicit grid and normalized objective, and showed that while the spin state is not refinable via photometric residual grid search, it remains observable through reconstruction fidelity (r peaks at the true period on 4/4 seeds). In §5.1, I rebuilt the static control on common observable support and identical parameterization using the B3 white-noise GLS estimator ($r = 0.4074 \pm 0.0254$), removing the "virtually zero information loss" claim

and explaining that identical pose-passes make the static time-domain design degenerate.

3. **Separation of Apparent Map and Physical Surface (Major item 3):** I explicitly separated the raw apparent map $\mathbf{m} = (1 - f_c)\mathbf{s} + f_c A_{\mathrm{cl}}\mathbf{1}$ from the physical bare surface $\mathbf{s}$. I formulated the centered residual and affine debiasing step ($\hat{\mathbf{s}} = (\hat{\mathbf{m}} - f_c A_{\mathrm{cl}}\mathbf{1})/(1 - f_c)$), and accounted for the $\Delta f_c \approx 0.008$–0.010 mismatch between specified and realized cloud fractions. In §7(iv), I evaluated the cross-pixel covariance $(\mathbf{C}_{\mathrm{cl}})_{kl}$ and quantified the cost of the pixel-conditional-independence approximation (underestimating formal parameter uncertainties by $\sim$ 15–20% while temporal decorrelation scrambles patterns into an effective noise floor). I elevated surface-correlated clouds ($\beta_{\mathrm{surf}} = 0.25$, $r = 0.1444 \pm 0.0124$, loss $\Delta r = -0.1978$, contrast 0.379) to a headline result in the Abstract, §7, and Conclusion 9 as an irreducible structural systematic that orbital revisits cannot overcome.

4. **Resource-Consistent Cadence Ledgers and Kinematics (Major items 5 and 6):** I implemented two mutually exclusive resource ledgers: Arm A (fixed 8 h per raster position, wall clocks 85.73 to 93.73 d) and Arm B (fixed 89.87 d campaign wall clock, dwells t_s adjusted from 7548.8 s down to 429.6 s at $M_p = 64$, matching the Keplerian tracking integral, Eq. 18). In Table 8, I evaluated the effective independent looks N_{eff} (defined in §5.8–5.9) and showed that what saturates is not r but N_{eff} (12.1 independent looks at $M_p = 64$, $\tau_{\mathrm{cloud}} = 4$ d). I diagnosed and solved the 3-node exposure quadrature aliasing ($R_{36}^{\mathrm{eq3}} = 1$ at $t_s = 7200$ s where sinc vanishes), replacing it with converged 16-node Gauss–Legendre quadrature (error $\leq 2.6 \times 10^{-13}$). In §3.3, I explicitly budgeted the 30-second raster translation kinematics (22.75 km s^{-1} impulsive lower bound for stop-and-dwell over 16,320 transitions), removing operational-feasibility claims for stop-and-dwell and explicitly scoping the raster positions as idealized prescribed samples.

5. **Observable Normalization and Finite-Source Discretization (Major item 7):** In §2.2, I reconciled the disk-mean Lambertian normalization: $\langle \mu \rangle = 8/(3\pi) \simeq 0.848826$, establishing the link between sub-stellar peak SNR (43.16) and disk-mean convolved SNR (36.6344). I specified the central-cell finite-aperture value $K_{00,\mathrm{cellmean}} = 0.042144$ ($n = 64$), and reported the core ablation sensitivity (22% drop in r, 43% in SSIM). In §5.18 and Table 12, I injected coronal systematics in true coronal background units ($\Delta y = \varepsilon Q_{\mathrm{cor}}/Q_{\mathrm{exo}} = 7.74 \times 10^4 \varepsilon$), identifying the single-dwell photon fluctuation as 2.99×10^{-7} and showing that the tolerance knee sits at 3–4σ ($\varepsilon \sim 10^{-6}$). In §7(i), I cited the extended-SGL quadrupole treatment of Turyshev & Toth (2021) and noted that solar quadrupole astigmatic caustics ($W_{J2} \approx 511 \sin^2 \beta_s$ m) require target-specific deconvolution libraries.

6. **Statistical Precision, Bootstrap Intervals, and Seed Disclosure (Major item 8):** I explicitly disclosed in §4.6 that the 10 paired seeds represent weather and photon shot noise variations conditional on a single fixed geography (`TRUTH_SEED = 7`). I defined the $\pm$ convention once as ensemble mean $\pm$ standard error, and provided individual per-seed values and 20,000-resample 95% percentile bootstrap confidence intervals in Appendix A ([Table 14]). I respected the exact sign-flip floor of $2/2^{10} = 1.95 \times 10^{-3}$, replacing all "$p < 10^{-3}$" claims with exact sign-flip and paired t-test p-values. I defined the spherical harmonic metric $r(\ell)$ in §5.3, noted the $\ell \approx 18$ grid ceiling, and constructed a 40-member physical cloud-only null in §5.17 (exceeding all 40 draws, $p \leq 0.024$).

7. **Numerical Validation, Double Precision, and Open Repository (Major item 9):** In §4.5 and Table 3, I reported a float32-vs-float64 precision audit at an honest reduced scale ($n = 32$, 16,384 samples), documenting relative errors in A_λ (3.12×10^{-7}), $\mathbf{b}$ (4.13×10^{-7}), round-off-only contributions (1.02×10^{-8} and 1.47×10^{-8}), normal-equation residual (1.34×10^{-15}), and normal-vs-QR map difference (1.67×10^{-13}), confirming numerical stability. I added a formal *Data and Code Availability* section with the GitHub repository link (https://github.com/sanskarsontakke/sgl-time-domain-imaging) and reproduction CLI commands. A concept DOI will be minted via Zenodo–GitHub integration at the release tag; the version DOI will be recorded in the manuscript upon release.

8. **Target Table Kinematics and Nominal Proxies (Major item 10):** I reduced Table 13 to an explicit nominal-proxy table, stating the scalar bolometric SNR scaling proxy (Eq. 17) explicitly. I corrected the Chen & Kipping radius dispersion to the Terran branch value (4.03%), noting the 2.04 $M_\oplus$ transition and flagging extrapolated rows. I derived the exact geometric de-inclination expectation $\langle(\sin i)^{-0.279}\rangle = 1.0978$ (Eq. 16), separating it from a posterior. I replaced tracking cost with the exact Keplerian time-integral $\Delta v = \oint |\mathbf{a}_\perp(t)|dt$ (Eq. 18), recovering the analytic $2/\pi = 0.6366$ edge-on/face-on ratio. I explicitly flagged τ Cet e as a refuted signal (Figueira et al. 2025), retaining it solely as an illustrative dynamical case.

Point-by-Point Response to the Major Comments of the Pre-Submission Review

Major Item 1: Implemented Novelty and Matched Dynamic Estimator

Review item (internal): *Identify the implemented novelty and compare against the strongest matched dynamic estimator. Use matched data, exposure integration, basis, nuisance knowledge, and tuning for the dynamic comparisons. Report paired contrasts against white/deflated GLS and a covariance-consistent nuisance fit. Give B2 either a faithful matched implementation or an explicitly restricted interpretation. Include a physically bounded-albedo sensitivity test, or explain why the unconstrained solution is the intended benchmark.*
Response: Fully addressed in Introduction (§1), §4.1–4.6, §5.1–5.6, and Table 6:

- *Novelty and Scope:* I have scoped my novelty clearly as an implementation, component testing, and controlled sampling study of an exposure-averaged, time-registered forward operator combined with exact anchored slot-nuisance profiling and covariance-aware inference across two declared resource ledgers.
- *Matched Dynamic Comparisons:* In Table 6 and §5.6, I report factor-by-factor paired contrasts holding data, basis, exposure quadrature, and nuisance knowledge identical. Comparing exact profiling under white noise (Row D: $r = 0.3454 \pm 0.0082$) against unprofiled white-noise GLS (Row A: $r = 0.2886 \pm 0.0097$) reveals that nuisance profiling alone yields $\Delta r = +0.0568 \pm 0.0046$ ($t = 12.4, p_t = 5.9 \times 10^{-7}$). Incorporating OU covariance (Row F: $r = 0.3422 \pm 0.0082$) slightly reduces correlation ($\Delta r = -0.0032 \pm 0.0019$).
- *Restricted Interpretation of B2:* In §4.6 and §5.2, §5.3, §5.6, I explicitly designate baseline B2 as an emulation rather than a reproduction of Turyshev (2026b). I document its severe coverage starvation (256 to 3,584 empty pixels per phase bin depending on arm and M_p), and withdraw B2 from the primary cadence and superiority claims.
- *Bounded-Albedo Benchmark Justification:* In §5.16, I added an explicit paragraph explaining why unconstrained regularized GLS is my primary benchmark: it admits closed-form analytic solutions, exact error propagation, and exact block-diagonal nuisance profiling without non-linear projection artifacts or boundary-clipping dynamics that obscure linear time-domain information limits.

Major Item 2: Spin Recovery, Figure 6b, and Longitude Degeneracy

Review item (internal): *The spin-recovery claim is contradicted by Figure 6b; absolute phase is not identifiable without a longitude reference. Regenerate the objective grid; specify a consistent likelihood and longitude convention; compare phase-perturbed maps before and after registration; and show recovery and interval coverage for identifiable period/pole parameters. Rebuild the static control on common observable support or remove its information-loss interpretation. Withdraw the absolute-phase precursor requirement and in-flight recovery claim.*
Response: Fully addressed in §5.1, §5.8–5.9, Figure 6b, Abstract, and Conclusion 5:

- *Withdrawal of In-Flight Phase Refinement:* I have withdrawn the claim that residual spin misalignments can be refined in flight by minimizing photometric residuals.
- *Analytical and Numerical Invariance:* In §5.9, I demonstrated that the whitened profile likelihood χ^2 is invariant to the assumed initial spin phase (relative spread $\leq 9.2 \times 10^{-12}$ across $\pm 10^\circ$). As derived in Eqs. 10–11 (the profile-likelihood decomposition), an initial phase perturbation δ is absorbed into the map's longitude origin via $F(P, \phi_0 + \delta) R_\delta \mathbf{s} = F(P, \phi_0)\mathbf{s}$, placing the phase derivative in the map-shift tangent space, which profile projection eliminates.
- *Period Error Drift:* I verified the period error drift relation $|\delta\phi| \simeq 2\pi(T/P)|\delta P/P|$ (derived in §5.9). I showed that while grid-search profile likelihood cannot refine spin parameters, the spin state is observable through reconstruction fidelity (r peaks sharply at the true period on 4/4 seeds).
- *Static Control Rebuilt:* In §5.1, I rebuilt the static frozen-planet control on common observable support and identical parameterization using the B3 white-noise GLS estimator ($r = 0.4074 \pm 0.0254$), removing the "virtually zero information loss" claim and explaining that identical pose-passes make the static time-domain design degenerate.

Major Item 3: Cloud Mean, Apparent Map, Covariance, and Surface-Coupled Clouds

Review item (internal): *Derive the cloud mean and measurement covariance before interpreting the fitted map as a surface. Define separate apparent-map and physical-surface variables, the centered residual, the full soft-threshold construction, and the calibration information supplied to the estimator. Measure covariance before and after deflation. Report results for the stated two-timescale model and parameter variations. Include surface-correlated clouds in the abstract and conclusions.*
Response: Fully addressed in §3.2, §4.1, §5.13, §7(iv), Abstract, and Conclusion 9:

- *Apparent Map vs. Physical Surface:* In §4.1, I formally distinguished the raw apparent map $\mathbf{m} = (1 - f_c)\mathbf{s} + f_c A_{\text{cl}} \mathbf{1}$ from the physical bare surface $\mathbf{s}$. I established the centered residual and affine debiasing relation $\hat{\mathbf{s}} = (\hat{\mathbf{m}} - f_c A_{\text{cl}} \mathbf{1})/(1 - f_c)$.
- *Affine Debiasing and Albedo Bias:* I documented that the $\Delta f_c \approx 0.008$–0.010 mismatch between specified cloud fraction (0.55) and realized stochastic cloud cover (0.558) explains the small residual albedo bias ($+0.008$ at 55.8% cover).
- *Cost of Pixel Conditional Independence:* In §7(iv), I evaluated the cross-pixel covariance $(\mathbf{C}_{\text{cl}})_{kl}$ (derived in §7) induced by the $12^\circ \approx 1334$ km cloud correlation scale, showing that neglecting cross-pixel blocks underestimates formal parameter uncertainties by $\sim$ 15–20% while temporal decorrelation scrambles patterns into an effective noise floor.
- *Surface-Correlated Clouds Headline:* I elevated surface-coupled clouds ($\beta_{\text{surf}} = 0.25$, $r = 0.1444 \pm 0.0124$, loss $\Delta r = -0.1978 \pm 0.0175$, contrast 0.379) to the Abstract, §7(iii), and Conclusion 9 as a structural systematic that orbital revisits cannot overcome.

Major Item 4: Slot Deflation, Rank, and Information Separation

Review item (internal): *Propagate slot deflation consistently and separate rank, information, regularization bias, and uncertainty. Verify equivalence to a direct nuisance fit or contrast-space solution. Measure the dominant cloud mode's alignment with the all-ones vector and residual covariance. Replace the information percentage with mode-dependent diagnostics, specify the spherical Laplacian and its null modes, and validate uncertainties.*
Response: Fully addressed in §4.1, §4.2, §5.2, §5.6, Table 2, and Table 6:

- *Exact Anchored Profile Likelihood:* I replaced heuristic slot deflation and the "6.25% dimension reduction preserves 93.75% of information" claim with the exact anchored profile likelihood (Eqs. 10–12), derived in closed form as a rank-one correction to the block-diagonal normal equations:

$$\mathsf{A}_{\text{anchored}} = \mathsf{A}_{\text{free}} + \frac{\mathbf{q}\mathbf{q}^{\mathsf{T}}}{S}, \qquad \mathbf{b}_{\text{anchored}} = \mathbf{b}_{\text{free}} + \frac{\mathbf{q}m}{S}.$$

- *Eigenvalue and Mode Audit:* In Table 2, I reported the complete eigenvalue and mode audit from `results/deflation_diag.json`, documenting unprofiled/free/anchored nullity (288/288/288), zero columns (144), constant mode gain ($913.68 \to 18.79 \to 774.05$), and retained fraction ($2.06\% \to 84.72\%$). I confirmed that the constant surface mode is preserved by the anchored profile.

- *Matched Exact vs. Approximate Contrast:* In §5.6 and Conclusion 4, I report the true matched contrast $F - C$ (holding OU covariance identical), showing that exact profiling differs from the approximate-deflation baseline by only $\Delta r = +(4.1 \pm 3.0) \times 10^{-5}$ ($p_t = 0.20$) and a map-norm difference of $0.83\% \pm 0.22\%$.

Major Item 5: Cadence Trend, Exposure Averaging, and Sinc Attenuation

Review item (internal): *The cadence experiment establishes a trend, not yet the claimed independent-weather mechanism. Publish the exact cadence/time construction and cloud-resampling unit. Use exposure quadrature converged against a substantially finer forward calculation. Measure covariance before and after deflation, include frozen-weather and several-τ_c controls, and vary campaign length independently of visit count.*
Response: Fully addressed in §4.3, §5.8–5.9, Table 1, Table 7, Table 8, Figure 5, Abstract, and Conclusion 3:

- *Effective Looks Saturation:* In Table 8 and §5.9, I evaluated the effective independent looks N_{eff} (defined in §5.8–5.9), demonstrating that what saturates is not r but N_{eff} (12.1 independent looks at $M_p = 64$, $\tau_{\text{cloud}} = 4$ d). I scoped all cadence requirements jointly with τ_{cloud}.

- *Exposure Quadrature Convergence:* In §4.3 and Table 1, I diagnosed the 3-node exposure quadrature aliasing ($R_{36}^{\text{eq3}} = 1$ at $t_s = 7200$ s where sinc vanishes), replacing it with converged 16-node Gauss–Legendre quadrature (error $\leq 2.6 \times 10^{-13}$).

- *Crossed Cadence Controls:* In Table 7, I reported cadence curves across both declared resource ledgers (Arm A and Arm B) with exposure modeling consistently applied, quoting paired standard errors for all configurations.

Major Item 6: Fixed-Resource Accounting and Raster Kinematics

Review item (internal): *The overhead-inclusive schedule violates the fixed-time constraint, and physical raster motion is not budgeted. Rebuild the cadence comparison with one explicit resource ledger and a realizable time list. Either provide a compatible stop-and-dwell trajectory/propulsion budget, use an integrated continuous-scan model, or remove operational-feasibility claims and label the positions as idealized prescribed samples.*
Response: Fully addressed in §3.3, §5.8–5.9, Table 7, and §6.2:

- *Two Consistent Resource Ledgers:* I established two mutually exclusive resource ledgers: Arm A (fixed 8 h per raster position, wall clocks 85.73 to 93.73 d) and Arm B (fixed 89.87 d campaign wall clock, dwells t_s adjusted from 7548.8 s down to 429.6 s at $M_p = 64$, matching the Keplerian tracking integral, Eq. 18).

- *Raster Translation Kinematics Budget:* In §3.3, I budgeted the 30-second raster translation kinematics: rest-to-rest displacement $\Delta = 20.913$ m in 30 s requires $\geq 1.394\ \text{m s}^{-1}$ impulse ($2.788\ \text{m s}^{-1}$ symmetric

bang-bang), summing to an idealized lower bound of 22.75 $\mathrm{km\,s^{-1}}$ per craft over 16,320 transitions. I removed operational-feasibility claims for stop-and-dwell, explicitly scoping the raster positions as idealized prescribed samples.

Major Item 7: Observable Normalization, Finite Discretization, and Systematics

Review item (internal): *The physical observable is more conditional than the quoted SNR suggests; validate the finite-source discretization and systematic budget. State brightness/albedo normalization conventions, central-cell treatment, padding, and boundary conditions. Demonstrate convergence against explicit cell/aperture integration on a reduced problem and isolate the Figure 2 discrepancy. Supply quantitative systematic-injection results, or withdraw that robustness claim. Add the quadrupole-imaging reference.*
Response: Fully addressed in §2.1–2.3, §5.18, Table 12, Figure 2, and §7(i)–(ii):

- *Lambertian Disk Normalization:* In §2.2, I reconciled the disk-mean Lambertian factor $\langle\mu\rangle = 8/(3\pi) \simeq 0.848826$ from `results/validation.npz`, linking the central-peak benchmark ($\mathrm{SNR}_C = 43.16$) to the disk-mean convolved SNR ($\mathrm{SNR}_{C,\text{no-}\mu} = 36.6344$).
- *Central-Cell Discretization:* I specified the central-cell finite-aperture value $K_{00,\text{cellmean}} = 0.042144$ ($n = 64$) from `results/kernel_convergence.json`, and documented the sensitivity from `results/core_ablation.json` (point $r = 0.9803$, SSIM 0.8513 vs. cell-mean $r = 0.7573$, SSIM 0.4856).
- *Figure 2 Discrepancy Resolved:* In §2.3, I stated the theoretical $\min(1.0, \cdot)$ cap and explained the sign reversal across n: discrete pixel averaging attenuates high-wavenumber noise below the continuum integral at coarse $n = 32$ (measured $<$ analytic), while spectral leakage and ill-conditioned transfer function division near Nyquist outpace the integral at fine $n = 64, 128$ (measured $>$ analytic).
- *Coronal Systematics Units:* In §5.18 and Table 12, I injected systematics in true coronal background units ($\Delta y = 7.74 \times 10^4 \varepsilon$), identifying the single-dwell photon fluctuation as 2.99×10^{-7} and showing that the tolerance knee sits at 3–4σ ($\varepsilon \sim 10^{-6}$).
- *Solar Quadrupole:* In §7(i), I cited Turyshev & Toth (2021) and noted that solar quadrupole astigmatic caustics ($W_{J2} \approx 511 \sin^2 \beta_s$ m) require target-specific deconvolution libraries.

Major Item 8: Statistical Precision, Seed Disclosure, and Null Models

Review item (internal): *Statistical precision, significance, and scientific resolution require reproducible definitions and suitable nulls. State exactly which of geography, campaign timing, clouds, and photon noise each seed changes. Give the ten paired values and bootstrap algorithm. Table 2 does not define its ± convention. Re-quote p-values respecting the exact sign-test floor. Define and validate the harmonic metric; report the null exceedance count and construction.*
Response: Fully addressed in §4.6, Tables 4–6, §5.3, §5.17, and Appendix A ([Table 14]):

- *Seed Disclosure:* In §4.6, I stated plainly that all 10 seeds share an identical planetary geography (`TRUTH_SEED = 7`) and vary only weather realizations (seed + 100) and photon shot noise (seed) conditional on that fixed map.
- *Per-Seed Table and Bootstrap Intervals:* In Appendix A ([Table 14]), I reported all 10 individual per-seed values and 20,000-resample 95% percentile bootstrap confidence intervals for key pipeline configurations and paired contrasts.

- *Exact Sign-Test Floors:* I respected the exact two-sided sign-flip floor of $2/2^{10} = 1.95 \times 10^{-3}$, replacing all "$p < 10^{-3}$" claims with exact sign-flip and paired t-test p-values.
- *Harmonic Metric and Nulls:* In §5.3 and §5.17, I defined the spherical harmonic metric $r(\ell)$, noted the $\ell \approx 18$ grid ceiling, and constructed a 40-member physical cloud-only null (exceeding all 40 draws, $p \leq 0.024$).

Major Item 9: Numerical Validation, Double Precision, and Auditable Outputs

Review item (internal): *Numerical validation and robustness claims need auditable outputs, not only summary assertions. Define every precision norm and report errors in both A_λ and* **b***, conditioning/residual diagnostics, and map differences against a full-double-precision reference. Provide a versioned executable archive or sufficiently complete algorithms and numerical inputs. Quantify the stated robustness experiments.*
Response: Fully addressed in §4.2, §4.5, Table 3, §5.13, and Data Availability:

- *Float32 vs. Float64 Precision Audit:* In §4.5 and Table 3, I reported a float32-vs-float64 precision audit at an honest reduced scale ($n = 32$, 16,384 samples), documenting relative errors in A_λ (3.12×10^{-7}), **b** (4.13×10^{-7}), round-off-only contributions (1.02×10^{-8} and 1.47×10^{-8}), normal-equation residual (1.34×10^{-15}), and normal-vs-QR map difference (1.67×10^{-13}), confirming numerical stability.
- *Two-Timescale Process Scored:* In §3.2 and §5.13, I evaluated the two-timescale cloud field with exact variance weighting $(0.3, 0.7)$, reporting $r = 0.3174 \pm 0.0088$ (marginal paired penalty -0.025 ± 0.009).
- *Auditable Repository:* I added a formal *Data and Code Availability* section with my GitHub repository (https://github.com/sanskarsontakke/sgl-time-domain-imaging), replication CLI commands, and complete archive provenance. A concept DOI will be minted via Zenodo–GitHub integration at the release tag.

Major Item 10: Target Uncertainty Table and Nominal Proxies

Review item (internal): *The target uncertainty table is not a reproducible propagation, and its mission interpretation exceeds the tested regime. Either regenerate Table 4 from documented joint input distributions and equations, or retain only explicitly labeled nominal proxies. State composition/branch assumptions, albedo and spectral conventions, interval definitions, orbit inputs, and selection epoch. Remove best-target or architecture conclusions that depend on untested cloud, spin, or trajectory regimes.*
Response: Fully addressed in §6.1, Table 13, and Figure 9:

- *Explicit Nominal Proxies:* I designated Table 13 explicitly as a nominal-proxy ranking, stating the scalar bolometric SNR proxy (Eq. 17) explicitly.
- *Chen & Kipping Branch Discipline:* I corrected the radius dispersion to the Terran branch value (4.03%), noted the 2.04 $M_\oplus$ transition, and flagged extrapolated rows (GJ 273 b, τ Cet e).
- *Geometric De-Inclination Expectation:* I derived the exact geometric expectation for the de-inclination factor, $\langle (\sin i)^{-0.279} \rangle = 1.0978$ (Eq. 16), reporting it separately from the nominal minimum-mass radii.
- *Keplerian Tracking Kinematics:* I replaced the tracking column with the exact Keplerian time-integral $\Delta v = \oint |\mathbf{a}_\perp(t)| dt$ (Eq. 18), recovering the analytic $2/\pi = 0.6366$ edge-on/face-on ratio.
- *Refuted τ Cet e Flagged:* I explicitly disclosed that τ Cet e is classified as a false positive (Figueira et al. 2025), retaining it solely as an illustrative dynamical case.

Point-by-Point Response to the Minor Comments of the Pre-Submission Review

1. **Notation and Dimensions (§2.1, §2.3):** Disconnected target distance z_0 from minimum heliocentric focal distance z. Defined D, Δ, n, N explicitly. Separated surface reflectance, geometric albedo, apparent mean map, and mean cloud opacity.

2. **Noise-Penalty Verbal Definition (§2.3):** Corrected definition to the ratio of reconstructed SNR to convolved input SNR on projected disk support, i.e., the reciprocal of the noise amplification factor σ_R/σ_C.

3. **Latitude Profile (§3.2):** Defined the latitudinal cloud modulation profile and transformation to opacity, verifying realized latitudinal structures.

4. **Earlier Cloud Result (§1):** Corrected Toth (2025) quote to state reconstructed SNR 1.13 at 6.8% cover and 0.80 at 13.7% cover.

5. **Figure Presentation:** Placed a common albedo scale on Figure 3, identified clipping and latitude mask, repositioned Figure 2 annotations outside plotted axes, and defined baseline B3 in §4.6 and Table 4.

6. **Engineering Arithmetic (§6.2):** Corrected solar flux dilution (2.5×10^5 at 500 AU, 4.225×10^5 at 650 AU); identified $v_\infty \simeq 154\ \mathrm{km\,s^{-1}}$ as mean transit scale (650 AU/20 yr); specified sail film areal density $\sigma_{\mathrm{sail}} \approx 1.5$–$2.0\ \mathrm{g\,m^{-2}}$ and $\alpha/\varepsilon \approx 0.1$–$0.2$.

7. **Data Volume and Scaling (§6.2):** Confirmed raw photometry count ($65,536 \times 32 = 2.10$ MB), showing science data is 10.4 kB per map and downlink is not bandwidth-limited.

8. **Architecture-Specific Wording:** Scoped all statements to the chosen single-detector scalar photometric raster architecture.

9. **Conclusions and Physical Interpretation (§8):** Replaced overclaiming words ("proves", "virtually zero information loss") with measured conditional statements; emphasized that persistent surface-locked cloud structure is a demonstrated failure mode.

Thank you for considering the submission. The repository shipped with the manuscript regenerates every number, figure and table quoted above from fixed seeds (`python3 -m reproduce_all --full`), and its test suite re-checks the shipped archives against the text on every push; I am glad to provide anything else a referee may need.

Sincerely,

Sanskar Sontakke
Independent Researcher
sanskarsontakke@gmail.com — ORCID: 0009-0008-4531-5707