

4. There are n kids with candies. You are given an integer array `candies`, where each `candies[i]` represents the number of candies the i th kid has, and an integer `extraCandies`, denoting the number of extra candies that you have. Return a Boolean array `result` of length n , where `result[i]` is `true` if, after giving the i th kid all the `extraCandies`, they will have the greatest number of candies among all the kids, or `false` otherwise.

Note that multiple kids can have the greatest number of candies.

Example 1: Input: `candies = [2,3,5,1,3]`, `extraCandies = 3` Output: `[true,true,true,false,true]` Explanation: If you give all `extraCandies` to: - Kid 1, they will have $2 + 3 = 5$ candies, which is the greatest among the kids. - Kid 2, they will have $3 + 3 = 6$ candies, which is the greatest among the kids. - Kid 3, they will have $5 + 3 = 8$ candies, which is the greatest among the kids. - Kid 4, they will have $1 + 3 = 4$ candies, which is not the greatest among the kids. - Kid 5, they will have $3 + 3 = 6$ candies, which is the greatest among the kids.

Example 2: Input: `candies = [4,2,1,1,2]`, `extraCandies = 1` Output: `[true,false,false,false,false]` Explanation: There is only 1 extra candy. Kid 1 will always have the greatest number of candies, even if a different kid is given the extra candy.

Example 3: Input: `candies = [12,1,12]`, `extraCandies = 10` Output: `[true,false,true]`