

### Chapter 1: Concepts of Motion

displacement:	$\Delta \vec{r} = \vec{r}_f - \vec{r}_i$
Average speed:	$s_{ave} = \frac{\text{total distance}}{\Delta t} = \frac{d}{\Delta t}$
Average velocity:	$\vec{v}_{ave} = \frac{\Delta \vec{r}}{\Delta t}$
Average Acceleration:	$\vec{a}_{ave} = \frac{\Delta \vec{v}}{\Delta t}$

### Chapter 2: Kinematics in one dimension

displacement:	$\Delta x = x_f - x_i$
	$x_f = x_i + \int_{t_i}^{t_f} v_x dx$
Average speed:	$s_{ave} = \frac{\text{total distance}}{\Delta t}$
Instantaneous velocity:	$v = \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} = \frac{dx}{dt}$
Instantaneous acceleration:	$a = \frac{dv}{dt} = \frac{d^2x}{dt^2}$

#### For Constant Acceleration

constant acceleration:	$a = a_{ave} = \frac{v_f - v_i}{\Delta t}$
velocity (with a and t):	$v(t) = v_0 + at$
position (with a and t):	$x(t) = \frac{1}{2}at^2 + v_0t + x_0$
velocity (with a and x):	$v^2 = v_0^2 + 2a(x - x_0)$
position (with v and t):	$x(t) = \frac{1}{2}(v_0 + v)t + x_0$
average velocity:	$v_{ave} = \frac{1}{2}(v_0 + v)$
average velocity:	$v_{ave} = v_0 + \frac{1}{2}at^2$
free fall acceleration:	$a_{freefall} = -g$

### Chapter 3: Vectors and Coordinate Systems

Unit Vectors:	have magnitude (length) = 1
	x-direction, $\hat{i}$
	y-direction, $\hat{j}$
	z-direction, $\hat{k}$
Vector equation:	$\vec{s} = \vec{a} + \vec{b}$
Vector commutative law:	$\vec{a} + \vec{b} = \vec{b} + \vec{a}$
Vector associative law:	$(\vec{a} + \vec{b}) + \vec{c} = \vec{a} + (\vec{b} + \vec{c})$
Vector subtraction:	$\vec{d} = \vec{a} - \vec{b} = \vec{a} + (-\vec{b})$
Vector components:	$\vec{a} = a_x\hat{i} + a_y\hat{j}$
	$a_x = a \cos \theta$ and $a_y = a \sin \theta$
	$a = \sqrt{a_x^2 + a_y^2}$ and $\tan \theta = \frac{a_y}{a_x}$
Vector addition:	$\vec{r} = \vec{a} + \vec{b}$
	$\vec{r} = r_x\hat{i} + r_y\hat{j} + r_z\hat{k}$
	$r_x = a_x + b_x$
	$r_y = a_y + b_y$
	$r_z = a_z + b_z$
vector by scalar multip.:	$2\vec{a} = 2a_x\hat{i} + 2a_y\hat{j} + 2a_z\hat{k}$
Scalar Product:	$\vec{a} \cdot \vec{b} = ab \cos \phi$
	$\vec{a} \cdot \vec{b} = a_xb_x + a_yb_y + a_zb_z$
	$\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$
Vector Product:	$\vec{c} = \vec{a} \times \vec{b}$
	$ c  = ab \sin \phi$
	direction from Right-hand rule
	$\vec{c} = (a_yb_z - a_zb_y)\hat{i} + (a_zb_x - a_xb_z)\hat{j}$
	$+ (a_xb_y - a_yb_x)\hat{k}$
	$\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a})$

Chapter 4: Kinematics in two dimensions	
Position:	$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$
Displacement:	$\Delta\vec{r} = \vec{r}_2 - \vec{r}_1$
	$\Delta\vec{r} = (x_2 - x_1)\hat{i} + (y_2 - y_1)\hat{j} + (z_2 - z_1)\hat{k}$
Average Velocity:	$\vec{v}_{ave} = \frac{\Delta\vec{r}}{\Delta t}$
Instantaneous Velocity:	$\vec{v} = \frac{d\vec{r}}{dt}$
	$\vec{v} = v_x\hat{i} + v_y\hat{j} + v_z\hat{k}$
	$v_x = \frac{dx}{dt}, v_y = \frac{dy}{dt}, v_z = \frac{dz}{dt}$
Average acceleration:	$\vec{a}_{ave} = \frac{\vec{v}_2 - \vec{v}_1}{\Delta t} = \frac{\Delta\vec{v}}{\Delta t}$
Instantaneous acceleration:	$\vec{a} = \frac{d\vec{v}}{dt}$
	$\vec{a} = a_x\hat{i} + a_y\hat{j} + a_z\hat{k}$
	$a_x = \frac{dv_x}{dt}, a_y = \frac{dv_y}{dt}, a_z = \frac{dv_z}{dt}$

Projectile Motion	
Initial velocity:	$v_0 = v_{0x}\hat{i} + v_{0y}\hat{j}$
	$v_{0x} = v_0 \cos \theta$ and $v_{0y} = v_0 \sin \theta$
Horizonatal Motion:	$x(t) = v_{0x}t + x_0$
Vertical Motion:	$y(t) = -\frac{1}{2}gt^2 + v_{0y}t + y_0$
	$v_y = v_{0y}t - gt$
	$v_y^2 = v_{0y}^2 - 2g(y - y_0)$
Equation of the path (trajectory):	$y = (\tan \theta_0)x - \frac{gx^2}{2(v_0 \cos \theta_0)^2}$
Horizontal Range:	$R = \frac{v_0^2}{g} \sin 2\theta_0$

Units	
Length:	1 m = 3.28 ft
	1 mi = 1.61 km
	1 ft = 12 in
Time:	1 min = 60 s
	60 min = 1 hr
	24 hr = 1 day
	365 days = 1 year
Speed:	1 m/s = 3.28 ft/s
	1 km/hr = 0.621 mi/hr

SI Prefixes		
Exponent	Prefix	SI Symbol
$10^{-18}$	atto	a
$10^{-15}$	femto	f
$10^{-12}$	pico	p
$10^{-9}$	nano	n
$10^{-6}$	micro	$\mu$
$10^{-3}$	milli	m
$10^{-2}$	cento	c
$10^{-1}$	deci	d
$10^1$	deca	da
$10^2$	hecto	h
$10^3$	kilo	k
$10^6$	mega	M
$10^9$	giga	G
$10^{12}$	tera	T

<u>Relative Motion:</u> Labels are for a point P in reference frames A,B	
1-dimension:	$x_{PA} = x_{PB} + x_{BA}$
	$v_{PA} = v_{PB} + v_{BA}$
	$a_{PA} = a_{PB}$
2-dimensions:	$\vec{r}_{PA} = \vec{r}_{PB} + \vec{r}_{BA}$
	$\vec{v}_{PA} = \vec{v}_{PB} + \vec{v}_{BA}$
	$\vec{a}_{PA} = \vec{a}_{PB}$
Uniform Circular Motion	
angular position:	$\theta(radians) = \frac{\text{arclength}}{\text{radius}} = \frac{s}{r}$
	$\theta_f = \theta_i + \omega\Delta t$
arclength:	$s = r\theta$
angular velocity:	$\omega_{ave} = \frac{\Delta\theta}{\Delta t}$
	$\omega = \frac{d\theta}{dt}$
centripedal acceleration:	$\vec{a} = \frac{v^2}{r}$ , towardscenterofcircle
Period of revolution:	$T = \frac{2\pi r}{v}$ (Uniform Circular Motion)

Chapter 5: Force and Motion-I	
Newton's 1st law:	if $\vec{F}_{net} = 0$ , then $\vec{a} = 0$
Newton's 2nd law:	$\vec{F}_{net} = m\vec{a}$
	$F_{net,x} = ma_x$
	$F_{net,y} = ma_y$
	$F_{net,z} = ma_z$
Superposition of Forces: (Net force)	$\vec{F}_{net} = \vec{F}_1 + \vec{F}_2 + \vec{F}_3 + \dots$
Gravitational Force:	$F_g = mg$ (magnitude)
Weight:	$W = mg$
Newton's 3rd law: (for two objects, B & C)	$\vec{F}_{BC} = -\vec{F}_{CB}$

Chapter 6,7,8: Forces	
Static Friction:	$f_{s,max} = \mu_s F_N$
Kinetic Friction:	$f_k = \mu_k F_N$
Drag Force:	$D = \frac{1}{2} C \rho A v^2$
Terminal speed:	$v_t = \sqrt{\frac{2F_g}{C\rho A}}$
Centripetal acceleration:	$a = \frac{v^2}{R}$
Centripetal Force:	$F = m \frac{v^2}{R}$

Constants	
Gravitational Acceleration:	$g = 9.8 \frac{m}{s^2}$
Atomic Mass Units:	1 u = $1.66 \times 10^{-27}$ kg

Geometry and other useful equations	
Surface Area:	Volume:
square: $l \times w$	Cube: $l \times w \times h$
circle: $\pi r^2$	Sphere: $\frac{4}{3} \pi r^3$
sphere: $4\pi r^2$	Cylinder: $\pi r^2 h$
cylinder: $2\pi r^2 + 2\pi r h$	
Circumference of a circle: $C = 2\pi r$	
Area of a circle: $A_{circle} = \pi r^2$	
Density: $\rho = \frac{m}{V}$	
Quadratic formula: $(ax^2 + bx + c = 0)$	
$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$	
Angular velocity, $\omega$ : $\omega = \frac{v}{r}$	