Chunk

Santiago Taborga

2025-05-01

```r
df <- read.csv("sample.csv")

# regression estimates
models <-
  list(
    "Masks" = felm(deaths.scaled ~ always.mask
                   + population + svi.index | state, data = df),
    "Vaccination" = felm(deaths.scaled ~ vax.complete
                         + population + svi.index | state, data = df),
    "Masks and Vaccination" = felm(deaths.scaled ~ always.mask + vax.complete
                                   + svi.index | state, data = df)
  )

# regression table
table <- modelsummary(
  models,
  coef_order = c("always.mask", "vax.complete", "population",
                 "svi.indexB", "svi.indexC", "svi.indexD"),
  coef_rename = c(
    "always.mask" = "Mask Usage",
    "vax.complete" = "Vaccine Completion",
    "population" = "Population",
    "svi.indexB" = "SVI - B",
    "svi.indexC" = "SVI - C",
    "svi.indexD" = "SVI - D"
  ),
  gof_map = c('nobs'),
  output = 'gt',
  star = TRUE
)

## add header above the table
table %>%
  tab_spanner(
    label = "Number of deaths",
    columns = c(`Masks`, `Vaccination`, `Masks and Vaccination`)
  )
```

	Number of deaths		
	Masks	Vaccination	Masks and Vaccination
Mask Usage	-112.887***		-83.296***
	(10.212)		(10.834)
Population	-0.000***	-0.000***	
	(0.000)	(0.000)	
SVI - B	11.415***	9.968***	9.581***
	(2.884)	(2.878)	(2.859)
SVI - C	15.142***	14.399***	13.128***
	(3.111)	(3.103)	(3.080)
SVI - D	19.811***	18.353***	17.491***
	(3.401)	(3.393)	(3.367)
Vaccine Completion		-1.123***	-0.923***
		(0.093)	(0.099)
Num.Obs.	3049	3049	3049

+ $p < 0.1$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$