

Original action

Independent duals ψ, χ
Background, source, P
289 fields / 252D matter

Ordered-word branch

P_6 -graded legal Dirac propagator words
Multiply the full word; then evaluate
Bare trace 252 $\neq$ prepared-state value 1

Trace reduction $\neq$ zero open output

One-way scalar trace 0; return -1
Intermediate state projection $\rightarrow 0$
Full $d\Gamma$ / Fock word $\rightarrow$ original readout

Original field branch

289 $\rightarrow$ 103 $\rightarrow$ rational S_{55}
Field / source readback $HF = JS$
At 0: 61D order 2 $\rightarrow$ 5D light kernel

Original state and real time

Full-space P ; double Laplace integral
raw / mean / connected
 S_{01} / dx^1 ; two halves, same source

Original equations $\rightarrow$ variations

Ward source completion $\rightarrow$ two G legs
Current variation $\rightarrow$ reader contact
Duhamel $\rightarrow$ two quantum source legs

Left source leg

Quantum Gram slot 1
Fixed window

Right source leg

Quantum Gram slot 2
Fixed window

Reader contact

Reader variation
Moving intersection

Left propagation

Conjugate G varies
Moving intersection

Right propagation

Field G varies
Moving intersection

Fixed-window source terms + moving-intersection field / contact terms

Body + both fluxes retained; $K = K_G + \frac{1}{2} S_N + K_C$

Retain $|d| \cdot$ full complex Gram $\cdot$ full angular integral + mixed bounds $\rightarrow$ raw / connected: two negative, one positive