

the 1990s, the number of people with a diagnosis of schizophrenia has increased in the United Kingdom (Meltzer et al. 2002). The prevalence of schizophrenia in the United Kingdom is estimated to be 1.2% (Meltzer et al. 2002).

There is a growing body of evidence that people with schizophrenia are at an increased risk of developing a substance use disorder (SUD) (Meltzer et al. 2002). The prevalence of SUD in people with schizophrenia is estimated to be 25% (Meltzer et al. 2002). The risk of developing a SUD is increased in people with schizophrenia who are also taking antipsychotic medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking antidepressant medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking mood stabilizing medication (Meltzer et al. 2002).

The risk of developing a SUD is also increased in people with schizophrenia who are also taking antipsychotic medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking antidepressant medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking mood stabilizing medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking antipsychotic medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking antidepressant medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking mood stabilizing medication (Meltzer et al. 2002).

The risk of developing a SUD is also increased in people with schizophrenia who are also taking antipsychotic medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking antidepressant medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking mood stabilizing medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking antipsychotic medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking antidepressant medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking mood stabilizing medication (Meltzer et al. 2002).

The risk of developing a SUD is also increased in people with schizophrenia who are also taking antipsychotic medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking antidepressant medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking mood stabilizing medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking antipsychotic medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking antidepressant medication (Meltzer et al. 2002). The risk of developing a SUD is also increased in people with schizophrenia who are also taking mood stabilizing medication (Meltzer et al. 2002).